



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 02:49 PM EDT

PDB ID : 1SVP  
Title : SINDBIS VIRUS CAPSID PROTEIN  
Authors : Lee, S.; Rossmann, M.G.  
Deposited on : 1996-03-22  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	 2% 81% 17% ..
1	B	161	 3% 81% 16% ..

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3411 atoms, of which 826 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SINDBIS VIRUS CAPSID PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	160	1506	770	280	217	234	5	0	0	0
1	B	160	1506	770	280	217	234	5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	215	ALA	SER	engineered mutation	UNP P03316
B	215	ALA	SER	engineered mutation	UNP P03316


- Molecule 2 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
2	A	65	195	130	65	0	0
2	B	68	204	136	68	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SINDBIS VIRUS CAPSID PROTEIN

Chain A: 



- Molecule 1: SINDBIS VIRUS CAPSID PROTEIN

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	29.00Å 56.50Å 60.83Å 93.10° 96.73° 94.93°	Depositor
Resolution (Å)	5.00 – 2.00 30.14 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (5.00-2.00) 69.1 (30.14-2.00)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.22 (at 2.00Å)	Xtrriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.184 , 0.279 0.179 , 0.271	Depositor DCC
$R_{free}$ test set	861 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.3	Xtrriage
Anisotropy	0.040	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 61.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3411	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/1254	0.82	1/1693 (0.1%)
1	B	0.50	0/1254	0.80	1/1693 (0.1%)
All	All	0.51	0/2508	0.81	2/3386 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	130	LEU	CA-CB-CG	5.10	127.03	115.30
1	B	130	LEU	CA-CB-CG	5.10	127.02	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1226	280	1199	17	0
1	B	1226	280	1199	23	0
2	A	65	130	0	3	0
2	B	68	136	0	5	0
All	All	2585	826	2398	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:136:VAL:HG22	1:B:169:LEU:HD23	1.53	0.89
1:B:179:THR:HG23	1:B:225:ARG:HG2	1.54	0.88
1:B:219:ILE:HD12	1:B:228:ALA:HB3	1.61	0.81
1:A:156:PHE:HB3	1:A:165:GLU:HG2	1.76	0.66
1:B:179:THR:HG22	2:B:282:HOH:O	1.98	0.63
1:B:248:ASN:ND2	1:B:252:LYS:H	1.97	0.62
1:A:128:HIS:HD2	2:A:270:HOH:O	1.87	0.58
1:A:147:ASP:HB3	2:A:292:HOH:O	2.06	0.56
1:B:252:LYS:HG3	2:B:329:HOH:O	2.05	0.56
1:A:207:ARG:HH21	1:A:237:GLY:HA2	1.71	0.56
1:A:248:ASN:ND2	1:A:252:LYS:H	2.07	0.53
1:B:109:LYS:HD2	1:B:109:LYS:C	2.31	0.50
1:B:128:HIS:HD2	2:B:270:HOH:O	1.95	0.49
1:A:214:ASP:O	1:A:217:ARG:HG3	2.13	0.48
1:B:207:ARG:NH1	1:B:237:GLY:HA2	2.27	0.48
1:B:217:ARG:NH2	2:B:325:HOH:O	2.46	0.48
1:B:207:ARG:HH11	1:B:237:GLY:HA2	1.80	0.47
1:A:136:VAL:HG22	1:A:169:LEU:HD23	1.97	0.47
1:A:110:LEU:HD23	1:A:110:LEU:HA	1.78	0.46
1:B:236:GLU:HB2	1:B:239:ARG:HD2	1.98	0.45
1:A:130:LEU:HD11	1:A:178:PHE:CD2	2.52	0.45
1:A:174:ARG:NH2	2:A:283:HOH:O	2.50	0.44
1:A:135:LYS:HB2	1:A:167:ALA:O	2.16	0.44
1:B:248:ASN:HD21	1:B:252:LYS:H	1.65	0.44
1:B:198:TYR:HA	1:B:202:ARG:O	2.17	0.44
1:B:247:TRP:HA	1:B:252:LYS:O	2.17	0.44
1:B:197:GLN:O	1:B:203:PHE:HA	2.18	0.44
1:A:114:ARG:HH21	1:A:176:GLU:HB2	1.82	0.44
1:B:207:ARG:HD2	1:B:237:GLY:O	2.19	0.43
1:A:115:LEU:HD12	1:A:129:ALA:O	2.19	0.43
1:A:109:LYS:O	1:A:110:LEU:HG	2.19	0.42
1:B:219:ILE:HD12	1:B:228:ALA:CB	2.39	0.42
1:A:156:PHE:CB	1:A:165:GLU:HG2	2.49	0.41
1:B:151:LEU:HG	1:B:173:MET:HE1	2.02	0.41
1:B:109:LYS:HD2	1:B:109:LYS:O	2.20	0.41
1:B:265:SER:HB3	2:B:333:HOH:O	2.20	0.41
1:A:248:ASN:HD22	1:A:248:ASN:C	2.24	0.41
1:B:148:HIS:HB3	1:B:151:LEU:HB2	2.03	0.41
1:B:166:PHE:CD1	1:B:166:PHE:N	2.89	0.40
1:A:197:GLN:NE2	1:A:239:ARG:HH12	2.19	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/161 (98%)	151 (96%)	7 (4%)	0	100	100
1	B	158/161 (98%)	151 (96%)	7 (4%)	0	100	100
All	All	316/322 (98%)	302 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/129 (99%)	121 (94%)	7 (6%)	21	17
1	B	128/129 (99%)	120 (94%)	8 (6%)	18	13
All	All	256/258 (99%)	241 (94%)	15 (6%)	19	15

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	LYS
1	A	155	LYS
1	A	183	GLU
1	A	191	TRP
1	A	207	ARG

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	248	ASN
1	A	253	THR
1	B	109	LYS
1	B	114	ARG
1	B	155	LYS
1	B	172	ASN
1	B	191	TRP
1	B	239	ARG
1	B	248	ASN
1	B	252	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	128	HIS
1	A	197	GLN
1	A	248	ASN
1	B	128	HIS
1	B	172	ASN
1	B	248	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	160/161 (99%)	0.12	4 (2%) 57 56	16, 27, 50, 71	0
1	B	160/161 (99%)	0.06	5 (3%) 49 48	15, 27, 49, 71	0
All	All	320/322 (99%)	0.09	9 (2%) 53 51	15, 27, 50, 71	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	107	ALA	5.7
1	A	266	ALA	5.4
1	A	107	ALA	5.1
1	A	122	ASP	3.4
1	B	186	GLU	3.1
1	B	108	LEU	2.5
1	B	175	SER	2.4
1	A	212	ARG	2.2
1	B	188	PHE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.