



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2023 – 01:24 AM EDT

PDB ID : 5T4C  
Title : Crystal structure of BhGH81 mutant in complex with laminaro-hexaose  
Authors : Pluinage, B.; Boraston, A.B.  
Deposited on : 2016-08-29  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	783	 95%
2	B	6	 33% 67%
3	C	5	 20% 60% 20%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BGC	C	5	X	-	-	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7120 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycoside Hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	752	6128	3892	1034	1186	16	0	13	0

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	7	MET	-	initiating methionine	UNP Q9KG76
A	8	GLY	-	expression tag	UNP Q9KG76
A	9	SER	-	expression tag	UNP Q9KG76
A	10	SER	-	expression tag	UNP Q9KG76
A	11	HIS	-	expression tag	UNP Q9KG76
A	12	HIS	-	expression tag	UNP Q9KG76
A	13	HIS	-	expression tag	UNP Q9KG76
A	14	HIS	-	expression tag	UNP Q9KG76
A	15	HIS	-	expression tag	UNP Q9KG76
A	16	HIS	-	expression tag	UNP Q9KG76
A	17	SER	-	expression tag	UNP Q9KG76
A	18	SER	-	expression tag	UNP Q9KG76
A	19	GLY	-	expression tag	UNP Q9KG76
A	20	LEU	-	expression tag	UNP Q9KG76
A	21	VAL	-	expression tag	UNP Q9KG76
A	22	PRO	-	expression tag	UNP Q9KG76
A	23	ARG	-	expression tag	UNP Q9KG76
A	24	GLY	-	expression tag	UNP Q9KG76
A	25	SER	-	expression tag	UNP Q9KG76
A	26	HIS	-	expression tag	UNP Q9KG76
A	27	MET	-	expression tag	UNP Q9KG76
A	542	GLN	GLU	engineered mutation	UNP Q9KG76

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	6	Total	C	O	0	0	0
			67	36	31			

- Molecule 3 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	5	Total	C	O	0	0	0
			56	30	26			

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O P 5 4 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	771	Total	O	0	0
			771	771		





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.48Å 96.50Å 156.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	82.19 – 1.80 60.87 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (82.19-1.80) 100.0 (60.87-1.80)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.73 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.141 , 0.163 0.154 , 0.174	Depositor DCC
$R_{free}$ test set	4635 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.2	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 47.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7120	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, EDO, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	2/6316 (0.0%)	0.69	1/8608 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	339	GLY	N-CA	11.84	1.63	1.46
1	A	339	GLY	CA-C	-6.00	1.42	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	339	GLY	N-CA-C	-7.84	93.50	113.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	338[A]	ILE	Peptide
1	A	338[B]	ILE	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6128	0	5638	9	0
2	B	67	0	57	0	0
3	C	56	0	48	1	0
4	A	48	0	72	0	0
5	A	50	0	0	0	0
6	A	771	0	0	3	0
All	All	7120	0	5815	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338[B]:ILE:HG22	6:A:905:HOH:O	1.96	0.66
1:A:324[A]:LEU:HG	6:A:905:HOH:O	1.96	0.65
1:A:324[A]:LEU:HD11	1:A:338[A]:ILE:HD12	1.80	0.64
1:A:338[B]:ILE:CG2	6:A:905:HOH:O	2.47	0.61
1:A:324[A]:LEU:HD11	1:A:338[A]:ILE:CD1	2.35	0.57
1:A:107:LEU:HD22	1:A:174:ILE:HG21	1.91	0.53
1:A:748:VAL:CG2	1:A:766:ILE:HD11	2.39	0.52
1:A:338[B]:ILE:HG22	1:A:339:GLY:N	2.34	0.42
1:A:617:ASN:HB2	3:C:5:BGC:C1	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	763/783 (97%)	743 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	633/648 (98%)	631 (100%)	2 (0%)	92	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	364	TYR
1	A	468	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BGC	B	1	2	12,12,12	0.62	0	17,17,17	0.95	1 (5%)
2	BGC	B	2	2	11,11,12	0.39	0	15,15,17	1.02	0
2	BGC	B	3	2	11,11,12	1.10	1 (9%)	15,15,17	1.13	1 (6%)
2	BGC	B	4	2	11,11,12	0.44	0	15,15,17	2.46	7 (46%)
2	BGC	B	5	2	11,11,12	0.40	0	15,15,17	0.88	0
2	BGC	B	6	2	11,11,12	0.68	0	15,15,17	1.26	1 (6%)
3	BGC	C	1	3	12,12,12	0.49	0	17,17,17	0.86	1 (5%)
3	BGC	C	2	3	11,11,12	0.39	0	15,15,17	1.22	1 (6%)
3	BGC	C	3	3	11,11,12	0.39	0	15,15,17	0.68	0
3	BGC	C	4	3	11,11,12	0.41	0	15,15,17	1.38	2 (13%)
3	BGC	C	5	3	11,11,12	0.41	0	15,15,17	2.57	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	1	2	-	1/2/22/22	0/1/1/1
2	BGC	B	2	2	-	0/2/19/22	0/1/1/1
2	BGC	B	3	2	-	0/2/19/22	0/1/1/1
2	BGC	B	4	2	-	0/2/19/22	0/1/1/1
2	BGC	B	5	2	-	0/2/19/22	0/1/1/1
2	BGC	B	6	2	-	0/2/19/22	0/1/1/1
3	BGC	C	1	3	-	0/2/22/22	0/1/1/1
3	BGC	C	2	3	-	0/2/19/22	0/1/1/1
3	BGC	C	3	3	-	0/2/19/22	0/1/1/1
3	BGC	C	4	3	-	0/2/19/22	0/1/1/1
3	BGC	C	5	3	1/1/4/5	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	BGC	O3-C3	-3.45	1.34	1.43

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	5	BGC	O5-C1-C2	7.87	122.92	110.77
3	C	5	BGC	C1-O5-C5	5.59	119.77	112.19
2	B	4	BGC	C1-O5-C5	4.89	118.81	112.19
2	B	4	BGC	C1-C2-C3	4.83	115.60	109.67
3	C	2	BGC	O3-C3-C2	-3.68	102.95	109.99
2	B	4	BGC	O3-C3-C2	-3.67	102.97	109.99
3	C	4	BGC	O3-C3-C2	-3.56	103.18	109.99
3	C	4	BGC	C1-O5-C5	3.55	117.00	112.19
2	B	4	BGC	O5-C1-C2	3.13	115.61	110.77
2	B	4	BGC	O5-C5-C6	2.67	111.39	107.20
2	B	6	BGC	C2-C3-C4	-2.51	106.56	110.89
2	B	4	BGC	C2-C3-C4	2.46	115.15	110.89
3	C	1	BGC	O5-C1-C2	-2.31	106.16	110.28
2	B	1	BGC	C3-C4-C5	2.08	113.95	110.24
2	B	4	BGC	C3-C4-C5	2.06	113.91	110.24
2	B	3	BGC	O3-C3-C2	2.03	113.89	109.99

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	C	5	BGC	C1

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	BGC	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	5	BGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	PO4	A	833	-	4,4,4	0.74	0	6,6,6	0.70	0
5	PO4	A	824	-	4,4,4	1.05	0	6,6,6	1.07	0
5	PO4	A	827	-	4,4,4	1.06	0	6,6,6	0.84	0
4	EDO	A	812	-	3,3,3	0.57	0	2,2,2	0.43	0
4	EDO	A	821	-	3,3,3	0.60	0	2,2,2	0.10	0
5	PO4	A	825	-	4,4,4	1.14	0	6,6,6	0.73	0
4	EDO	A	823	-	3,3,3	0.50	0	2,2,2	0.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	815	-	3,3,3	0.47	0	2,2,2	0.30	0
4	EDO	A	814	-	3,3,3	0.44	0	2,2,2	0.33	0
4	EDO	A	818	-	3,3,3	0.60	0	2,2,2	0.27	0
4	EDO	A	820	-	3,3,3	0.53	0	2,2,2	0.09	0
4	EDO	A	816	-	3,3,3	0.62	0	2,2,2	0.28	0
5	PO4	A	831	-	4,4,4	0.62	0	6,6,6	0.77	0
4	EDO	A	817	-	3,3,3	0.34	0	2,2,2	0.67	0
5	PO4	A	826	-	4,4,4	1.23	0	6,6,6	0.81	0
5	PO4	A	829	-	4,4,4	0.74	0	6,6,6	0.45	0
5	PO4	A	832	-	4,4,4	0.86	0	6,6,6	0.49	0
5	PO4	A	830	-	4,4,4	1.02	0	6,6,6	0.41	0
5	PO4	A	828	-	4,4,4	0.87	0	6,6,6	0.72	0
4	EDO	A	822	-	3,3,3	0.36	0	2,2,2	0.36	0
4	EDO	A	813	-	3,3,3	0.60	0	2,2,2	0.60	0
4	EDO	A	819	-	3,3,3	0.39	0	2,2,2	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	823	-	-	1/1/1/1	-
4	EDO	A	817	-	-	0/1/1/1	-
4	EDO	A	815	-	-	1/1/1/1	-
4	EDO	A	822	-	-	0/1/1/1	-
4	EDO	A	814	-	-	0/1/1/1	-
4	EDO	A	818	-	-	0/1/1/1	-
4	EDO	A	813	-	-	0/1/1/1	-
4	EDO	A	820	-	-	1/1/1/1	-
4	EDO	A	816	-	-	1/1/1/1	-
4	EDO	A	812	-	-	0/1/1/1	-
4	EDO	A	821	-	-	0/1/1/1	-
4	EDO	A	819	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	A	816	EDO	O1-C1-C2-O2
4	A	820	EDO	O1-C1-C2-O2
4	A	815	EDO	O1-C1-C2-O2
4	A	823	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	752/783 (96%)	-0.49	9 (1%) 79 76	7, 13, 24, 47	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	339	GLY	3.8
1	A	778	GLY	3.3
1	A	775	ILE	3.3
1	A	27	MET	3.0
1	A	48	GLY	2.8
1	A	50	ASN	2.6
1	A	766	ILE	2.1
1	A	777	ASN	2.0
1	A	691	TYR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	B	1	12/12	0.83	0.33	22,38,47,47	0
3	BGC	C	5	11/12	0.86	0.18	25,30,35,35	0
3	BGC	C	1	12/12	0.88	0.13	14,18,23,26	0


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BGC	B	6	11/12	0.92	0.12	16,20,23,23	0
3	BGC	C	4	11/12	0.96	0.07	12,15,18,20	0
2	BGC	B	2	11/12	0.96	0.07	11,13,16,16	0
3	BGC	C	3	11/12	0.97	0.08	10,11,11,13	0
3	BGC	C	2	11/12	0.98	0.07	11,12,14,14	0
2	BGC	B	5	11/12	0.98	0.07	8,9,11,12	0
2	BGC	B	3	11/12	0.98	0.06	8,9,10,10	0
2	BGC	B	4	11/12	0.98	0.05	9,9,10,10	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	A	823	4/4	0.66	0.35	42,44,44,44	0
4	EDO	A	817	4/4	0.77	0.19	36,40,41,41	0
4	EDO	A	818	4/4	0.81	0.16	29,29,30,31	0
4	EDO	A	816	4/4	0.82	0.23	30,31,31,31	0
4	EDO	A	820	4/4	0.84	0.16	42,43,44,46	0
4	EDO	A	819	4/4	0.87	0.21	33,36,36,37	0
5	PO4	A	833	5/5	0.87	0.18	33,38,40,40	0
4	EDO	A	821	4/4	0.89	0.15	23,24,24,25	0
4	EDO	A	822	4/4	0.89	0.21	31,31,32,33	0
5	PO4	A	832	5/5	0.90	0.16	39,46,48,48	0
4	EDO	A	815	4/4	0.90	0.16	34,35,35,36	0
5	PO4	A	831	5/5	0.92	0.21	34,34,38,40	0
4	EDO	A	814	4/4	0.92	0.10	27,27,28,29	0
4	EDO	A	813	4/4	0.92	0.13	14,15,16,17	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	PO4	A	829	5/5	0.93	0.17	34,38,39,39	0
5	PO4	A	828	5/5	0.95	0.14	36,38,42,42	0
4	EDO	A	812	4/4	0.96	0.09	16,16,16,17	0
5	PO4	A	830	5/5	0.97	0.12	44,45,45,46	0
5	PO4	A	825	5/5	0.98	0.10	21,21,21,23	0
5	PO4	A	826	5/5	0.98	0.09	21,22,23,23	0
5	PO4	A	827	5/5	0.98	0.07	26,26,27,27	0
5	PO4	A	824	5/5	0.99	0.07	16,17,17,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.