

Full wwPDB X-ray Structure Validation Report (i)

Nov 20, 2023 – 10:47 AM EST

PDB ID	:	8T5Q
Title	:	SARS-CoV-2 ORF3a peptide in complex with TRAF2 TRAF domain
Authors	:	Busscher, B.M.; Xiao, T.S.
Deposited on	:	2023-06-14
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution
IVIETIC	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	Δ	188	5%	70/	110/
1	11	100	4%	/ % •	11%
1	В	188	78%	11% •	11%
1	С	188	75%	14%	11%
1	D	100	13%		
<u> </u>	D	188	74%	14% •	11%
1	Е	188	80%	10%	11%



Mol	Chain	Length	Quality of chain		
1	F	188	9%	17%	11%
2	G	5	80%		20%
2	Н	5	100%		
2	Ι	5	80% 80%		20%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		A	toms			ZeroOcc	AltConf	Trace
1	Δ	168	Total	С	Ν	0	\mathbf{S}	0	0	0
	A	108	1334	853	229	242	10	0	0	0
1	р	168	Total	С	Ν	0	S	0	0	0
	D	108	1334	853	229	242	10	0	0	0
1	C	168	Total	С	Ν	0	S	0	0	0
			1334	853	229	242	10			
1	П	168	Total	С	Ν	0	S	0	0	0
	D	108	1334	853	229	242	10	0	0	0
1	Б	169	Total	С	Ν	0	S	0	0	0
		108	1334	853	229	242	10	0	0	0
1	Б	169	Total	С	Ν	0	S	0	0	0
	Г	108	1334	853	229	242	10	0	U	

• Molecule 1 is a protein called TNF receptor-associated factor 2.

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	314	SER	-	expression tag	UNP Q12933
В	314	SER	-	expression tag	UNP Q12933
С	314	SER	-	expression tag	UNP Q12933
D	314	SER	-	expression tag	UNP Q12933
Е	314	SER	-	expression tag	UNP Q12933
F	314	SER	-	expression tag	UNP Q12933

• Molecule 2 is a protein called ORF3a protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	5	Total 35	C 22	N 6	O 7	0	0	0
2	Н	5	Total 35	C 22	N 6	0 7	0	0	0
2	Ι	5	Total 35	C 22	N 6	0 7	1	0	0



• Molecule 3 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Δ	1	Total C O	0	0
0	Л	1	13 8 5	0	0
2	В	1	Total C O	0	0
5	D	1	13 8 5	0	0
3	Л	1	Total C O	0	0
0	D	1	13 8 5	0	0
3	F	1	Total C O	0	0
J	Г	1	$13 \ 8 \ 5$	0	0

• Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).







Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 7 & 4 & 3 \end{array}$	0	0
4	Ε	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0
4	Ε	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	65	$\begin{array}{cc} \text{Total} & \text{O} \\ 65 & 65 \end{array}$	0	0
5	В	60	Total O 60 60	0	0
5	С	61	$\begin{array}{cc} \text{Total} & \text{O} \\ 61 & 61 \end{array}$	0	0
5	D	27	TotalO2727	0	0
5	Е	67	Total O 67 67	0	0
5	F	34	$\begin{array}{cc} \text{Total} & \text{O} \\ 34 & 34 \end{array}$	0	0
5	G	1	Total O 1 1	0	0
5	Н	1	Total O 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: TNF receptor-associated factor 2



N438 N439 R440 A446 V451 S454 P459 P459 V460	A466 3467 2467 C473 P474 V475 A480 M478 A480 M478 A480 A480 A480 C498 C498 C498 C498 C498 C498 C498 C498	
• Molecule 1: TN	NF receptor-associated factor 2	
Chain E:	80%	10% 11%
SER ALA ALA ALA LEU SER SER SER SER SER CLU CLU	ARG ARG CLY TLE CLY LAS ASP LASA A335 A335 A335 A335 A335 A335 A335 A	M433 R440 R440 E441 P445 P445 P445 P445 R441 R478 R478 R478 R478 R481 R481 R481 R483 R483 R483 R485 R485 R485 R485 R485 R485 R485 R485
• Molecule 1: TN	NF receptor-associated factor 2	
Chain F:	72%	17% 11%
SER GLU ALA ILEU SER SER LVY CLN GLN GLU	ARG CLY TLE CLY CLY CLY CLY CLEU CLY CLEU CLY CLY CLY CLY CLY CLY CLY CLY CLY CLY	A375 M300 M300 M304 M304 M304 M305 M400 M400 M400 M400 M410 M411 M415
R423 N427 N427 N422 M433 N433 N438 R440 R441	V443 1444 1444 1444 V451 V451 P459 P459 R493 K493 K493 L501	
• Molecule 2: Of	RF3a protein	
Chain G:	80%	20%
P204		
• Molecule 2: OI	RF3a protein	
Chain H [.]	100%	
P204 • 1205 • A206 • A207 • S208 •		
• Molecule 2: Of	RF3a protein	
	80%	
Chain I:	80%	20%
P204 1205 A207 S208		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	135.20Å 84.22Å 124.39Å	Depositor
a, b, c, α , β , γ	90.00° 118.82° 90.00°	Depositor
$Perclution(\hat{\lambda})$	29.61 - 1.90	Depositor
Resolution (A)	29.61 - 1.90	EDS
% Data completeness	98.7 (29.61-1.90)	Depositor
(in resolution range)	$98.8\ (29.61-1.90)$	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.56 (at 1.89 Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487, PHENIX 1.20.1_4487	Depositor
D D.	0.207 , 0.240	Depositor
Π, Π_{free}	0.206 , 0.238	DCC
R_{free} test set	4759 reflections $(4.98%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	29.6	Xtriage
Anisotropy	0.448	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 45.4	EDS
L-test for twinning ²	$< L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8505	wwPDB-VP
Average B, all atoms $(Å^2)$	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 29.28 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5917e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PG4 $\,$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.41	0/1364	0.63	0/1842
1	В	0.41	0/1364	0.66	0/1842
1	С	0.41	0/1364	0.62	0/1842
1	D	0.36	0/1364	0.60	0/1842
1	Ε	0.45	0/1364	0.65	0/1842
1	F	0.37	0/1364	0.64	0/1842
2	G	0.32	0/35	0.47	0/46
2	Н	0.25	0/35	0.40	0/46
2	Ι	0.25	0/35	0.35	0/46
All	All	0.40	0/8289	0.63	0/11190

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Ε	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	Ε	365	ARG	Sidechain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1334	0	1328	8	0
1	В	1334	0	1328	16	0
1	С	1334	0	1328	23	0
1	D	1334	0	1328	20	0
1	Е	1334	0	1328	7	0
1	F	1334	0	1328	19	0
2	G	35	0	36	1	0
2	Н	35	0	36	0	0
2	Ι	35	0	36	1	0
3	А	13	0	18	1	0
3	В	13	0	18	4	0
3	D	13	0	18	0	0
3	F	13	0	18	1	0
4	С	14	0	20	3	0
4	Ε	14	0	20	0	0
5	А	65	0	0	0	0
5	В	60	0	0	0	0
5	С	61	0	0	0	0
5	D	27	0	0	0	0
5	Е	67	0	0	0	0
5	F	34	0	0	0	0
5	G	1	0	0	0	0
5	Н	1	0	0	0	0
All	All	8505	0	8188	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:436:ASP:H	1:C:442:HIS:HD2	1.24	0.85
1:C:482:ASN:HD21	1:C:485:VAL:H	1.32	0.78
1:B:493:LYS:HZ2	3:B:601:PG4:H61	1.50	0.76
1:B:486:ARG:HG3	1:B:486:ARG:HH11	1.53	0.73



			Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:C:479:GLU:H	1:C:482:ASN:HB3	1.57	0.68
1:D:338:LEU:HD21	1:F:338:LEU:HB3	1.77	0.66
1:D:338:LEU:HD13	1:F:339:GLU:HG2	1.79	0.65
1:E:393:ARG:NH1	1:E:395:TYR:OH	2.32	0.63
1:F:367:GLU:HG2	1:F:372:ARG:HH12	1.64	0.62
1:F:341:LYS:O	1:F:345:MET:HG2	2.00	0.62
1:D:338:LEU:O	1:D:342:VAL:HG23	2.01	0.61
1:E:479:GLU:OE2	1:E:481:LYS:HE3	2.00	0.60
1:B:341:LYS:O	1:B:345:MET:HG3	2.01	0.59
1:A:353:VAL:HG11	3:A:601:PG4:H21	1.84	0.59
1:F:334:ALA:N	1:F:337:ASP:OD2	2.35	0.59
1:D:395:TYR:HB2	1:D:408:SER:HB2	1.85	0.58
1:A:345:MET:HE1	1:C:342:VAL:HG13	1.85	0.58
1:C:441:GLU:OE1	1:C:477:LYS:NZ	2.36	0.57
1:C:400:GLY:H	1:C:403:ARG:NH1	2.03	0.57
1:F:395:TYR:HB2	1:F:408:SER:HB2	1.89	0.55
1:C:482:ASN:ND2	1:C:485:VAL:H	2.01	0.55
1:B:434:LEU:HD21	1:B:478:MET:HE1	1.88	0.55
1:D:394:ILE:HD11	1:D:407:LEU:HD21	1.90	0.54
1:C:436:ASP:N	1:C:442:HIS:HD2	2.01	0.54
1:D:390:MET:HG2	1:D:411:PHE:CZ	2.42	0.54
1:C:478:MET:HA	1:C:482:ASN:HD22	1.73	0.54
1:D:479:GLU:HG3	1:D:480:ALA:N	2.23	0.54
1:C:482:ASN:HD21	1:C:485:VAL:N	2.04	0.52
1:D:345:MET:HE1	1:F:345:MET:HG3	1.90	0.52
1:B:493:LYS:NZ	3:B:601:PG4:H61	2.20	0.52
1:B:486:ARG:HG3	1:B:486:ARG:NH1	2.24	0.51
1:D:479:GLU:HG2	1:D:481:LYS:H	1.75	0.51
1:C:396:LEU:HA	1:C:407:LEU:HD12	1.92	0.50
1:C:396:LEU:HD23	1:C:407:LEU:HD11	1.93	0.50
1:C:407:LEU:HD13	1:C:478:MET:CE	2.42	0.50
1:D:407:LEU:N	1:D:473:CYS:O	2.44	0.50
1:B:340:GLN:O	1:B:344:GLU:HG3	2.12	0.49
1:D:415:LYS:HB2	1:D:459:PRO:HB2	1.94	0.49
1:E:425:PRO:HG2	1:E:451:VAL:HG22	1.94	0.49
1:F:415:LYS:HA	1:F:459:PRO:HG2	1.93	0.48
1:F:372:ARG:NH1	1:F:372:ARG:HB3	2.28	0.48
1:D:467:SER:HA	2:I:206:GLN:HA	1.96	0.48
1:C:478:MET:HA	1:C:482:ASN:ND2	2.29	0.47
1:E:361:PHE:HZ	1:E:478:MET:HG3	1.79	0.47
1:D:407:LEU:HD12	1:D:478:MET:SD	2.55	0.47



	lo de page	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:B:394:ILE:HD11	1:B:407:LEU:HD21	1.95	0.47	
1:C:360:ASP:OD2	1:C:363:ARG:NH2	2.24	0.47	
1:E:493:LYS:HE2	1:E:495:ILE:HD11	1.97	0.47	
1:F:335:MET:O	1:F:339:GLU:HG3	2.13	0.47	
1:B:477:LYS:HD3	1:B:484:TYR:CE2	2.50	0.46	
1:C:493:LYS:HD3	4:C:602:PEG:H41	1.96	0.46	
1:F:394:ILE:HD12	1:F:409:LEU:HD13	1.98	0.46	
1:C:493:LYS:HZ2	4:C:602:PEG:H41	1.80	0.46	
1:F:427:ASN:HA	1:F:449:PRO:HG2	1.98	0.45	
1:F:390:MET:HG2	1:F:411:PHE:CZ	2.51	0.45	
1:D:440:ARG:HD3	1:D:440:ARG:HA	1.59	0.45	
1:B:493:LYS:HZ2	3:B:601:PG4:H42	1.82	0.44	
1:A:346:GLU:HG2	1:B:345:MET:SD	2.58	0.44	
1:B:493:LYS:HZ2	3:B:601:PG4:C6	2.26	0.44	
1:F:493:LYS:HD3	3:F:601:PG4:H22	1.99	0.44	
1:E:336:ALA:HA	1:E:339:GLU:OE2	2.18	0.44	
1:E:338:LEU:HD12	1:E:338:LEU:HA	1.75	0.44	
1:A:467:SER:OG	2:G:204:PRO:HB2	2.17	0.43	
1:B:367:GLU:CD	1:B:373:ILE:HD13	2.39	0.43	
1:D:365:ARG:NH2	1:D:475:VAL:HG11	2.34	0.42	
1:D:412:VAL:HG22	1:D:466:ALA:HB2	2.01	0.42	
1:D:497:ASP:OD2	1:D:499:THR:OG1	2.20	0.42	
1:D:345:MET:SD	1:F:346:GLU:HG3	2.60	0.42	
1:A:382:TYR:HA	1:A:388:TYR:O	2.20	0.42	
1:C:493:LYS:NZ	4:C:601:PEG:O2	2.52	0.42	
1:F:396:LEU:HA	1:F:407:LEU:CD2	2.50	0.42	
1:B:485:VAL:HG22	1:B:490:ILE:CG2	2.50	0.41	
1:A:338:LEU:HD21	1:C:338:LEU:HD23	2.01	0.41	
1:C:433:MET:HA	1:C:443:VAL:O	2.20	0.41	
1:C:477:LYS:O	1:C:482:ASN:HB2	2.21	0.41	
1:A:434:LEU:HB3	1:A:443:VAL:HB	2.03	0.41	
1:C:382:TYR:HA	1:C:388:TYR:O	2.21	0.41	
1:C:454:SER:HA	1:C:457:GLN:HG2	2.03	0.41	
1:D:338:LEU:HD12	1:D:338:LEU:HA	1.76	0.41	
1:F:375:ALA:HB2	1:F:395:TYR:CE1	2.55	0.41	
1:F:390:MET:HE3	1:F:390:MET:HB2	2.01	0.41	
1:F:433:MET:HA	1:F:443:VAL:O	2.21	0.41	
1:A:412:VAL:HG22	1:A:466:ALA:HB2	2.03	0.41	
1:B:433:MET:HA	1:B:443:VAL:O	2.20	0.41	
1:B:454:SER:HA	1:B:457:GLN:HG2	2.02	0.41	
1:D:479:GLU:HG3	1:D:480:ALA:H	1.86	0.41	



There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	166/188~(88%)	163~(98%)	3~(2%)	0	100	100
1	В	166/188~(88%)	164 (99%)	2(1%)	0	100	100
1	С	166/188~(88%)	162 (98%)	4 (2%)	0	100	100
1	D	166/188~(88%)	165 (99%)	1 (1%)	0	100	100
1	Ε	166/188~(88%)	164 (99%)	2(1%)	0	100	100
1	F	166/188~(88%)	163 (98%)	3(2%)	0	100	100
2	G	3/5~(60%)	3 (100%)	0	0	100	100
2	Н	3/5~(60%)	3 (100%)	0	0	100	100
2	Ι	3/5~(60%)	3 (100%)	0	0	100	100
All	All	1005/1143 (88%)	990 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	144/162~(89%)	140~(97%)	4 (3%)	43 36
1	В	144/162~(89%)	142~(99%)	2(1%)	67 65
1	С	144/162~(89%)	140 (97%)	4 (3%)	43 36



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	D	144/162~(89%)	139~(96%)	5(4%)	36	27
1	Ε	144/162~(89%)	140~(97%)	4(3%)	43	36
1	F	144/162~(89%)	138 (96%)	6 (4%)	30	20
2	G	4/4 (100%)	4 (100%)	0	100	100
2	Н	4/4 (100%)	4 (100%)	0	100	100
2	Ι	4/4 (100%)	4 (100%)	0	100	100
All	All	876/984~(89%)	851 (97%)	25 (3%)	42	35

Continued from previous page...

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	335	MET
1	А	345	MET
1	А	433	MET
1	А	481	LYS
1	В	340	GLN
1	В	414	MET
1	С	335	MET
1	С	337	ASP
1	С	445	ASP
1	С	486	ARG
1	D	335	MET
1	D	423	ARG
1	D	438	ASN
1	D	478	MET
1	D	482	ASN
1	Е	335	MET
1	Е	403	ARG
1	Е	433	MET
1	Е	445	ASP
1	F	365	ARG
1	F	403	ARG
1	F	423	ARG
1	F	438	ASN
1	F	441	GLU
1	F	454	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:



Mol	Chain	Res	Type
1	С	442	HIS
1	С	482	ASN
1	D	439	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	ain Dec Link		Bog Link Bond lengths			Bond angles		
	mor Type Cha	Ullaili	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PG4	В	601	-	12,12,12	0.23	0	11,11,11	0.39	0
4	PEG	Е	602	-	$6,\!6,\!6$	0.14	0	$5,\!5,\!5$	0.17	0
4	PEG	С	602	-	$6,\!6,\!6$	0.17	0	$5,\!5,\!5$	0.18	0
3	PG4	F	601	-	12,12,12	0.25	0	11,11,11	0.37	0
4	PEG	С	601	-	$6,\!6,\!6$	0.20	0	$5,\!5,\!5$	0.08	0
4	PEG	Е	601	-	$6,\!6,\!6$	0.09	0	$5,\!5,\!5$	0.22	0
3	PG4	D	601	-	12,12,12	0.20	0	11,11,11	0.47	0
3	PG4	А	601	-	12,12,12	0.20	0	11,11,11	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	В	601	-	-	9/10/10/10	-
4	PEG	Е	602	-	-	2/4/4/4	-
4	PEG	С	602	-	-	2/4/4/4	-
3	PG4	F	601	-	-	5/10/10/10	-
4	PEG	С	601	-	-	0/4/4/4	-
4	PEG	Е	601	-	-	2/4/4/4	-
3	PG4	D	601	-	-	3/10/10/10	-
3	PG4	А	601	-	-	1/10/10/10	-

'-' means no outliers of that kind were identified.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	601	PG4	O2-C3-C4-O3
3	В	601	PG4	O1-C1-C2-O2
4	С	602	PEG	O1-C1-C2-O2
3	В	601	PG4	C8-C7-O4-C6
4	Е	601	PEG	O1-C1-C2-O2
3	D	601	PG4	O2-C3-C4-O3
3	В	601	PG4	O4-C7-C8-O5
4	Е	601	PEG	O2-C3-C4-O4
4	Е	602	PEG	O1-C1-C2-O2
3	В	601	PG4	C1-C2-O2-C3
3	В	601	PG4	C3-C4-O3-C5
3	F	601	PG4	C1-C2-O2-C3
3	F	601	PG4	C4-C3-O2-C2
3	В	601	PG4	C5-C6-O4-C7
3	D	601	PG4	O1-C1-C2-O2
4	С	602	PEG	C1-C2-O2-C3
3	D	601	PG4	C4-C3-O2-C2
3	В	601	PG4	C4-C3-O2-C2
3	F	601	PG4	C8-C7-O4-C6
3	F	601	PG4	C5-C6-O4-C7
3	В	601	PG4	O3-C5-C6-O4
4	Е	602	PEG	C4-C3-O2-C2
3	А	601	PG4	O3-C5-C6-O4
3	В	601	PG4	O2-C3-C4-O3



There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	601	PG4	4	0
4	С	602	PEG	2	0
3	F	601	PG4	1	0
4	С	601	PEG	1	0
3	А	601	PG4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	168/188~(89%)	0.32	9 (5%) 25 29	26, 35, 63, 86	7 (4%)
1	В	168/188~(89%)	0.30	7 (4%) 36 39	25, 36, 67, 95	6 (3%)
1	С	168/188~(89%)	0.57	19 (11%) 5 5	26, 36, 80, 123	3 (1%)
1	D	168/188~(89%)	0.86	24 (14%) 2 2	32, 50, 95, 128	1 (0%)
1	Е	168/188~(89%)	0.24	11 (6%) 18 21	24, 32, 70, 104	3 (1%)
1	F	168/188~(89%)	0.69	17 (10%) 7 8	28, 49, 75, 118	3 (1%)
2	G	5/5~(100%)	1.15	1 (20%) 1 1	39, 40, 44, 53	3 (60%)
2	Н	5/5~(100%)	4.18	5 (100%) 0 0	60, 66, 73, 74	3 (60%)
2	Ι	5/5~(100%)	3.05	4 (80%) 0 0	60, 64, 71, 91	2 (40%)
All	All	1023/1143~(89%)	0.53	97 (9%) 8 9	24, 38, 80, 128	31 (3%)

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	334	ALA	9.5
1	D	334	ALA	9.5
1	С	480	ALA	8.2
1	D	480	ALA	7.8
1	С	479	GLU	7.4
1	Е	335	MET	6.9
2	Н	208	SER	6.7
1	А	336	ALA	6.5
1	А	480	ALA	6.3
1	С	439	ASN	6.3
1	D	338	LEU	6.1
1	В	480	ALA	5.9
1	D	335	MET	5.8
1	D	336	ALA	5.7
1	D	481	LYS	5.6



0100

Mol	Chain	Res	Type	RSRZ
1	С	440	ARG	5.5
1	D	337	ASP	5.3
1	Е	336	ALA	5.1
1	А	335	MET	5.0
1	F	335	MET	4.8
1	D	479	GLU	4.6
2	Ι	207	ALA	4.5
1	Е	337	ASP	4.5
1	В	481	LYS	4.3
1	С	336	ALA	4.3
1	F	451	VAL	4.2
1	С	335	MET	4.2
1	В	335	MET	4.2
1	С	476	SER	4.2
2	Н	207	ALA	4.1
1	С	481	LYS	4.1
1	F	336	ALA	4.1
1	А	481	LYS	4.0
2	Н	205	ILE	4.0
1	F	338	LEU	4.0
1	D	372	ARG	4.0
1	F	439	ASN	3.9
2	Ι	205	ILE	3.8
1	А	334	ALA	3.7
1	В	334	ALA	3.7
2	Н	206	GLN	3.3
1	С	482	ASN	3.3
2	Ι	208	SER	3.2
1	С	483	SER	3.2
1	F	339	GLU	3.2
2	Ι	206	GLN	3.2
1	D	343	LEU	3.1
1	Е	338	LEU	3.1
1	F	432	LEU	3.0
1	D	451	VAL	2.9
1	D	501	LEU	2.9
1	D	369	VAL	2.9
1	F	401	THR	2.8
1	А	337	ASP	2.8
2	Н	204	PRO	2.7
2	G	208	SER	2.6
1	В	482	ASN	2.6



Mol	Chain	Res	Type	RSRZ
1	С	438	ASN	2.6
1	F	476	SER	2.5
1	D	339	GLU	2.5
1	F	438	ASN	2.5
1	А	476	SER	2.5
1	С	441	GLU	2.5
1	Е	441	GLU	2.5
1	D	392	LEU	2.5
1	D	340	GLN	2.5
1	Е	339	GLU	2.5
1	С	452	THR	2.4
1	D	460	VAL	2.4
1	Е	440	ARG	2.4
1	F	343	LEU	2.3
1	С	445	ASP	2.3
1	D	482	ASN	2.3
1	С	475	VAL	2.3
1	D	405	THR	2.2
1	F	440	ARG	2.2
1	D	446	ALA	2.2
1	В	430	VAL	2.2
1	Е	444	ILE	2.1
1	А	432	LEU	2.1
1	Е	439	ASN	2.1
1	F	400	GLY	2.1
1	В	460	VAL	2.1
1	Е	341	LYS	2.1
1	А	440	ARG	2.1
1	С	477	LYS	2.1
1	С	343	LEU	2.1
1	D	454	SER	2.1
1	F	441	GLU	2.1
1	F	423	ARG	2.1
1	C	446	ALA	2.1
1	С	405	THR	2.1
1	D	409	LEU	2.1
1	D	366	GLN	2.0
1	F	444	ILE	2.0
1	F	462	ASP	2.0
1	D	344	GLU	2.0

Continued from previous page...



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q<0.9
4	PEG	С	601	7/7	0.75	0.14	42,46,49,49	0
3	PG4	А	601	13/13	0.80	0.19	36,39,42,44	13
4	PEG	Е	602	7/7	0.80	0.19	43,44,46,50	7
4	PEG	Е	601	7/7	0.82	0.14	43,45,45,47	0
3	PG4	D	601	13/13	0.82	0.20	40,43,46,48	13
3	PG4	F	601	13/13	0.84	0.14	43,45,48,51	0
3	PG4	В	601	13/13	0.88	0.13	42,45,49,50	0
4	PEG	С	602	7/7	0.91	0.12	44,47,49,54	0

6.5 Other polymers (i)

There are no such residues in this entry.

