



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 30, 2023 – 06:03 PM EST

PDB ID : 8T9Q
Title : Crystal structure of fused YR, an asymmetric 4-OT trimer
Authors : Moreno, R.Y.; Zhang, Y.J.
Deposited on : 2023-06-24
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	25% (Poor fit) 40% (≥3 outliers), 30% (2 outliers), 28% (1 outlier), 2% (0 outliers), 2% (Not modelled)
1	B	149	27% (Poor fit) 48% (≥3 outliers), 27% (2 outliers), 22% (1 outlier), 2% (0 outliers), 2% (Not modelled)
1	C	149	24% (Poor fit) 53% (≥3 outliers), 24% (2 outliers), 19% (1 outlier), 2% (0 outliers), 2% (Not modelled)

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 2552 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tautomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	108	782	495	128	158	1	0	0	0
1	B	116	848	539	140	168	1	0	0	0
1	C	121	890	565	146	178	1	0	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	66	SER	-	linker	UNP J2VT77
A	67	ALA	-	linker	UNP J2VT77
A	68	ALA	-	linker	UNP J2VT77
A	69	ASP	-	linker	UNP J2VT77
A	70	GLY	-	linker	UNP J2VT77
A	71	ALA	-	linker	UNP J2VT77
A	72	PRO	-	linker	UNP J2VT77
A	73	PRO	-	linker	UNP J2VT77
A	74	SER	-	linker	UNP J2VT77
A	75	LEU	-	linker	UNP J2VT77
B	66	SER	-	linker	UNP J2VT77
B	67	ALA	-	linker	UNP J2VT77
B	68	ALA	-	linker	UNP J2VT77
B	69	ASP	-	linker	UNP J2VT77
B	70	GLY	-	linker	UNP J2VT77
B	71	ALA	-	linker	UNP J2VT77
B	72	PRO	-	linker	UNP J2VT77
B	73	PRO	-	linker	UNP J2VT77
B	74	SER	-	linker	UNP J2VT77
B	75	LEU	-	linker	UNP J2VT77
C	66	SER	-	linker	UNP J2VT77
C	67	ALA	-	linker	UNP J2VT77
C	68	ALA	-	linker	UNP J2VT77

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	69	ASP	-	linker	UNP J2VT77
C	70	GLY	-	linker	UNP J2VT77
C	71	ALA	-	linker	UNP J2VT77
C	72	PRO	-	linker	UNP J2VT77
C	73	PRO	-	linker	UNP J2VT77
C	74	SER	-	linker	UNP J2VT77
C	75	LEU	-	linker	UNP J2VT77


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	13	Total O 13 13	0	0
2	B	12	Total O 12 12	0	0
2	C	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tautomerase



- Molecule 1: Tautomerase



- Molecule 1: Tautomerase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.09Å 49.41Å 49.32Å 102.36° 102.16° 102.11°	Depositor
Resolution (Å)	46.51 – 2.29 46.42 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.6 (46.51-2.29) 94.6 (46.42-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.32Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.269 , 0.303 0.266 , 0.299	Depositor DCC
R_{free} test set	1824 reflections (10.09%)	wwPDB-VP
Wilson B-factor (Å ²)	60.7	Xtrriage
Anisotropy	0.136	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 79.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.398 for l,h,k 0.398 for k,l,h 0.410 for -l,-k,-h 0.400 for -k,-h,-l 0.447 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2552	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/791	0.87	0/1078
1	B	0.71	0/857	0.89	0/1167
1	C	0.73	0/901	0.88	0/1229
All	All	0.71	0/2549	0.88	0/3474

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	136	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	782	0	787	51	0
1	B	848	0	868	49	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	890	0	901	32	0
2	A	13	0	0	5	0
2	B	12	0	0	1	0
2	C	7	0	0	2	0
All	All	2552	0	2556	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:GLU:CD	1:B:106:THR:HG21	1.52	1.26
1:A:137:GLU:OE1	1:B:106:THR:HG21	1.27	1.26
1:A:137:GLU:OE1	1:B:106:THR:CG2	1.97	1.11
1:A:39:ARG:NH1	1:B:129:ASP:OD2	1.99	0.96
1:B:71:ALA:HA	1:B:135:ARG:NH2	1.81	0.95
1:B:133:ILE:O	1:B:135:ARG:NH2	2.04	0.90
1:A:137:GLU:OE2	1:B:106:THR:HG21	1.77	0.85
1:A:114:LEU:HD22	1:C:144:ASP:HA	1.58	0.85
1:A:39:ARG:CZ	1:B:129:ASP:OD2	2.29	0.80
1:C:72:PRO:HD2	1:C:133:ILE:O	1.80	0.80
1:A:106:THR:HG21	2:C:201:HOH:O	1.85	0.76
1:A:60:ILE:C	2:A:207:HOH:O	2.24	0.76
1:A:5:MET:SD	1:A:19:LEU:HD21	2.26	0.75
1:B:23:LEU:O	1:B:27:THR:OG1	2.06	0.73
1:B:133:ILE:HB	1:B:135:ARG:NH2	2.04	0.72
1:B:71:ALA:HA	1:B:135:ARG:HH22	1.55	0.72
1:B:2:ASN:HB2	1:B:71:ALA:HB3	1.73	0.71
1:C:139:ARG:NH2	2:C:201:HOH:O	2.26	0.68
1:A:72:PRO:HD2	1:A:133:ILE:O	1.94	0.68
1:A:132:GLY:HA3	2:A:211:HOH:O	1.94	0.67
1:B:65:GLU:HG3	1:B:127:ALA:O	1.95	0.67
1:A:23:LEU:O	1:A:27:THR:OG1	2.14	0.66
1:A:4:GLU:HG3	1:A:69:ASP:HB2	1.78	0.64
1:A:35:ILE:O	1:B:53:GLY:HA2	1.98	0.64
1:A:39:ARG:NH2	1:B:129:ASP:OD2	2.31	0.63
1:A:103:ILE:O	1:A:106:THR:OG1	2.11	0.62
1:B:72:PRO:HD2	1:B:133:ILE:O	1.98	0.62
1:A:4:GLU:CG	1:A:69:ASP:HB2	2.31	0.61
1:A:18:ALA:HB1	1:A:120:ILE:HG12	1.84	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:SER:H	1:A:136:ALA:HB2	1.67	0.60
1:A:39:ARG:HH12	1:B:129:ASP:CG	2.04	0.59
1:B:22:ALA:HB1	1:B:116:LYS:HB3	1.84	0.58
1:A:65:GLU:N	2:A:202:HOH:O	2.37	0.57
1:B:5:MET:SD	1:B:19:LEU:HD21	2.45	0.57
1:C:18:ALA:CB	1:C:120:ILE:HG12	2.34	0.57
1:A:102:ILE:O	1:A:106:THR:HG23	2.04	0.57
1:A:103:ILE:HG23	1:A:107:PHE:HE2	1.69	0.56
1:A:103:ILE:HG23	1:A:107:PHE:CE2	2.40	0.56
1:A:60:ILE:O	2:A:201:HOH:O	2.17	0.56
1:A:131:VAL:HG13	1:C:135:ARG:HG2	1.88	0.56
1:A:18:ALA:CB	1:A:120:ILE:HG12	2.34	0.56
1:C:101:SER:HB3	1:C:105:VAL:HG23	1.88	0.55
1:A:50:PHE:O	1:A:57:VAL:HG23	2.07	0.55
1:A:30:ALA:HB1	1:A:108:ILE:HG22	1.89	0.54
1:A:30:ALA:HB2	1:A:112:THR:HG21	1.89	0.54
1:B:144:ASP:HA	1:C:114:LEU:HD22	1.87	0.54
1:B:141:ILE:HB	1:C:130:THR:HB	1.90	0.54
1:C:18:ALA:HB1	1:C:120:ILE:HG12	1.90	0.54
1:A:114:LEU:HD22	1:C:144:ASP:CA	2.34	0.53
1:B:30:ALA:HB2	1:B:112:THR:HG21	1.90	0.53
1:B:143:LYS:C	1:C:114:LEU:HD22	2.28	0.53
1:A:16:LYS:NZ	1:A:44:GLU:OE2	2.34	0.52
1:B:35:ILE:O	1:C:53:GLY:HA2	2.09	0.52
1:C:30:ALA:HB1	1:C:108:ILE:HG22	1.92	0.51
1:A:137:GLU:OE2	1:B:106:THR:CG2	2.56	0.51
1:C:114:LEU:HD12	1:C:128:VAL:HG22	1.92	0.50
1:B:15:GLN:OE1	1:B:121:ALA:HA	2.13	0.49
1:A:137:GLU:OE1	1:B:106:THR:HG22	2.02	0.49
1:B:13:ASP:O	1:B:16:LYS:HB2	2.12	0.49
1:B:133:ILE:HB	1:B:135:ARG:HH22	1.77	0.49
1:B:48:THR:OG1	2:B:201:HOH:O	2.20	0.48
1:B:108:ILE:O	1:B:112:THR:HG23	2.14	0.47
1:C:4:GLU:HG3	1:C:69:ASP:HB2	1.97	0.47
1:A:132:GLY:CA	2:A:211:HOH:O	2.58	0.47
1:B:144:ASP:N	1:C:114:LEU:HD22	2.29	0.47
1:A:53:GLY:HA2	1:C:35:ILE:O	2.15	0.47
1:C:71:ALA:HB1	1:C:135:ARG:HE	1.80	0.47
1:B:4:GLU:HG2	1:B:69:ASP:HB2	1.97	0.46
1:C:55:LYS:HE2	1:C:59:ALA:HB1	1.95	0.46
1:B:5:MET:HA	1:B:67:ALA:O	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:12:SER:O	1:B:13:ASP:C	2.54	0.46
1:C:138:VAL:HG23	1:C:139:ARG:HG3	1.96	0.46
1:B:144:ASP:CA	1:C:114:LEU:HD22	2.46	0.45
1:C:2:ASN:HB2	1:C:71:ALA:HB3	1.98	0.45
1:A:6:PHE:CD1	1:A:6:PHE:N	2.85	0.45
1:A:114:LEU:HD11	1:A:128:VAL:HG13	1.98	0.45
1:A:73:PRO:HA	1:A:135:ARG:O	2.18	0.44
1:B:143:LYS:C	1:C:114:LEU:CD2	2.86	0.44
1:A:39:ARG:HH22	1:B:129:ASP:HB2	1.83	0.44
1:B:117:GLU:N	1:B:117:GLU:CD	2.71	0.44
1:A:135:ARG:NH1	1:B:131:VAL:HG11	2.33	0.43
1:C:18:ALA:HB3	1:C:120:ILE:HG12	2.00	0.43
1:B:114:LEU:HG	1:B:128:VAL:HG11	2.00	0.43
1:A:28:VAL:HG21	1:A:35:ILE:HG13	1.99	0.43
1:C:23:LEU:HG	1:C:117:GLU:CD	2.39	0.43
1:A:3:ILE:HA	1:A:69:ASP:O	2.19	0.42
1:A:39:ARG:HD3	1:B:65:GLU:OE2	2.19	0.42
1:A:103:ILE:HG22	1:A:104:GLN:N	2.34	0.42
1:C:145:ILE:HG22	1:C:146:PRO:HD2	2.01	0.42
1:A:16:LYS:HD3	1:B:49:GLN:HG2	2.01	0.42
1:A:102:ILE:O	1:A:105:VAL:HG22	2.20	0.42
1:B:19:LEU:HD11	1:B:117:GLU:HG3	2.02	0.42
1:B:114:LEU:HG	1:B:128:VAL:CG1	2.50	0.42
1:C:23:LEU:HD23	1:C:113:GLU:HG2	2.01	0.42
1:A:21:GLY:O	1:A:25:GLN:HG2	2.20	0.42
1:B:101:SER:HB3	1:B:105:VAL:HG23	2.02	0.41
1:C:142:LEU:HD23	1:C:142:LEU:H	1.85	0.41
1:B:22:ALA:O	1:B:26:ALA:N	2.43	0.41
1:B:101:SER:HB3	1:B:105:VAL:CG2	2.50	0.41
1:A:125:ASP:O	1:A:128:VAL:HG12	2.20	0.41
1:B:141:ILE:HG12	1:C:110:GLY:HA3	2.02	0.41
1:C:4:GLU:HA	1:C:41:TRP:O	2.21	0.41
1:C:19:LEU:HD12	1:C:19:LEU:HA	1.94	0.41
1:A:102:ILE:HG13	1:A:105:VAL:HG22	2.03	0.40
1:C:13:ASP:HA	1:C:16:LYS:HD2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/149 (70%)	93 (89%)	11 (11%)	0	100	100
1	B	112/149 (75%)	103 (92%)	9 (8%)	0	100	100
1	C	117/149 (78%)	110 (94%)	7 (6%)	0	100	100
All	All	333/447 (74%)	306 (92%)	27 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/110 (73%)	71 (89%)	9 (11%)	6	6
1	B	88/110 (80%)	80 (91%)	8 (9%)	9	11
1	C	93/110 (84%)	79 (85%)	14 (15%)	3	2
All	All	261/330 (79%)	230 (88%)	31 (12%)	5	5

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ILE
1	A	36	ASP
1	A	37	SER
1	A	40	VAL
1	A	50	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	75	LEU
1	A	103	ILE
1	A	116	LYS
1	A	117	GLU
1	B	36	ASP
1	B	37	SER
1	B	48	THR
1	B	50	PHE
1	B	116	LYS
1	B	117	GLU
1	B	144	ASP
1	B	145	ILE
1	C	35	ILE
1	C	37	SER
1	C	50	PHE
1	C	52	ILE
1	C	75	LEU
1	C	102	ILE
1	C	114	LEU
1	C	116	LYS
1	C	117	GLU
1	C	125	ASP
1	C	128	VAL
1	C	135	ARG
1	C	141	ILE
1	C	145	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	104	GLN
1	B	2	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	108/149 (72%)	1.95	37 (34%) 0 0	47, 74, 108, 140	0
1	B	116/149 (77%)	1.83	40 (34%) 0 0	48, 73, 106, 115	0
1	C	121/149 (81%)	1.81	36 (29%) 0 0	42, 69, 97, 120	0
All	All	345/447 (77%)	1.86	113 (32%) 0 0	42, 72, 106, 140	0

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	75	LEU	8.7
1	B	58	ALA	8.4
1	A	122	ALA	7.3
1	B	116	LYS	7.2
1	B	105	VAL	7.1
1	C	149	ASP	6.9
1	B	73	PRO	6.7
1	C	135	ARG	6.7
1	C	127	ALA	6.6
1	A	108	ILE	6.6
1	A	40	VAL	6.2
1	C	117	GLU	5.9
1	C	128	VAL	5.8
1	A	121	ALA	5.8
1	A	117	GLU	5.6
1	A	31	ILE	5.5
1	C	23	LEU	5.4
1	B	5	MET	5.4
1	B	112	THR	5.3
1	B	126	ALA	5.3
1	C	20	VAL	5.2
1	C	140	VAL	5.2
1	A	119	LEU	5.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1	PRO	5.1
1	C	141	ILE	5.1
1	B	141	ILE	5.1
1	C	147	ASN	5.1
1	A	50	PHE	4.7
1	C	130	THR	4.7
1	C	16	LYS	4.6
1	B	108	ILE	4.5
1	A	114	LEU	4.5
1	C	150	TYR	4.5
1	A	105	VAL	4.3
1	B	51	GLY	4.3
1	C	142	LEU	4.2
1	B	117	GLU	4.2
1	A	107	PHE	4.1
1	C	119	LEU	4.0
1	A	27	THR	3.9
1	A	3	ILE	3.9
1	B	36	ASP	3.8
1	B	140	VAL	3.8
1	C	144	ASP	3.7
1	A	75	LEU	3.7
1	B	22	ALA	3.7
1	C	40	VAL	3.7
1	A	102	ILE	3.7
1	A	111	ARG	3.7
1	A	125	ASP	3.6
1	A	52	ILE	3.6
1	A	11	TYR	3.5
1	B	145	ILE	3.5
1	B	34	PRO	3.5
1	B	114	LEU	3.3
1	A	8	VAL	3.3
1	C	74	SER	3.2
1	A	116	LYS	3.2
1	A	42	LEU	3.2
1	C	7	VAL	3.1
1	A	32	GLY	3.1
1	B	111	ARG	3.1
1	C	39	ARG	3.0
1	B	107	PHE	3.0
1	B	70	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	26	ALA	2.9
1	B	1	PRO	2.9
1	C	106	THR	2.8
1	B	56	THR	2.8
1	B	113	GLU	2.7
1	A	39	ARG	2.7
1	A	25	GLN	2.7
1	A	112	THR	2.7
1	B	32	GLY	2.6
1	A	136	ALA	2.6
1	A	45	VAL	2.6
1	C	120	ILE	2.6
1	C	22	ALA	2.5
1	C	146	PRO	2.5
1	B	33	ALA	2.5
1	C	108	ILE	2.5
1	A	123	LEU	2.5
1	B	143	LYS	2.5
1	C	123	LEU	2.5
1	A	46	PRO	2.4
1	C	31	ILE	2.4
1	A	104	GLN	2.4
1	A	133	ILE	2.4
1	C	111	ARG	2.4
1	B	139	ARG	2.4
1	C	52	ILE	2.3
1	B	37	SER	2.3
1	B	55	LYS	2.3
1	C	113	GLU	2.3
1	B	19	LEU	2.3
1	B	135	ARG	2.2
1	C	132	GLY	2.2
1	A	120	ILE	2.2
1	A	118	ARG	2.2
1	B	120	ILE	2.2
1	B	144	ASP	2.2
1	C	121	ALA	2.2
1	B	23	LEU	2.2
1	C	116	LYS	2.2
1	A	5	MET	2.2
1	C	38	VAL	2.1
1	C	71	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	57	VAL	2.1
1	B	133	ILE	2.1
1	C	24	THR	2.1
1	B	54	GLY	2.1
1	B	124	THR	2.1
1	A	57	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.