



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 6, 2022 – 04:37 pm BST

PDB ID : 6TAV
Title : Crystal structure of endopeptidase-induced alpha2-macroglobulin
Authors : Gomis-Ruth, F.X.; Duquerroy, S.; Trapani, S.; Marrero, A.; Goulas, T.; Guevara, T.; Andersen, G.A.; Sottrup-Jensen, L.; Navaza, J.
Deposited on : 2019-10-30
Resolution : 4.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.27
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1005 (4.62-3.78)
Clashscore	141614	1044 (4.60-3.80)
Ramachandran outliers	138981	1000 (4.60-3.80)
Sidechain outliers	138945	1007 (4.62-3.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	1474	74% (green), 20% (yellow), 6% (orange), 0% (red), 0% (grey)
1	B	1474	66% (green), 18% (yellow), 14% (orange), 2% (red), 2% (grey)
1	C	1474	67% (green), 18% (yellow), 14% (orange), 1% (red), 2% (grey)
1	D	1474	65% (green), 20% (yellow), 14% (orange), 1% (red), 2% (grey)
2	E	4	50% (yellow), 50% (orange)
2	G	4	50% (yellow), 50% (orange)
2	J	4	25% (green), 75% (yellow)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	O	4	 75% 25%
2	V	4	 25% 75%
3	F	3	 33% 67%
3	I	3	 33% 67%
3	M	3	 33% 67%
3	R	3	 67% 33%
4	H	2	 50% 50%
4	K	2	 50% 50%
4	L	2	 50% 50%
4	N	2	 100%
4	P	2	 50% 50%
4	Q	2	 50% 50%
4	S	2	 50% 50%
4	T	2	 50% 50%
4	U	2	 50% 50%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 82015 atoms, of which 40382 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-2-macroglobulin.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	1416	Total 21976	C 7027	H 10926	N 1849	O 2125	S 49	10926	0	0
1	B	1273	Total 19741	C 6312	H 9812	N 1665	O 1909	S 43	9812	0	0
1	C	1274	Total 19770	C 6315	H 9834	N 1668	O 1910	S 43	9834	0	0
1	D	1271	Total 19716	C 6298	H 9810	N 1664	O 1901	S 43	9810	0	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	4	Total 50	C 28	N 2	O 20	0	0	0
2	G	4	Total 50	C 28	N 2	O 20	0	0	0
2	J	4	Total 50	C 28	N 2	O 20	0	0	0
2	O	4	Total 50	C 28	N 2	O 20	0	0	0
2	V	4	Total 50	C 28	N 2	O 20	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	3	39	22	2	15	0	0	0
3	I	3	39	22	2	15	0	0	0
3	M	3	39	22	2	15	0	0	0
3	R	3	39	22	2	15	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	H	2	28	16	2	10	0	0	0
4	K	2	28	16	2	10	0	0	0
4	L	2	28	16	2	10	0	0	0
4	N	2	28	16	2	10	0	0	0
4	P	2	28	16	2	10	0	0	0
4	Q	2	28	16	2	10	0	0	0
4	S	2	28	16	2	10	0	0	0
4	T	2	28	16	2	10	0	0	0
4	U	2	28	16	2	10	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	Total 14	8	1	5	0	0
5	A	1	Total 14	8	1	5	0	0
5	A	1	Total 14	8	1	5	0	0
5	B	1	Total 14	8	1	5	0	0
5	C	1	Total 14	8	1	5	0	0
5	C	1	Total 14	8	1	5	0	0
5	C	1	Total 14	8	1	5	0	0
5	C	1	Total 14	8	1	5	0	0
5	D	1	Total 14	8	1	5	0	0
5	D	1	Total 14	8	1	5	0	0
5	D	1	Total 14	8	1	5	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-2-macroglobulin



Chain O:  75% 25%

MAG1
MAG2
BMA3
MAM4

- Molecule 2: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain V:  25% 75%


MAG1
MAG2
BMA3
MAM4

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  33% 67%


MAG1
MAG2
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  33% 67%

MAG1
MAG2
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  33% 67%

MAG1
MAG2
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain R:  67% 33%

MAG1
MAG2
BMA3

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  50% 50%

MAG1
BMA3

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  50% 50%

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  50% 50%

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:  100%

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P:  50% 50%


MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Q:  50% 50%

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain S:  50% 50%

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain T:  50% 50%

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain U: 

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	130.80Å 260.30Å 281.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.24 – 4.20 95.86 – 4.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (34.24-4.20) 99.2 (95.86-4.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 4.15Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.235 , 0.282 0.268 , 0.294	Depositor DCC
R_{free} test set	834 reflections (1.19%)	wwPDB-VP
Wilson B-factor (Å ²)	221.9	Xtriage
Anisotropy	0.306	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.78	EDS
Total number of atoms	82015	wwPDB-VP
Average B, all atoms (Å ²)	256.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/11297	0.75	1/15352 (0.0%)
1	B	0.54	0/10149	0.72	0/13786
1	C	0.56	0/10156	0.74	2/13798 (0.0%)
1	D	0.53	0/10126	0.74	0/13757
All	All	0.55	0/41728	0.74	3/56693 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	952	GLY	C-N-CA	6.84	138.81	121.70
1	C	953	ASP	N-CA-C	5.73	126.47	111.00
1	A	952	GLY	C-N-CA	5.60	135.70	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11050	10926	10934	88	0
1	B	9929	9812	9818	83	0
1	C	9936	9834	9836	91	0
1	D	9906	9810	9812	97	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	50	0	43	1	0
2	G	50	0	43	1	0
2	J	50	0	43	0	0
2	O	50	0	43	0	0
2	V	50	0	43	0	0
3	F	39	0	34	0	0
3	I	39	0	34	0	0
3	M	39	0	34	0	0
3	R	39	0	34	1	0
4	H	28	0	25	0	0
4	K	28	0	25	0	0
4	L	28	0	25	0	0
4	N	28	0	25	0	0
4	P	28	0	25	0	0
4	Q	28	0	25	0	0
4	S	28	0	25	1	0
4	T	28	0	25	0	0
4	U	28	0	25	0	0
5	A	42	0	39	0	0
5	B	14	0	13	0	0
5	C	56	0	52	0	0
5	D	42	0	39	0	0
All	All	41633	40382	41119	353	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 353 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1147:LYS:HB3	1:D:1170:LEU:HD21	1.54	0.90
1:D:77:LEU:HD22	1:D:87:VAL:HG11	1.53	0.89
1:C:77:LEU:HD22	1:C:87:VAL:HG11	1.55	0.89
1:A:77:LEU:HD22	1:A:87:VAL:HG11	1.56	0.87
1:C:535:ALA:HB1	1:C:536:PRO:CD	2.20	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1412/1474 (96%)	1184 (84%)	168 (12%)	60 (4%)	3	26
1	B	1269/1474 (86%)	1064 (84%)	146 (12%)	59 (5%)	2	24
1	C	1270/1474 (86%)	1082 (85%)	141 (11%)	47 (4%)	3	28
1	D	1267/1474 (86%)	1068 (84%)	145 (11%)	54 (4%)	2	25
All	All	5218/5896 (88%)	4398 (84%)	600 (12%)	220 (4%)	3	26

5 of 220 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125	GLU
1	A	282	ASP
1	A	286	PHE
1	A	374	GLY
1	A	535	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1241/1290 (96%)	1075 (87%)	166 (13%)	4	20
1	B	1110/1290 (86%)	971 (88%)	139 (12%)	4	22
1	C	1112/1290 (86%)	967 (87%)	145 (13%)	4	21
1	D	1108/1290 (86%)	957 (86%)	151 (14%)	3	20
All	All	4571/5160 (89%)	3970 (87%)	601 (13%)	4	21

5 of 601 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	149	VAL
1	D	1056	ILE
1	D	295	ASN
1	D	142	GLN
1	D	644	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1325	GLN
1	D	584	HIS
1	D	1252	GLN
1	D	747	ASN
1	B	827	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

50 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	1,2	14,14,15	0.42	0	17,19,21	1.24	2 (11%)
2	NAG	E	2	2	14,14,15	0.46	0	17,19,21	1.49	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	E	3	2	11,11,12	0.53	0	15,15,17	1.16	1 (6%)
2	MAN	E	4	2	11,11,12	0.58	0	15,15,17	1.06	2 (13%)
3	NAG	F	1	1,3	14,14,15	0.46	0	17,19,21	0.97	1 (5%)
3	NAG	F	2	3	14,14,15	0.41	0	17,19,21	0.91	1 (5%)
3	BMA	F	3	3	11,11,12	0.39	0	15,15,17	0.73	0
2	NAG	G	1	1,2	14,14,15	0.39	0	17,19,21	0.84	1 (5%)
2	NAG	G	2	2	14,14,15	0.48	0	17,19,21	0.95	2 (11%)
2	BMA	G	3	2	11,11,12	0.60	0	15,15,17	1.00	1 (6%)
2	MAN	G	4	2	11,11,12	0.71	0	15,15,17	1.51	2 (13%)
4	NAG	H	1	1,4	14,14,15	0.40	0	17,19,21	1.03	1 (5%)
4	NAG	H	2	4	14,14,15	0.37	0	17,19,21	0.55	0
3	NAG	I	1	1,3	14,14,15	0.33	0	17,19,21	0.66	0
3	NAG	I	2	3	14,14,15	0.40	0	17,19,21	0.89	2 (11%)
3	BMA	I	3	3	11,11,12	0.49	0	15,15,17	0.71	1 (6%)
2	NAG	J	1	1,2	14,14,15	0.54	0	17,19,21	1.20	1 (5%)
2	NAG	J	2	2	14,14,15	0.45	0	17,19,21	1.15	2 (11%)
2	BMA	J	3	2	11,11,12	0.48	0	15,15,17	0.77	0
2	MAN	J	4	2	11,11,12	0.54	0	15,15,17	1.14	2 (13%)
4	NAG	K	1	1,4	14,14,15	0.45	0	17,19,21	0.88	1 (5%)
4	NAG	K	2	4	14,14,15	0.41	0	17,19,21	0.63	0
4	NAG	L	1	1,4	14,14,15	0.44	0	17,19,21	1.16	2 (11%)
4	NAG	L	2	4	14,14,15	0.46	0	17,19,21	0.78	0
3	NAG	M	1	1,3	14,14,15	0.38	0	17,19,21	0.85	1 (5%)
3	NAG	M	2	3	14,14,15	0.45	0	17,19,21	1.06	2 (11%)
3	BMA	M	3	3	11,11,12	0.38	0	15,15,17	0.58	0
4	NAG	N	1	1,4	14,14,15	0.41	0	17,19,21	1.05	2 (11%)
4	NAG	N	2	4	14,14,15	0.46	0	17,19,21	1.58	3 (17%)
2	NAG	O	1	1,2	14,14,15	0.40	0	17,19,21	0.78	0
2	NAG	O	2	2	14,14,15	0.45	0	17,19,21	1.16	3 (17%)
2	BMA	O	3	2	11,11,12	0.58	0	15,15,17	0.76	0
2	MAN	O	4	2	11,11,12	0.43	0	15,15,17	0.86	0
4	NAG	P	1	1,4	14,14,15	0.41	0	17,19,21	0.59	0
4	NAG	P	2	4	14,14,15	0.51	0	17,19,21	0.99	1 (5%)
4	NAG	Q	1	1,4	14,14,15	0.40	0	17,19,21	1.03	2 (11%)
4	NAG	Q	2	4	14,14,15	0.42	0	17,19,21	0.57	0
3	NAG	R	1	1,3	14,14,15	0.29	0	17,19,21	0.69	0
3	NAG	R	2	3	14,14,15	0.46	0	17,19,21	1.52	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BMA	R	3	3	11,11,12	0.50	0	15,15,17	0.93	1 (6%)
4	NAG	S	1	1,4	14,14,15	0.40	0	17,19,21	0.79	1 (5%)
4	NAG	S	2	4	14,14,15	0.42	0	17,19,21	1.14	3 (17%)
4	NAG	T	1	1,4	14,14,15	0.39	0	17,19,21	0.79	0
4	NAG	T	2	4	14,14,15	0.42	0	17,19,21	1.16	2 (11%)
4	NAG	U	1	1,4	14,14,15	0.38	0	17,19,21	1.04	2 (11%)
4	NAG	U	2	4	14,14,15	0.37	0	17,19,21	0.68	0
2	NAG	V	1	1,2	14,14,15	0.40	0	17,19,21	1.38	4 (23%)
2	NAG	V	2	2	14,14,15	0.47	0	17,19,21	1.49	2 (11%)
2	BMA	V	3	2	11,11,12	0.49	0	15,15,17	0.91	0
2	MAN	V	4	2	11,11,12	0.46	0	15,15,17	1.03	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	1/6/23/26	0/1/1/1
2	BMA	E	3	2	-	1/2/19/22	0/1/1/1
2	MAN	E	4	2	-	0/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
2	NAG	G	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
2	BMA	G	3	2	-	2/2/19/22	0/1/1/1
2	MAN	G	4	2	-	1/2/19/22	0/1/1/1
4	NAG	H	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	2/6/23/26	0/1/1/1
3	NAG	I	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	I	2	3	-	0/6/23/26	0/1/1/1
3	BMA	I	3	3	-	0/2/19/22	0/1/1/1
2	NAG	J	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	J	2	2	-	0/6/23/26	0/1/1/1
2	BMA	J	3	2	-	1/2/19/22	0/1/1/1
2	MAN	J	4	2	-	0/2/19/22	0/1/1/1
4	NAG	K	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	K	2	4	-	1/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	L	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	L	2	4	-	2/6/23/26	0/1/1/1
3	NAG	M	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	M	2	3	-	1/6/23/26	0/1/1/1
3	BMA	M	3	3	-	0/2/19/22	0/1/1/1
4	NAG	N	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	N	2	4	-	2/6/23/26	0/1/1/1
2	NAG	O	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	O	2	2	-	3/6/23/26	0/1/1/1
2	BMA	O	3	2	-	2/2/19/22	0/1/1/1
2	MAN	O	4	2	-	0/2/19/22	0/1/1/1
4	NAG	P	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	P	2	4	-	0/6/23/26	0/1/1/1
4	NAG	Q	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	Q	2	4	-	1/6/23/26	0/1/1/1
3	NAG	R	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	R	2	3	-	2/6/23/26	0/1/1/1
3	BMA	R	3	3	-	0/2/19/22	0/1/1/1
4	NAG	S	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	S	2	4	-	1/6/23/26	0/1/1/1
4	NAG	T	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	T	2	4	-	1/6/23/26	0/1/1/1
4	NAG	U	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	U	2	4	-	2/6/23/26	0/1/1/1
2	NAG	V	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	V	2	2	-	0/6/23/26	0/1/1/1
2	BMA	V	3	2	-	2/2/19/22	0/1/1/1
2	MAN	V	4	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	N	2	NAG	C1-C2-N2	4.91	118.87	110.49
2	G	4	MAN	C1-O5-C5	4.72	118.59	112.19
2	V	2	NAG	C1-C2-N2	4.06	117.42	110.49
2	E	2	NAG	C1-O5-C5	4.05	117.69	112.19
2	E	2	NAG	C1-C2-N2	3.73	116.86	110.49

There are no chirality outliers.

5 of 44 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	V	3	BMA	O5-C5-C6-O6
4	P	1	NAG	O5-C5-C6-O6
2	O	2	NAG	O5-C5-C6-O6
4	P	1	NAG	C4-C5-C6-O6
2	O	2	NAG	C4-C5-C6-O6



There are no ring outliers.



7 monomers are involved in 4 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	2	NAG	1	0
2	E	2	NAG	1	0
2	G	1	NAG	1	0
3	R	1	NAG	1	0
2	E	1	NAG	1	0
4	S	1	NAG	1	0
3	R	2	NAG	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.






































5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	D	2007	1	14,14,15	0.36	0	17,19,21	0.53	0
5	NAG	C	2004	1	14,14,15	0.41	0	17,19,21	0.75	0
5	NAG	D	2004	1	14,14,15	0.44	0	17,19,21	1.07	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	C	2001	1	14,14,15	0.43	0	17,19,21	1.50	3 (17%)
5	NAG	A	2019	1	14,14,15	0.49	0	17,19,21	1.43	2 (11%)
5	NAG	D	2003	1	14,14,15	0.45	0	17,19,21	0.82	0
5	NAG	A	2008	1	14,14,15	0.36	0	17,19,21	0.69	0
5	NAG	B	2012	1	14,14,15	0.48	0	17,19,21	1.24	3 (17%)
5	NAG	A	2013	1	14,14,15	0.40	0	17,19,21	0.48	0
5	NAG	C	2002	1	14,14,15	0.38	0	17,19,21	0.70	0
5	NAG	C	2003	1	14,14,15	0.34	0	17,19,21	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	D	2007	1	-	2/6/23/26	0/1/1/1
5	NAG	C	2004	1	-	2/6/23/26	0/1/1/1
5	NAG	D	2004	1	-	1/6/23/26	0/1/1/1
5	NAG	C	2001	1	-	2/6/23/26	0/1/1/1
5	NAG	A	2019	1	-	1/6/23/26	0/1/1/1
5	NAG	D	2003	1	-	2/6/23/26	0/1/1/1
5	NAG	A	2008	1	-	0/6/23/26	0/1/1/1
5	NAG	B	2012	1	-	1/6/23/26	0/1/1/1
5	NAG	A	2013	1	-	2/6/23/26	0/1/1/1
5	NAG	C	2002	1	-	0/6/23/26	0/1/1/1
5	NAG	C	2003	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2019	NAG	C1-O5-C5	4.69	118.55	112.19
5	C	2001	NAG	C1-C2-N2	3.65	116.73	110.49
5	C	2001	NAG	O5-C1-C2	-3.41	105.90	111.29
5	C	2001	NAG	C2-N2-C7	3.07	127.27	122.90
5	B	2012	NAG	C2-N2-C7	2.91	127.05	122.90

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	2007	NAG	O5-C5-C6-O6
5	A	2013	NAG	O5-C5-C6-O6
5	D	2003	NAG	C4-C5-C6-O6
5	D	2007	NAG	C4-C5-C6-O6
5	D	2003	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain O:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain V:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)













Electron density around Chain H:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain K:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around Chain N:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain P:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain Q:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain T:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.