



Full wwPDB X-ray Structure Validation Report

May 16, 2020 – 04:56 am BST

PDB ID : 1TGH
Title : TATA BINDING PROTEIN (TBP)/DNA COMPLEX
Authors : Juo, Z.S.; Dickerson, R.E.
Deposited on : 1996-02-13
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	12	 17% 83%
1	C	12	 17% 75% 8%
2	A	185	 % 45% 44% 9% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2355 atoms, of which 422 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*GP*TP*AP*TP*AP*TP*AP*TP*AP*CP*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
1	B	12	267	118	24	44	70	11	0	0	0
1	C	12	267	118	24	44	70	11	0	0	0

- Molecule 2 is a protein called PROTEIN (TATA BINDING PROTEIN (TBP)).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	A	180	1767	927	338	252	243	7	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	151	GLY	ALA	CONFLICT	UNP P20226
A	153	ARG	GLU	CONFLICT	UNP P20226
A	154	GLY	SER	CONFLICT	UNP P20226

- Molecule 3 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
3	B	5	15	10	5	0	0
3	C	1	3	2	1	0	0
3	A	12	36	24	12	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*CP*GP*TP*AP*TP*AP*TP*AP*TP*AP*CP*G)-3')

Chain B: 



- Molecule 1: DNA (5'-D(*CP*GP*TP*AP*TP*AP*TP*AP*TP*AP*CP*G)-3')

Chain C: 



- Molecule 2: PROTEIN (TATA BINDING PROTEIN (TBP))

Chain A: 







4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.97Å 67.40Å 86.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.90 36.32 – 2.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-2.90) 98.9 (36.32-2.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.15 (at 2.90Å)	Xtrriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.214 , 0.294 0.213 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	61.7	Xtrriage
Anisotropy	0.398	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 73.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.028 for k,h,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2355	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	1.03	0/272	1.07	0/418
1	C	0.99	1/272 (0.4%)	1.03	0/418
2	A	0.68	0/1455	0.91	0/1958
All	All	0.79	1/1999 (0.1%)	0.95	0/2794

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	120	DA	C5-C6	-5.92	1.35	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	243	24	138	33	0
1	C	243	24	138	21	0
2	A	1429	338	1521	99	0
3	A	12	24	0	0	0
3	B	5	10	0	0	0
3	C	1	2	0	0	0
All	All	1933	422	1797	132	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 36.

All (132) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:108:DA:H2''	1:B:109:DT:C5'	1.70	1.20
1:C:123:DC:H2''	1:C:124:DG:C8	1.87	1.10
1:B:108:DA:H2''	1:B:109:DT:H5''	1.20	1.10
1:B:102:DG:H2''	1:B:103:DT:H5'	1.38	1.05
1:B:109:DT:H2''	1:B:110:DA:H5''	1.37	1.01
1:B:102:DG:H2''	1:B:103:DT:C5'	1.91	0.98
1:B:105:DT:H5''	2:A:288:ILE:HD13	1.54	0.90
1:B:109:DT:C6	1:B:109:DT:H5'	2.08	0.89
1:C:117:DT:H5''	2:A:197:ILE:HD13	1.56	0.88
1:B:101:DC:H2''	1:B:102:DG:C8	2.08	0.87
1:B:108:DA:H2''	1:B:109:DT:H5'	1.55	0.87
2:A:277:SER:HB3	2:A:287:LEU:HD11	1.57	0.86
1:B:109:DT:C2'	1:B:110:DA:H5''	2.07	0.85
2:A:176:LEU:HD22	2:A:196:VAL:HG12	1.60	0.84
1:B:108:DA:C2'	1:B:109:DT:H5''	2.07	0.81
1:B:109:DT:H1'	2:A:210:PHE:CE2	2.15	0.81
2:A:192:ARG:HH22	2:A:197:ILE:HD12	1.47	0.78
2:A:196:VAL:HG13	2:A:209:ILE:HB	1.69	0.75
1:B:109:DT:H6	1:B:109:DT:H5'	1.47	0.74
1:C:123:DC:C2'	1:C:124:DG:C8	2.69	0.73
2:A:313:VAL:HG12	2:A:315:ALA:H	1.56	0.71
1:B:102:DG:H2''	1:B:103:DT:H5''	1.72	0.71
2:A:270:VAL:HG12	2:A:274:GLN:NE2	2.04	0.70
1:B:108:DA:C2'	1:B:109:DT:C5'	2.61	0.70
2:A:266:LEU:O	2:A:270:VAL:HG23	1.92	0.69
2:A:232:LYS:O	2:A:236:VAL:HG23	1.94	0.68
1:B:109:DT:C2'	1:B:110:DA:C5'	2.73	0.67
2:A:290:ARG:HH11	2:A:290:ARG:HG3	1.60	0.66
2:A:155:SER:HB2	2:A:325:TYR:CD2	2.31	0.66
1:C:116:DA:C5'	2:A:192:ARG:HD3	2.25	0.65
1:B:105:DT:C5'	2:A:288:ILE:HD13	2.25	0.65
1:B:101:DC:C2'	1:B:102:DG:C8	2.79	0.65
2:A:266:LEU:HD22	2:A:287:LEU:HB2	1.80	0.63
1:B:105:DT:H5''	2:A:288:ILE:CD1	2.27	0.62
2:A:272:THR:HG1	2:A:331:PHE:HZ	1.48	0.62
2:A:270:VAL:HG21	2:A:279:TYR:HD2	1.65	0.62
2:A:209:ILE:HD13	2:A:215:MET:HG3	1.82	0.61
1:C:123:DC:C2'	1:C:124:DG:N7	2.63	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:172:CYS:O	2:A:174:LEU:HD13	2.01	0.60
1:B:106:DA:H5''	2:A:297:VAL:HG21	1.84	0.59
2:A:279:TYR:CE1	2:A:281:PRO:HB3	2.37	0.59
1:C:113:DC:H2''	1:C:114:DG:OP2	2.02	0.59
2:A:177:LYS:O	2:A:181:LEU:HG	2.03	0.59
2:A:209:ILE:CD1	2:A:215:MET:SD	2.91	0.58
1:C:123:DC:H5'	1:C:123:DC:H6	1.69	0.57
1:B:102:DG:C2'	1:B:103:DT:C5'	2.77	0.57
2:A:314:ARG:HG2	2:A:318:TYR:CE2	2.38	0.57
2:A:192:ARG:NH2	2:A:197:ILE:HD12	2.17	0.57
2:A:164:ILE:HD13	2:A:254:MET:HG2	1.86	0.57
1:C:123:DC:H2''	1:C:124:DG:N7	2.21	0.56
2:A:260:VAL:HG22	2:A:304:GLY:CA	2.36	0.55
2:A:269:LEU:HD13	2:A:287:LEU:CD2	2.35	0.55
2:A:188:TYR:CZ	2:A:190:PRO:HD3	2.43	0.54
2:A:270:VAL:HG12	2:A:274:GLN:HE22	1.72	0.54
1:B:101:DC:HO5'	1:B:101:DC:H6	1.54	0.54
1:C:120:DA:H2''	1:C:121:DT:O5'	2.08	0.53
2:A:262:PHE:CZ	2:A:329:LYS:HA	2.43	0.53
2:A:209:ILE:HD11	2:A:215:MET:SD	2.48	0.53
2:A:175:ASP:O	2:A:179:ILE:HG13	2.10	0.52
2:A:191:LYS:NZ	2:A:192:ARG:HB3	2.23	0.52
1:C:116:DA:H4'	2:A:192:ARG:CD	2.39	0.52
2:A:250:LYS:HD3	2:A:252:GLN:HE22	1.75	0.52
1:B:105:DT:O2	2:A:299:LEU:HD21	2.08	0.52
1:B:106:DA:H5''	2:A:297:VAL:CG2	2.39	0.52
2:A:209:ILE:HD13	2:A:215:MET:CG	2.40	0.52
2:A:227:ARG:O	2:A:230:ALA:HB3	2.10	0.52
2:A:278:SER:O	2:A:287:LEU:HD12	2.10	0.51
2:A:187:GLU:HB2	2:A:197:ILE:HB	1.92	0.50
2:A:157:ILE:HG22	2:A:158:VAL:N	2.26	0.50
1:C:123:DC:H2''	1:C:124:DG:H8	1.67	0.50
2:A:260:VAL:O	2:A:261:LYS:HB2	2.11	0.49
1:B:102:DG:C2'	1:B:103:DT:H5''	2.41	0.49
1:B:107:DT:C5	1:B:108:DA:N6	2.81	0.48
1:C:116:DA:H4'	2:A:192:ARG:HD3	1.95	0.48
1:C:123:DC:H5'	1:C:123:DC:C6	2.47	0.48
2:A:236:VAL:O	2:A:240:LEU:HD13	2.13	0.48
1:B:108:DA:C2'	1:B:109:DT:H5'	2.36	0.48
1:B:107:DT:H4'	2:A:252:GLN:O	2.14	0.48
2:A:259:ASP:HB3	2:A:261:LYS:NZ	2.29	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:209:ILE:HD13	2:A:215:MET:SD	2.54	0.47
1:B:102:DG:C2'	1:B:103:DT:H5'	2.28	0.47
2:A:324:ILE:O	2:A:328:LEU:HG	2.15	0.47
2:A:233:TYR:O	2:A:237:VAL:HG23	2.14	0.47
2:A:290:ARG:CG	2:A:290:ARG:HH11	2.27	0.47
2:A:227:ARG:HD3	2:A:249:PHE:CD2	2.50	0.47
1:B:108:DA:H4'	2:A:252:GLN:HG3	1.97	0.47
1:B:104:DA:OP1	2:A:283:LEU:HD21	2.16	0.46
2:A:260:VAL:HG22	2:A:304:GLY:HA3	1.97	0.46
1:C:116:DA:O5'	2:A:192:ARG:HD3	2.16	0.46
2:A:238:GLN:HE21	2:A:244:ALA:HB3	1.81	0.46
2:A:269:LEU:HD13	2:A:287:LEU:HD23	1.96	0.46
2:A:220:ALA:O	2:A:221:LYS:HG2	2.16	0.45
2:A:252:GLN:OE1	2:A:252:GLN:HA	2.15	0.45
1:B:108:DA:C4'	2:A:252:GLN:HG3	2.46	0.45
2:A:210:PHE:HB2	2:A:214:LYS:O	2.17	0.45
2:A:272:THR:OG1	2:A:273:HIS:HD2	1.99	0.45
1:C:119:DT:H6	1:C:119:DT:H2'	1.61	0.45
2:A:206:THR:HG22	2:A:207:ALA:N	2.31	0.44
2:A:284:PHE:CG	2:A:285:PRO:HD2	2.52	0.44
1:C:116:DA:C2	2:A:208:LEU:HD13	2.53	0.44
2:A:270:VAL:HG22	2:A:287:LEU:HD13	2.00	0.44
2:A:235:ARG:HH11	2:A:235:ARG:HG2	1.82	0.44
2:A:200:ILE:HG23	2:A:233:TYR:CE1	2.53	0.43
2:A:191:LYS:HZ2	2:A:192:ARG:HB3	1.82	0.43
1:C:116:DA:C4'	2:A:192:ARG:HD3	2.49	0.43
2:A:165:VAL:HG13	2:A:218:THR:HG22	2.00	0.43
2:A:159:PRO:HA	2:A:258:CYS:HB3	2.00	0.43
2:A:266:LEU:HB3	2:A:279:TYR:CE2	2.53	0.43
2:A:284:PHE:CD1	2:A:285:PRO:HD2	2.53	0.43
2:A:260:VAL:HG22	2:A:304:GLY:O	2.19	0.43
1:C:118:DA:OP1	2:A:199:ARG:NH1	2.51	0.43
2:A:219:GLY:O	2:A:220:ALA:O	2.36	0.43
2:A:261:LYS:O	2:A:262:PHE:HB3	2.18	0.42
2:A:170:LEU:HD12	2:A:174:LEU:HD21	2.00	0.42
2:A:298:LEU:N	2:A:298:LEU:HD12	2.35	0.42
1:C:123:DC:H2'	1:C:124:DG:N7	2.34	0.42
2:A:266:LEU:HD22	2:A:287:LEU:CB	2.48	0.42
2:A:195:ALA:HB2	2:A:210:PHE:CE1	2.55	0.42
1:B:106:DA:N7	1:B:107:DT:C4	2.88	0.42
2:A:185:ASN:OD1	2:A:201:ARG:NH1	2.49	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:188:TYR:CE2	2:A:190:PRO:HD3	2.55	0.41
1:C:116:DA:C2	2:A:193:PHE:CZ	3.08	0.41
2:A:297:VAL:O	2:A:309:THR:HG22	2.20	0.41
2:A:189:ASN:N	2:A:189:ASN:OD1	2.54	0.41
2:A:324:ILE:HG13	2:A:328:LEU:HG	2.03	0.41
2:A:265:ARG:HB2	2:A:267:GLU:OE2	2.20	0.41
2:A:177:LYS:HD2	2:A:181:LEU:HD21	2.03	0.41
1:C:119:DT:H5''	2:A:221:LYS:HZ1	1.85	0.41
2:A:329:LYS:HB3	2:A:329:LYS:HE2	1.82	0.40
2:A:279:TYR:CZ	2:A:281:PRO:HB3	2.57	0.40
2:A:206:THR:CG2	2:A:207:ALA:N	2.84	0.40
2:A:291:MET:HE3	2:A:298:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	178/185 (96%)	150 (84%)	24 (14%)	4 (2%)	6 24

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	220	ALA
2	A	172	CYS
2	A	184	ARG
2	A	262	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	155/158 (98%)	137 (88%)	18 (12%)	5 16

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	174	LEU
2	A	184	ARG
2	A	185	ASN
2	A	189	ASN
2	A	191	LYS
2	A	192	ARG
2	A	196	VAL
2	A	211	SER
2	A	214	LYS
2	A	228	LEU
2	A	240	LEU
2	A	267	GLU
2	A	269	LEU
2	A	290	ARG
2	A	292	ILE
2	A	293	LYS
2	A	299	LEU
2	A	302	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	273	HIS
2	A	274	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	12/12 (100%)	-0.42	0 100 100	17, 31, 40, 43	0
1	C	12/12 (100%)	-0.63	0 100 100	22, 28, 34, 34	0
2	A	180/185 (97%)	-0.33	2 (1%) 80 80	10, 29, 55, 71	0
All	All	204/209 (97%)	-0.35	2 (0%) 82 82	10, 29, 55, 71	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	192	ARG	2.9
2	A	260	VAL	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.