



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 23, 2024 – 12:51 AM EDT

PDB ID : 7THO
Title : Integrin alpha IIB beta3 complex with Eptifibatide
Authors : Zhu, J.; Lin, F.-Y.; Zhu, J.; Springer, T.A.
Deposited on : 2022-01-11
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.002 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1606 (2.78-2.74)
Clashscore	180529	1689 (2.78-2.74)
Ramachandran outliers	177936	1665 (2.78-2.74)
Sidechain outliers	177891	1665 (2.78-2.74)
RSRZ outliers	164620	1606 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	
1	C	454	
2	B	471	
2	D	471	
3	E	216	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	216	<p>% 96%</p>
4	F	214	<p>7% 97%</p>
4	L	214	<p>2% 95% 5%</p>
5	M	8	<p>12% 50% 38% 12%</p>
5	N	8	<p>62% 25% 12%</p>
6	G	4	<p>25% 25% 50%</p>
7	I	2	<p>50% 50%</p>
7	K	2	<p>50% 50%</p>
8	J	3	<p>33% 67%</p>

2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 21512 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin alpha-IIb.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	454	Total	C	N	O	S	0	5	0
			3517	2235	609	665	8			
1	C	453	Total	C	N	O	S	0	2	0
			3484	2214	600	662	8			

- Molecule 2 is a protein called Integrin beta-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	466	Total	C	N	O	S	4	2	0
			3601	2243	615	710	33			
2	D	471	Total	C	N	O	S	13	2	0
			3638	2269	620	715	34			

- Molecule 3 is a protein called Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	214	Total	C	N	O	S	0	0	0
			1631	1035	264	326	6			
3	H	216	Total	C	N	O	S	0	0	0
			1642	1041	266	329	6			

- Molecule 4 is a protein called Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	F	214	Total	C	N	O	S	0	0	0
			1637	1019	268	341	9			
4	L	214	Total	C	N	O	S	0	0	0
			1637	1019	268	341	9			

- Molecule 5 is a protein called Eptifibatide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				S
5	M	8	57	35	11	9	2	0	0	1
5	N	8	57	35	11	9	2	0	0	1

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	G	4	50	28	2	20	0	0	0

- Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
7	I	2	28	16	2	10	0	0	0
7	K	2	28	16	2	10	0	0	0

- Molecule 8 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
8	J	3	39	22	2	15	0	0	0

- Molecule 9 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	4	Total Ca 4 4	0	0
9	B	2	Total Ca 2 2	0	0
9	C	4	Total Ca 4 4	0	0
9	D	2	Total Ca 2 2	0	0

- Molecule 10 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total O S 5 4 1	0	0
10	C	1	Total O S 5 4 1	0	0
10	C	1	Total O S 5 4 1	0	0
10	L	1	Total O S 5 4 1	0	0

- Molecule 11 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	B	1	Total Mg 1 1	0	0
11	D	1	Total Mg 1 1	0	0

- Molecule 12 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
12	B	1	14	8	1	5	0	0
12	D	1	14	8	1	5	0	0

- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
13	A	176	176	176	0	0
13	B	68	68	68	0	0
13	C	109	109	109	0	0
13	D	32	32	32	0	0
13	E	2	2	2	0	0
13	F	1	1	1	0	0
13	H	7	7	7	0	0
13	L	8	8	8	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	M	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Integrin alpha-IIb

Chain A: 




- Molecule 1: Integrin alpha-IIb

Chain C: 





- Molecule 2: Integrin beta-3

Chain B: 




- Molecule 2: Integrin beta-3

Chain D: 




- Molecule 3: Fab heavy chain




- Molecule 3: Fab heavy chain



- Molecule 4: Fab light chain




- Molecule 4: Fab light chain



- Molecule 5: Eptifibatide



- Molecule 5: Eptifibatide



- Molecule 6: beta-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  50% 50%

MAG1
MAG2

- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  50% 50%

MAG1
MAG2

- Molecule 8: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  33% 67%

MAG1
MAG2
BMA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	260.34Å 145.04Å 104.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.44 – 2.75 48.44 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.44-2.75) 99.9 (48.44-2.75)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 2.77Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.189 , 0.222 0.191 , 0.225	Depositor DCC
R_{free} test set	1230 reflections (1.19%)	wwPDB-VP
Wilson B-factor (Å ²)	68.3	Xtrriage
Anisotropy	0.382	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 61.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	21512	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPT, CA, NH2, HRG, NAG, MG, BMA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3623	0.53	0/4936
1	C	0.25	0/3587	0.51	0/4888
2	B	0.24	0/3674	0.48	0/4982
2	D	0.24	0/3713	0.49	0/5036
3	E	0.24	0/1673	0.47	0/2290
3	H	0.24	0/1684	0.47	0/2305
4	F	0.25	0/1673	0.48	0/2269
4	L	0.24	0/1673	0.47	0/2269
5	M	1.78	1/41 (2.4%)	1.97	1/56 (1.8%)
5	N	1.85	1/41 (2.4%)	1.91	1/56 (1.8%)
All	All	0.27	2/21382 (0.0%)	0.51	2/29087 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	N	5	TRP	C-N	5.91	1.45	1.34
5	M	5	TRP	C-N	5.42	1.44	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	N	6	PRO	O-C-N	-6.63	112.09	122.70
5	M	6	PRO	O-C-N	-6.43	112.41	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3517	0	3362	15	0
1	C	3484	0	3321	10	0
2	B	3601	0	3523	23	0
2	D	3638	0	3559	22	0
3	E	1631	0	1590	6	0
3	H	1642	0	1600	5	0
4	F	1637	0	1553	3	0
4	L	1637	0	1553	5	0
5	M	57	0	46	2	0
5	N	57	0	45	0	0
6	G	50	0	43	1	0
7	I	28	0	25	1	0
7	K	28	0	25	2	0
8	J	39	0	34	0	0
9	A	4	0	0	0	0
9	B	2	0	0	0	0
9	C	4	0	0	0	0
9	D	2	0	0	0	0
10	A	5	0	0	0	0
10	C	10	0	0	0	0
10	L	5	0	0	0	0
11	B	1	0	0	0	0
11	D	1	0	0	0	0
12	B	14	0	13	0	0
12	D	14	0	13	2	0
13	A	176	0	0	4	1
13	B	68	0	0	2	0
13	C	109	0	0	2	1
13	D	32	0	0	1	0
13	E	2	0	0	0	0
13	F	1	0	0	0	0
13	H	7	0	0	0	0
13	L	8	0	0	0	0
13	M	1	0	0	0	0
All	All	21512	0	20305	89	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:127:ASP:OD2	13:B:2101:HOH:O	2.06	0.72
1:A:400[B]:ARG:NH1	13:A:601:HOH:O	2.28	0.65
2:D:3:ASN:O	2:D:7:THR:OG1	2.15	0.64
1:A:400[A]:ARG:NH1	13:A:602:HOH:O	2.30	0.64
1:C:90:ARG:NH2	13:C:604:HOH:O	2.32	0.62
2:B:141:GLN:HG3	2:B:341:LEU:HD21	1.83	0.59
2:B:65:GLU:OE1	13:B:2102:HOH:O	2.18	0.56
6:G:3:BMA:O2	6:G:4:BMA:O5	2.22	0.56
2:D:306:LEU:HB3	2:D:328:THR:HG22	1.89	0.55
2:D:108:GLU:HG2	2:D:391:ILE:HG22	1.89	0.55
2:D:219:PRO:HB3	2:D:253:LYS:HB2	1.89	0.55
1:C:422:ARG:NH2	13:C:607:HOH:O	2.36	0.55
3:E:149:LYS:HZ3	3:E:182:THR:HB	1.72	0.54
2:B:386:CYS:HB3	2:B:389:LEU:HD11	1.88	0.54
2:B:14:GLN:HB2	2:B:438:GLN:HE22	1.72	0.54
2:B:374:CYS:O	2:B:376:ASN:N	2.41	0.53
2:D:249:THR:HG22	2:D:309:ALA:HB3	1.91	0.52
4:L:4:MET:HE1	4:L:90:GLN:HB3	1.92	0.51
2:D:114:ILE:HG12	2:D:245:LEU:HD12	1.92	0.51
1:C:390:LEU:HD23	1:C:403:PRO:HG3	1.92	0.51
2:D:400:GLU:HB2	7:K:1:NAG:H83	1.92	0.51
1:C:230:TYR:OH	1:C:279:ARG:NH2	2.44	0.50
2:D:125:LYS:HA	2:D:212:VAL:HG11	1.92	0.50
1:A:422:ARG:NH2	13:A:606:HOH:O	2.38	0.49
2:B:400:GLU:HB2	7:I:1:NAG:H83	1.95	0.49
2:D:177:CYS:HB3	2:D:182:THR:HG23	1.95	0.49
2:B:288:ASP:OD1	2:B:289:TYR:N	2.46	0.48
3:H:213:ASP:OD1	3:H:213:ASP:N	2.46	0.48
2:D:288:ASP:OD1	2:D:289:TYR:N	2.42	0.48
2:B:139:ALA:HB2	2:B:200:VAL:HG11	1.96	0.48
1:C:107:CYS:HA	1:C:130:CYS:HA	1.95	0.48
2:B:249:THR:HG22	2:B:309:ALA:HB3	1.97	0.47
1:C:224:ASP:OD1	1:C:225:SER:N	2.42	0.47
2:D:449:ASN:HB3	2:D:452:ASN:HB2	1.96	0.47
1:A:158:ASN:ND2	13:A:610:HOH:O	2.42	0.47
2:B:376:ASN:O	2:B:378:GLU:HG3	2.15	0.47
1:C:259:THR:HA	1:C:263:THR:HA	1.97	0.46
3:E:194:TRP:CG	3:E:195:PRO:HA	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:223:PHE:CZ	2:B:254:THR:HG21	2.51	0.46
2:B:51:PRO:HA	2:B:54:ILE:HD13	1.98	0.46
4:F:37:GLN:HB2	4:F:47:LEU:HD11	1.98	0.46
1:A:3:LEU:O	1:A:5:PRO:HD3	2.16	0.46
3:H:125:PRO:HB3	3:H:151:TYR:HB3	1.98	0.45
3:H:194:TRP:CG	3:H:195:PRO:HA	2.51	0.44
2:D:223:PHE:CZ	2:D:254:THR:HG21	2.53	0.44
1:A:5:PRO:HA	1:A:8:LEU:HD21	2.00	0.44
1:A:141:ALA:HA	1:A:207:TYR:CD2	2.53	0.44
1:A:224:ASP:OD1	5:M:2:HRG:NH1	2.35	0.44
1:A:394:GLY:HA2	1:A:399:LEU:HD23	2.00	0.44
2:B:464:PRO:HA	2:B:465:GLY:HA2	1.66	0.43
1:C:3:LEU:O	1:C:5:PRO:HD3	2.18	0.43
2:D:32:PRO:HB2	2:D:35:SER:HB2	1.99	0.43
2:D:217:ASP:OD2	2:D:255:HIS:NE2	2.51	0.43
2:D:281:TYR:CE1	2:D:283:ALA:HB3	2.53	0.43
4:F:193:THR:HG23	4:F:206:VAL:HG13	2.00	0.43
1:A:215:HIS:CD2	3:H:32:THR:HG22	2.54	0.43
1:C:303:ARG:NH2	1:C:335:ARG:HG2	2.35	0.42
12:D:2004:NAG:O6	7:K:2:NAG:O6	2.27	0.42
1:A:187:GLY:HA2	1:A:191:PHE:HA	2.01	0.42
1:A:235:TRP:CZ2	1:A:270:LEU:HD11	2.54	0.42
3:E:30:LYS:HG3	3:E:54:ALA:HA	2.02	0.42
3:E:20:LEU:HD22	3:E:113:THR:HG21	2.01	0.42
3:E:102:ASP:HB3	3:E:105:ALA:HB2	2.01	0.42
3:H:102:ASP:HB3	3:H:105:ALA:HB2	2.01	0.42
2:B:215:ASN:ND2	5:M:4:ASP:OD1	2.53	0.42
2:D:409:GLU:HB2	2:D:412:LYS:HE3	2.02	0.42
1:A:262:TRP:CD1	2:B:253:LYS:HE2	2.55	0.42
2:D:313:ASN:ND2	2:D:314[B]:VAL:HG13	2.34	0.41
1:C:319:ASP:O	1:C:321:LYS:HG2	2.20	0.41
2:D:375:LEU:HD22	2:D:375:LEU:H	1.85	0.41
2:B:3:ASN:ND2	2:B:40:LEU:HG	2.36	0.41
2:B:311:THR:HA	2:B:333:LEU:O	2.20	0.41
4:L:21:ILE:O	4:L:72:SER:HA	2.20	0.41
12:D:2004:NAG:H3	12:D:2004:NAG:C8	2.50	0.41
2:B:292:LEU:HD22	2:B:325:ILE:HD11	2.03	0.41
4:F:120:PRO:HG2	4:F:130:ALA:HB1	2.01	0.41
4:L:37:GLN:HB2	4:L:47:LEU:HD11	2.02	0.41
2:B:25:TRP:HA	2:B:37:ARG:O	2.19	0.41
3:E:213:ASP:OD1	3:E:213:ASP:N	2.54	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:CYS:HA	1:A:130:CYS:HA	2.01	0.41
2:D:77:SER:HA	2:D:78:SER:HA	1.54	0.41
4:L:167:ASP:OD1	4:L:168:SER:N	2.53	0.41
2:B:202:ARG:NH2	2:B:206:GLU:OE2	2.54	0.41
2:D:67:ARG:HB2	2:D:86:GLN:NE2	2.36	0.41
1:A:262:TRP:HB3	2:B:317:LEU:HD13	2.03	0.40
2:B:449:ASN:HB2	2:B:452:ASN:HB2	2.03	0.40
4:L:149:LYS:HB2	4:L:193:THR:HB	2.03	0.40
2:D:3:ASN:ND2	2:D:40:LEU:HG	2.36	0.40
2:D:216:ARG:NH1	13:D:2105:HOH:O	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
13:A:760:HOH:O	13:C:672:HOH:O[1_554]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/454 (101%)	437 (96%)	17 (4%)	3 (1%)	19	32
1	C	453/454 (100%)	434 (96%)	19 (4%)	0	100	100
2	B	466/471 (99%)	440 (94%)	24 (5%)	2 (0%)	30	47
2	D	471/471 (100%)	448 (95%)	23 (5%)	0	100	100
3	E	210/216 (97%)	201 (96%)	9 (4%)	0	100	100
3	H	212/216 (98%)	202 (95%)	10 (5%)	0	100	100
4	F	212/214 (99%)	199 (94%)	13 (6%)	0	100	100
4	L	212/214 (99%)	201 (95%)	11 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	M	5/8 (62%)	5 (100%)	0	0	100	100
5	N	5/8 (62%)	2 (40%)	2 (40%)	1 (20%)	0	0
All	All	2703/2726 (99%)	2569 (95%)	128 (5%)	6 (0%)	44	63

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	157	GLU
2	B	76	ASP
2	B	375	LEU
1	A	123	GLU
1	A	158	ASN
5	N	6	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/362 (101%)	362 (99%)	5 (1%)	62	78
1	C	363/362 (100%)	359 (99%)	4 (1%)	70	83
2	B	414/416 (100%)	412 (100%)	2 (0%)	86	92
2	D	418/416 (100%)	414 (99%)	4 (1%)	73	84
3	E	186/187 (100%)	186 (100%)	0	100	100
3	H	187/187 (100%)	187 (100%)	0	100	100
4	F	188/188 (100%)	188 (100%)	0	100	100
4	L	188/188 (100%)	188 (100%)	0	100	100
5	M	4/4 (100%)	4 (100%)	0	100	100
5	N	4/4 (100%)	4 (100%)	0	100	100
All	All	2319/2314 (100%)	2304 (99%)	15 (1%)	84	90

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	LEU
1	A	166	TYR
1	A	190	TYR
1	A	219	GLN
1	A	288	TYR
2	B	182	THR
2	B	215	ASN
1	C	23	LEU
1	C	166	TYR
1	C	270	LEU
1	C	288	TYR
2	D	7	THR
2	D	86	GLN
2	D	127	ASP
2	D	375	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	GLN
1	A	338	HIS
2	B	438	GLN
1	C	7	GLN
1	C	451	GLN
2	D	132	GLN
2	D	280	HIS
2	D	428	GLN
3	H	82	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	HRG	M	2	5	10,11,12	2.56	3 (30%)	6,12,14	2.12	2 (33%)
5	HRG	N	2	5	10,11,12	2.66	2 (20%)	6,12,14	2.21	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HRG	M	2	5	-	1/9/10/12	-
5	HRG	N	2	5	-	2/9/10/12	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	N	2	HRG	CB-CA	5.70	1.62	1.53
5	M	2	HRG	CB-CA	5.38	1.61	1.53
5	N	2	HRG	CZ-NE	5.13	1.43	1.33
5	M	2	HRG	CZ-NE	4.82	1.42	1.33
5	M	2	HRG	CZ-NH1	2.09	1.39	1.32

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	N	2	HRG	NE-CZ-NH1	4.54	128.47	120.67
5	M	2	HRG	NH2-CZ-NE	3.96	128.26	119.27
5	M	2	HRG	NE-CZ-NH1	-2.19	116.91	120.67

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	N	2	HRG	NE-CD-CG-CG'
5	N	2	HRG	CD-CG-CG'-CB
5	M	2	HRG	NE-CD-CG-CG'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	M	2	HRG	1	0

5.5 Carbohydrates [i](#)

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	G	1	6,2	14,14,15	0.39	0	17,19,21	0.66	0
6	NAG	G	2	6	14,14,15	0.58	1 (7%)	17,19,21	0.63	0
6	BMA	G	3	6	11,11,12	2.04	3 (27%)	15,15,17	1.80	3 (20%)
6	BMA	G	4	6	11,11,12	0.72	0	15,15,17	1.20	2 (13%)
7	NAG	I	1	7,2	14,14,15	0.66	1 (7%)	17,19,21	0.70	0
7	NAG	I	2	7	14,14,15	0.22	0	17,19,21	0.64	0
8	NAG	J	1	8,2	14,14,15	0.55	0	17,19,21	0.83	1 (5%)
8	NAG	J	2	8	14,14,15	0.21	0	17,19,21	0.50	0
8	BMA	J	3	8	11,11,12	0.90	0	15,15,17	1.30	1 (6%)
7	NAG	K	1	7,2	14,14,15	0.82	1 (7%)	17,19,21	0.83	0
7	NAG	K	2	7	14,14,15	0.28	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	G	1	6,2	-	0/6/23/26	0/1/1/1
6	NAG	G	2	6	-	0/6/23/26	0/1/1/1
6	BMA	G	3	6	-	2/2/19/22	0/1/1/1
6	BMA	G	4	6	-	0/2/19/22	0/1/1/1
7	NAG	I	1	7,2	-	2/6/23/26	0/1/1/1
7	NAG	I	2	7	-	3/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	J	1	8,2	-	0/6/23/26	0/1/1/1
8	NAG	J	2	8	-	1/6/23/26	0/1/1/1
8	BMA	J	3	8	-	2/2/19/22	1/1/1/1
7	NAG	K	1	7,2	-	2/6/23/26	0/1/1/1
7	NAG	K	2	7	-	0/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	G	3	BMA	C2-C3	5.01	1.60	1.52
6	G	3	BMA	O3-C3	3.12	1.50	1.43
6	G	3	BMA	C4-C3	2.38	1.58	1.52
7	K	1	NAG	C1-C2	2.34	1.55	1.52
7	I	1	NAG	O5-C1	-2.06	1.40	1.43
6	G	2	NAG	O5-C1	-2.05	1.40	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	3	BMA	O3-C3-C2	5.39	121.05	110.05
8	J	3	BMA	C1-O5-C5	4.29	117.94	112.19
6	G	4	BMA	C1-O5-C5	3.53	116.92	112.19
8	J	1	NAG	C1-O5-C5	2.75	115.87	112.19
6	G	4	BMA	O2-C2-C3	-2.28	105.44	110.15
6	G	3	BMA	C3-C4-C5	2.18	114.19	110.23
6	G	3	BMA	O2-C2-C3	2.02	114.33	110.15

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	J	3	BMA	O5-C5-C6-O6
6	G	3	BMA	C4-C5-C6-O6
7	I	2	NAG	C8-C7-N2-C2
7	I	2	NAG	O7-C7-N2-C2
7	I	2	NAG	O5-C5-C6-O6
7	K	1	NAG	O5-C5-C6-O6
7	I	1	NAG	O5-C5-C6-O6
6	G	3	BMA	O5-C5-C6-O6
7	I	1	NAG	C4-C5-C6-O6
7	K	1	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
8	J	3	BMA	C4-C5-C6-O6
8	J	2	NAG	C4-C5-C6-O6


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	J	3	BMA	C1-C2-C3-C4-C5-O5


5 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	G	3	BMA	1	0
7	K	2	NAG	1	0
7	I	1	NAG	1	0
7	K	1	NAG	1	0
6	G	4	BMA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 14 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	SO4	C	506	-	4,4,4	0.25	0	6,6,6	0.05	0
12	NAG	B	2004	2	14,14,15	1.08	2 (14%)	17,19,21	0.88	0
10	SO4	A	505	-	4,4,4	0.24	0	6,6,6	0.05	0
10	SO4	C	505	-	4,4,4	0.23	0	6,6,6	0.08	0
12	NAG	D	2004	2	14,14,15	0.63	0	17,19,21	2.29	3 (17%)
10	SO4	L	301	-	4,4,4	0.24	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	NAG	B	2004	2	-	1/6/23/26	0/1/1/1
12	NAG	D	2004	2	-	3/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	B	2004	NAG	C1-C2	3.09	1.56	1.52
12	B	2004	NAG	C3-C2	2.04	1.56	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	D	2004	NAG	C2-N2-C7	7.98	133.60	122.90
12	D	2004	NAG	C1-C2-N2	3.34	115.69	110.43
12	D	2004	NAG	C3-C4-C5	2.00	113.86	110.23

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
12	D	2004	NAG	C3-C2-N2-C7
12	D	2004	NAG	C8-C7-N2-C2
12	D	2004	NAG	O7-C7-N2-C2
12	B	2004	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	D	2004	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	H	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	H	134:CYS	C	138:THR	N	6.62

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	454/454 (100%)	-0.39	0 100 100	29, 56, 92, 143	5 (1%)
1	C	453/454 (99%)	0.06	2 (0%) 89 90	34, 77, 119, 166	2 (0%)
2	B	466/471 (98%)	0.08	4 (0%) 81 83	36, 91, 170, 226	3 (0%)
2	D	471/471 (100%)	0.70	22 (4%) 37 39	56, 108, 160, 218	5 (1%)
3	E	214/216 (99%)	0.82	12 (5%) 31 33	80, 140, 189, 218	0
3	H	216/216 (100%)	0.33	2 (0%) 81 83	59, 107, 160, 183	0
4	F	214/214 (100%)	0.67	16 (7%) 22 24	87, 140, 210, 268	1 (0%)
4	L	214/214 (100%)	0.32	4 (1%) 66 67	72, 101, 137, 179	0
5	M	5/8 (62%)	0.78	1 (20%) 3 4	83, 87, 101, 130	0
5	N	5/8 (62%)	1.26	0 100 100	105, 106, 127, 157	0
All	All	2712/2726 (99%)	0.25	63 (2%) 61 61	29, 96, 172, 268	16 (0%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	129[A]	TRP	4.7
4	F	132	VAL	4.7
4	F	116	SER	4.2
4	F	133	VAL	3.5
4	F	135	PHE	3.5
4	L	135	PHE	3.3
2	D	142	MET	3.2
2	D	145	LEU	3.2
4	F	181	LEU	3.1
2	D	243	SER	3.0
4	F	117	ILE	3.0
4	F	115	VAL	2.9
2	B	375	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	466	TRP	2.8
3	E	144	LEU	2.8
3	E	130	LEU	2.7
3	E	147	LEU	2.7
3	H	130	LEU	2.7
1	C	453	VAL	2.7
2	D	118	MET	2.6
4	L	105	GLU	2.5
2	D	194	LEU	2.5
3	E	48	ILE	2.5
2	D	116	TYR	2.4
2	D	122	TYR	2.4
2	D	77	SER	2.4
4	L	113	PRO	2.4
3	E	133	VAL	2.3
2	D	331	GLY	2.3
4	L	137	ASN	2.3
3	E	146	CYS	2.3
3	E	124	ALA	2.3
4	F	134	CYS	2.3
4	F	131	SER	2.2
3	E	64	PHE	2.2
4	F	205	ILE	2.2
4	F	209	PHE	2.2
1	C	1	LEU	2.2
2	D	397	PHE	2.2
3	H	214	LYS	2.2
4	F	112	ALA	2.2
4	F	120	PRO	2.1
2	D	136	THR	2.1
2	D	102	ILE	2.1
2	D	200	VAL	2.1
2	D	340	VAL	2.1
3	E	120	ALA	2.1
2	D	375	LEU	2.1
3	E	145	GLY	2.1
5	M	7	CYS	2.1
4	F	179	LEU	2.1
2	B	456	GLU	2.0
3	E	149	LYS	2.0
2	D	151	ILE	2.0
3	E	12	VAL	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	348	TYR	2.0
2	D	471	CYS	2.0
4	F	106	ILE	2.0
2	D	107	VAL	2.0
2	D	379	VAL	2.0
2	B	179	ASP	2.0
2	D	432	ASP	2.0
4	F	186	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	HRG	N	2	12/13	0.86	0.20	99,111,130,130	0
5	HRG	M	2	12/13	0.90	0.20	84,101,118,127	0

6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
12	NAG	D	2004	14/15	0.51	0.14	116,131,136,136	0
10	SO4	C	506	5/5	0.57	0.14	198,201,204,205	0
9	CA	D	2002	1/1	0.68	0.17	194,194,194,194	0
10	SO4	A	505	5/5	0.71	0.14	194,199,206,207	0
10	SO4	L	301	5/5	0.74	0.10	129,133,135,140	0
12	NAG	B	2004	14/15	0.78	0.09	116,138,146,147	0
9	CA	B	2002	1/1	0.83	0.15	181,181,181,181	0
10	SO4	C	505	5/5	0.87	0.14	135,142,144,147	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	CA	A	502	1/1	0.97	0.04	52,52,52,52	0
9	CA	D	2003	1/1	0.97	0.04	72,72,72,72	0
9	CA	C	502	1/1	0.97	0.05	86,86,86,86	0
9	CA	C	504	1/1	0.97	0.05	64,64,64,64	0
9	CA	C	501	1/1	0.98	0.03	114,114,114,114	0
9	CA	A	503	1/1	0.98	0.06	47,47,47,47	0
11	MG	D	2001	1/1	0.98	0.04	95,95,95,95	0
9	CA	C	503	1/1	0.98	0.05	79,79,79,79	0
9	CA	B	2003	1/1	0.98	0.07	61,61,61,61	0
9	CA	A	501	1/1	0.99	0.03	67,67,67,67	0
9	CA	A	504	1/1	0.99	0.03	55,55,55,55	0
11	MG	B	2001	1/1	0.99	0.04	51,51,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.