



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 15, 2022 – 04:06 PM EST

PDB ID : 7THW  
Title : Crystal Structure of the Soluble Domain of the Putative OmpA -Family Membrane Protein YPO0514 from Yersinia pestis  
Authors : Kim, Y.; Tesar, C.; Chhor, G.; Clancy, S.; Babnigg, G.; Schneewind, O.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CS-GID)  
Deposited on : 2022-01-12  
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.25  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.25


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	 4% 75% 8% • 17%
1	B	190	 7% 71% 5% 24%
1	C	190	 5% 76% 5% • 18%
1	D	190	 7% 72% 5% • 23%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	D	402	-	-	X	-

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5021 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative OmpA-family membrane protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	158	1302	834	224	235	3	6	0	0	0
1	B	144	1194	770	203	213	3	5	0	0	0
1	C	155	1276	820	217	230	3	6	0	0	0
1	D	146	1207	778	205	215	3	6	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	132	SER	-	expression tag	UNP A0A454Y5W9
A	133	ASN	-	expression tag	UNP A0A454Y5W9
A	134	ALA	-	expression tag	UNP A0A454Y5W9
B	132	SER	-	expression tag	UNP A0A454Y5W9
B	133	ASN	-	expression tag	UNP A0A454Y5W9
B	134	ALA	-	expression tag	UNP A0A454Y5W9
C	132	SER	-	expression tag	UNP A0A454Y5W9
C	133	ASN	-	expression tag	UNP A0A454Y5W9
C	134	ALA	-	expression tag	UNP A0A454Y5W9
D	132	SER	-	expression tag	UNP A0A454Y5W9
D	133	ASN	-	expression tag	UNP A0A454Y5W9
D	134	ALA	-	expression tag	UNP A0A454Y5W9

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total O P 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	13	Total O 13 13	0	0
4	B	6	Total O 6 6	0	0
4	C	8	Total O 8 8	0	0
4	D	6	Total O 6 6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative OmpA-family membrane protein




- Molecule 1: Putative OmpA-family membrane protein



- Molecule 1: Putative OmpA-family membrane protein



- Molecule 1: Putative OmpA-family membrane protein



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.86Å 123.10Å 54.18Å 90.00° 89.97° 90.00°	Depositor
Resolution (Å)	49.59 – 2.20 49.59 – 2.20	Depositor EDS
% Data completeness (in resolution range)	93.4 (49.59-2.20) 93.5 (49.59-2.20)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.58 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.19_4092	Depositor
R, $R_{free}$	0.226 , 0.269 0.226 , 0.267	Depositor DCC
$R_{free}$ test set	1477 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.2	Xtrriage
Anisotropy	0.422	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 33.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.379 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5021	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/1328	0.46	0/1789
1	B	0.25	0/1217	0.43	0/1637
1	C	0.26	0/1301	0.44	0/1751
1	D	0.25	0/1229	0.42	0/1651
All	All	0.25	0/5075	0.44	0/6828

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1302	0	1260	13	0
1	B	1194	0	1162	6	0
1	C	1276	0	1233	10	0
1	D	1207	0	1176	5	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	D	5	0	0	2	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	13	0	0	0	0
4	B	6	0	0	0	0
4	C	8	0	0	0	0
4	D	6	0	0	0	0
All	All	5021	0	4831	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:TRP:CD2	1:C:149:MSE:HG2	2.34	0.62
1:B:156:GLU:OE2	1:B:204:TYR:OH	2.20	0.58
1:A:152:LEU:O	1:A:156:GLU:HG2	2.09	0.53
1:C:152:LEU:O	1:C:156:GLU:HG2	2.10	0.51
1:A:149:MSE:HG2	1:C:140:TRP:CD2	2.45	0.51
1:C:179:TYR:CG	1:D:175:ARG:HD3	2.45	0.51
1:B:152:LEU:O	1:B:156:GLU:HG2	2.10	0.51
1:A:155:VAL:O	1:A:159:ARG:HG3	2.12	0.50
1:A:179:TYR:CG	1:B:175:ARG:HD3	2.46	0.50
1:A:204:TYR:CZ	1:A:208:ILE:HD11	2.48	0.48
1:D:181:GLN:OE1	1:D:191:GLN:NE2	2.46	0.47
1:A:158:ILE:O	1:A:161:TRP:HB2	2.15	0.47
1:A:281:ASP:O	1:A:285:ARG:HG3	2.15	0.47
1:D:181:GLN:NE2	1:D:185:GLN:OE1	2.48	0.46
1:A:197:LEU:HD21	1:A:253:VAL:HG13	1.96	0.46
1:A:140:TRP:CG	1:C:149:MSE:HG2	2.51	0.45
1:A:156:GLU:HG3	1:C:137:LEU:HD13	1.98	0.45
1:C:161:TRP:CD1	1:C:162:PRO:HD2	2.50	0.45
1:A:179:TYR:CD1	1:B:175:ARG:HD3	2.52	0.45
1:B:151:LEU:HD11	1:B:173:ILE:HG23	1.99	0.44
1:C:158:ILE:O	1:C:161:TRP:HB2	2.19	0.41
1:D:206:ASP:HA	3:D:402:PO4:O1	2.20	0.41
1:D:209:PHE:HB2	3:D:402:PO4:O1	2.21	0.41
1:B:159:ARG:HH21	1:B:208:ILE:HD11	1.85	0.41
1:C:266:ASP:OD1	1:C:266:ASP:N	2.48	0.40
1:A:149:MSE:HG2	1:C:140:TRP:CG	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/190 (81%)	151 (98%)	3 (2%)	0	100	100
1	B	138/190 (73%)	136 (99%)	2 (1%)	0	100	100
1	C	149/190 (78%)	147 (99%)	2 (1%)	0	100	100
1	D	140/190 (74%)	138 (99%)	2 (1%)	0	100	100
All	All	581/760 (76%)	572 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/159 (87%)	136 (99%)	2 (1%)	67	80
1	B	127/159 (80%)	125 (98%)	2 (2%)	62	76
1	C	135/159 (85%)	133 (98%)	2 (2%)	65	78
1	D	128/159 (80%)	123 (96%)	5 (4%)	32	41
All	All	528/636 (83%)	517 (98%)	11 (2%)	53	67

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	161	TRP
1	A	220	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	181	GLN
1	B	271	MSE
1	C	161	TRP
1	C	262	CYS
1	D	137	LEU
1	D	149	MSE
1	D	159	ARG
1	D	181	GLN
1	D	271	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	185	GLN
1	D	181	GLN
1	D	185	GLN
1	D	191	GLN
1	D	195	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	D	402	-	4,4,4	0.93	0	6,6,6	0.56	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	402	PO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	152/190 (80%)	0.45	8 (5%) 26 25	49, 67, 108, 119	0
1	B	139/190 (73%)	0.52	13 (9%) 8 7	51, 76, 102, 114	0
1	C	149/190 (78%)	0.51	10 (6%) 17 16	48, 68, 108, 129	0
1	D	140/190 (73%)	0.51	13 (9%) 8 7	53, 76, 106, 117	0
All	All	580/760 (76%)	0.50	44 (7%) 13 12	48, 71, 107, 129	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	219	ASN	8.5
1	B	209	PHE	5.0
1	D	155	VAL	5.0
1	D	206	ASP	4.7
1	D	140	TRP	4.3
1	B	140	TRP	4.0
1	B	144	TYR	3.9
1	D	208	ILE	3.6
1	D	174	VAL	3.4
1	D	228	ARG	3.4
1	D	158	ILE	3.4
1	B	256	PHE	3.2
1	B	190	PRO	3.2
1	C	155	VAL	3.2
1	C	151	LEU	2.8
1	B	165	ASN	2.8
1	C	169	VAL	2.7
1	B	189	TYR	2.7
1	B	228	ARG	2.7
1	D	253	VAL	2.6
1	D	209	PHE	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	154	LEU	2.5
1	A	144	TYR	2.4
1	C	166	ALA	2.4
1	A	193	ASP	2.4
1	C	222	LEU	2.3
1	D	184	LEU	2.3
1	A	180	PHE	2.3
1	D	292	GLY	2.3
1	A	295	ALA	2.2
1	C	146	ALA	2.2
1	B	184	LEU	2.2
1	B	210	ASN	2.2
1	D	159	ARG	2.2
1	A	152	LEU	2.2
1	C	296	THR	2.1
1	B	222	LEU	2.1
1	D	286	LEU	2.1
1	B	219	ASN	2.1
1	C	161	TRP	2.1
1	A	211	GLY	2.1
1	A	197	LEU	2.1
1	B	197	LEU	2.0
1	C	158	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

*Continued on next page...*

*Continued from previous page...*


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

3	PO4	D	402	5/5	0.80	0.22	105,106,115,169	0
2	CA	B	401	1/1	0.95	0.19	112,112,112,112	0
2	CA	C	401	1/1	0.96	0.10	91,91,91,91	0
2	CA	D	401	1/1	0.97	0.05	99,99,99,99	0
2	CA	A	401	1/1	0.98	0.10	91,91,91,91	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around CA B 401:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around CA C 401:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around CA D 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.