

#### Feb 4, 2023 – 10:30 AM EST

PDB ID	:	7TQT
EMDB ID	:	EMD-26071
Title	:	Coxsackievirus A21 capsid subdomain in complex with mouse polyclonal an-
		tibody pAbC-5
Authors	:	Antanasijevic, A.; Ward, A.B.
Deposited on	:	2022-01-27
Resolution	:	4.10  Å(reported)
Based on initial model	:	1Z7S

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.32.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\#Entries)$	${f EM} {f structures} \ (\#{f Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
			49%	
1	L	98	95%	5%
_			30%	
2	Н	108	98%	•
_			40%	
3	a	298	90%	5% 6%
		200	34%	
3	е	298	91%	• 5%
		200	45%	
3	1	298	90%	5% 5%
		200	44%	
3	m	298	86%	6% 7%
0		200	43%	
3	q	298	90%	5% 5%
4	1	979	37%	
4	b	272	93%	• •



Mol	Chain	Length	Quality of chain	
4	f	272	34%	• 5%
4	j	272	48% 91%	• 6%
4	n	272	94%	• •
4	r	272	91%	• 7%
5	с	240	99%	•
5	g	240	95%	••
5	k	240	95%	••
5	О	240	95%	••
5	s	240	93%	5% •
6	d	69	96%	••
6	h	69	93%	6% •
6	1	69	94%	• •
6	р	69	70% 94%	••
6	$\mathbf{t}$	69	<u>68%</u> 93%	6% •



# 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 33877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called pAbC-5 light chain.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	L	98	Total 490	C 294	N 98	O 98	0	0

• Molecule 2 is a protein called pAbC-5 heavy chain.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	Н	108	Total	C 204	N 100	0	0	0
			540	324	108	108		

• Molecule 3 is a protein called VP1.

Mol	Chain	Residues		At	oms			AltConf	Trace
2	0	201	Total	С	Ν	Ο	$\mathbf{S}$	0	0
0	a	201	2213	1402	378	423	10	0	0
3	0	282	Total	С	Ν	Ο	$\mathbf{S}$	0	0
0	е	282	2219	1405	379	425	10	0	0
3	i	<u> </u>	Total	С	Ν	0	$\mathbf{S}$	0	0
0	1	200	2226	1410	380	426	10	0	0
2	3 m	276	Total	С	Ν	0	S	0	0
0			2176	1382	371	413	10	0	0
2	9	263	Total	С	Ν	0	S	0	0
5	q	200	2226	1410	380	426	10	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	290	ALA	THR	conflict	UNP Q7T7N6
e	290	ALA	THR	conflict	UNP Q7T7N6
i	290	ALA	THR	conflict	UNP Q7T7N6
m	290	ALA	THR	conflict	UNP Q7T7N6
q	290	ALA	THR	conflict	UNP Q7T7N6

• Molecule 4 is a protein called VP2.



Mol	Chain	Residues		At	oms			AltConf	Trace
4	h	264	Total	С	Ν	0	$\mathbf{S}$	0	0
4	D	204	2053	1306	346	390	11	0	0
4	f	258	Total	С	Ν	Ο	S	0	0
4	1	200	2008	1279	339	379	11	0	0
4	i	257	Total	С	Ν	Ο	$\mathbf{S}$	0	0
4	J		2003	1276	339	377	11	0	0
4	n	261	Total	С	Ν	0	S	0	0
4	± 11	201	2029	1290	343	385	11	0	0
4	r	252	Total	С	Ν	0	S	0	0
±		200	1975	1261	333	370	11	0	

• Molecule 5 is a protein called VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	0	220	Total	С	Ν	0	$\mathbf{S}$	0	0
0	C	239	1848	1178	307	343	20	0	0
5	ď	234	Total	С	Ν	Ο	$\mathbf{S}$	0	0
0	g	204	1810	1154	299	337	20	0	0
5	ŀ	033	Total	С	Ν	Ο	$\mathbf{S}$	0	0
0	K	200	1801	1148	297	336	20	0	0
5	0	224	Total	С	Ν	0	S	0	0
0	0	234	1810	1154	299	337	20	0	0
5	G	225	Total	С	Ν	0	S	0	0
0	G	235	1818	1160	300	338	20	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
с	225	ARG	LYS	conflict	UNP Q7T7N6
g	225	ARG	LYS	conflict	UNP Q7T7N6
k	225	ARG	LYS	conflict	UNP Q7T7N6
0	225	ARG	LYS	conflict	UNP Q7T7N6
s	225	ARG	LYS	conflict	UNP Q7T7N6

• Molecule 6 is a protein called VP4.

Mol	Chain	Residues	Atoms				AltConf	Trace	
6	d	68	Total	С	Ν	0	S	0	0
0 0	08	512	311	90	110	1	0	0	
6	h	68	Total	С	Ν	0	S	0	0
0	11	08	512	311	90	110	1		0
6	1	69	Total	С	Ν	0	S	0	0
0	1	00	512	311	90	110	1	U	U



Mol	Chain	Residues	Atoms				AltConf	Trace	
6	n	67	Total	С	Ν	0	S	0	0
0	o p	07	504	307	88	108	1	0	0
6	+	68	Total	С	Ν	0	S	0	0
0	U	t 08	512	311	90	110	1	0	0



Mol	Chain	Residues	Atoms	AltConf
7	0	1	Total C O	0
1	a	1	16  14  2	0
7	0	1	Total C O	0
1	C	T	16  14  2	0
7	i	1	Total C O	0
'	1	1	16  14  2	0
7	m	1	Total C O	0
'	111	1	16  14  2	0
7	a	1	Total C O	0
'	Ч		16  14  2	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: pAbC-5 light chain



• Molecule 3: VP1



L D W I D E

• Molecule 3: VP1



















# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	12654	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.019	Depositor
Minimum map value	-0.013	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.007	Depositor
Map size (Å)	292.59998, 292.59998, 292.59998	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.045, 1.045, 1.045	Depositor



# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	ond lengths	Bond angles		
	Ullaili	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
3	a	1.20	9/2275~(0.4%)	1.00	7/3102~(0.2%)	
3	е	1.17	8/2281~(0.4%)	0.98	5/3111~(0.2%)	
3	i	1.18	7/2288~(0.3%)	1.07	12/3121~(0.4%)	
3	m	1.22	17/2237~(0.8%)	1.02	7/3048~(0.2%)	
3	q	1.18	10/2288~(0.4%)	1.00	9/3121~(0.3%)	
4	b	1.21	10/2110~(0.5%)	0.97	7/2887~(0.2%)	
4	f	1.07	4/2063~(0.2%)	0.95	5/2821~(0.2%)	
4	j	1.09	5/2057~(0.2%)	0.97	5/2811~(0.2%)	
4	n	1.07	5/2084~(0.2%)	0.91	2/2850~(0.1%)	
4	r	1.10	5/2029~(0.2%)	0.91	1/2773~(0.0%)	
5	с	1.00	0/1897	0.93	3/2582~(0.1%)	
5	g	1.11	5/1859~(0.3%)	0.93	3/2531~(0.1%)	
5	k	1.06	0/1850	0.96	6/2520~(0.2%)	
5	0	1.07	2/1859~(0.1%)	0.99	7/2531~(0.3%)	
5	s	1.06	3/1867~(0.2%)	1.01	11/2542~(0.4%)	
6	d	0.99	1/519~(0.2%)	0.95	2/705~(0.3%)	
6	h	0.97	2/519~(0.4%)	0.94	2/705~(0.3%)	
6	1	1.06	0/519	0.97	3/704~(0.4%)	
6	р	0.99	1/511~(0.2%)	0.83	1/694~(0.1%)	
6	t	1.06	3/519~(0.6%)	0.93	3/705~(0.4%)	
All	All	1.12	97/33631~(0.3%)	0.97	101/45864~(0.2%)	

All (97) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	b	85	CYS	CB-SG	-10.02	1.65	1.82
4	b	224	HIS	CB-CG	-8.55	1.34	1.50
6	t	14	GLU	CG-CD	-8.02	1.40	1.51
4	b	129	GLU	CG-CD	-7.67	1.40	1.51
4	b	9	TYR	CG-CD2	7.66	1.49	1.39
4	b	9	TYR	CE1-CZ	7.64	1.48	1.38



001000	nucu jion	v proov	o uo pugo	• • •			
Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	a	137	TYR	CB-CG	-7.55	1.40	1.51
5	0	62	VAL	CB-CG2	-7.51	1.37	1.52
3	m	298	PHE	CG-CD1	7.51	1.50	1.38
4	b	9	TYR	CG-CD1	7.49	1.48	1.39
4	r	79	TRP	CB-CG	-7.45	1.36	1.50
3	q	154	TYR	CB-CG	-7.31	1.40	1.51
4	f	129	GLU	CG-CD	-7.21	1.41	1.51
3	m	264	TRP	CB-CG	-7.20	1.37	1.50
3	a	298	PHE	CG-CD2	7.14	1.49	1.38
4	f	129	GLU	CD-OE2	-6.97	1.18	1.25
3	a	298	PHE	CG-CD1	6.95	1.49	1.38
3	i	298	PHE	CG-CD1	6.79	1.49	1.38
3	е	298	PHE	CG-CD1	6.77	1.49	1.38
5	g	105	TYR	CB-CG	-6.74	1.41	1.51
3	m	154	TYR	CD1-CE1	-6.74	1.29	1.39
3	m	182	TYR	CB-CG	-6.74	1.41	1.51
3	i	255	TYR	CB-CG	-6.71	1.41	1.51
4	r	210	VAL	CB-CG2	-6.66	1.38	1.52
3	q	255	TYR	CB-CG	-6.65	1.41	1.51
3	q	298	PHE	CG-CD1	6.62	1.48	1.38
3	е	298	PHE	CG-CD2	6.56	1.48	1.38
3	i	298	PHE	CG-CD2	6.46	1.48	1.38
4	j	179	TYR	CB-CG	-6.46	1.42	1.51
3	m	298	PHE	CG-CD2	6.41	1.48	1.38
3	i	144	GLU	CD-OE1	-6.38	1.18	1.25
4	b	9	TYR	CE2-CZ	6.34	1.46	1.38
4	r	142	TYR	CB-CG	-6.26	1.42	1.51
4	n	142	TYR	CB-CG	-6.21	1.42	1.51
3	m	154	TYR	CB-CG	-6.20	1.42	1.51
4	j	129	GLU	CG-CD	-6.06	1.42	1.51
6	t	14	GLU	CD-OE1	-5.99	1.19	1.25
3	a	255	TYR	CB-CG	-5.96	1.42	1.51
5	g	105	TYR	CD1-CE1	-5.96	1.30	1.39
5	g	105	TYR	CG-CD1	-5.96	1.31	1.39
3	m	298	PHE	CE2-CZ	5.91	1.48	1.37
3	m	43	GLU	CG-CD	-5.88	1.43	1.51
3	е	122	TYR	CB-CG	-5.88	1.42	1.51
3	q	100	PHE	CB-CG	-5.86	1.41	1.51
6	t	14	GLU	CD-OE2	-5.80	1.19	1.25
5	0	193	TYR	CB-CG	-5.79	1.43	1.51
4	r	257	CYS	CB-SG	-5.78	1.72	1.81
4	n	260	PHE	CB-CG	-5.77	1.41	1.51



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	i	69	GLU	CD-OE2	-5.74	1.19	1.25
4	n	129	GLU	CD-OE2	-5.72	1.19	1.25
3	е	69	GLU	CG-CD	-5.64	1.43	1.51
3	q	154	TYR	CD1-CE1	-5.64	1.30	1.39
3	a	120	PHE	CB-CG	-5.55	1.42	1.51
4	n	129	GLU	CG-CD	-5.54	1.43	1.51
3	m	168	TYR	CB-CG	-5.54	1.43	1.51
3	i	137	TYR	CB-CG	-5.53	1.43	1.51
3	е	128	GLU	CD-OE1	-5.51	1.19	1.25
3	m	43	GLU	CD-OE1	-5.51	1.19	1.25
3	a	210	VAL	CB-CG2	-5.50	1.41	1.52
3	a	43	GLU	CD-OE2	-5.47	1.19	1.25
4	b	103	ARG	CD-NE	-5.45	1.37	1.46
3	m	243	SER	CB-OG	-5.45	1.35	1.42
5	g	185	GLU	CD-OE2	-5.41	1.19	1.25
3	m	150	TYR	CB-CG	-5.39	1.43	1.51
3	е	149	VAL	CB-CG1	-5.38	1.41	1.52
4	b	239	PHE	CB-CG	-5.34	1.42	1.51
3	е	298	PHE	CE1-CZ	5.34	1.47	1.37
3	q	238	ARG	CD-NE	-5.31	1.37	1.46
3	m	100	PHE	CB-CG	-5.28	1.42	1.51
3	m	69	GLU	CD-OE1	-5.27	1.19	1.25
4	n	134	CYS	CB-SG	-5.27	1.73	1.81
3	m	128	GLU	CD-OE2	-5.23	1.19	1.25
3	q	234	VAL	CB-CG2	-5.22	1.41	1.52
4	r	259	GLU	CD-OE1	-5.20	1.20	1.25
3	a	128	GLU	CD-OE2	-5.19	1.20	1.25
4	j	174	PHE	CB-CG	-5.18	1.42	1.51
4	f	37	GLU	CD-OE1	-5.16	1.20	1.25
5	g	82	PHE	CB-CG	-5.16	1.42	1.51
6	h	36	SER	CB-OG	-5.16	1.35	1.42
3	q	144	GLU	CD-OE2	-5.16	1.20	1.25
4	f	259	GLU	CD-OE1	-5.15	1.20	1.25
3	i	298	PHE	CE1-CZ	5.15	1.47	1.37
3	a	277	GLU	CD-OE2	-5.13	1.20	1.25
4	j	37	GLU	CD-OE1	-5.12	1.20	1.25
5	s	67	MET	CG-SD	-5.11	1.67	1.81
3	q	128	GLU	CD-OE1	-5.11	1.20	1.25
4	j	259	GLU	CD-OE2	-5.11	1.20	1.25
6	р	14	GLU	CD-OE1	-5.05	1.20	1.25
3	m	213	GLU	CD-OE1	-5.04	1.20	1.25
6	h	14	GLU	CD-OE1	-5.03	1.20	1.25



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	m	96	GLU	CD-OE1	-5.03	1.20	1.25
4	b	103	ARG	CG-CD	-5.03	1.39	1.51
5	s	192	PHE	CB-CG	-5.02	1.42	1.51
6	d	32	TYR	CB-CG	-5.02	1.44	1.51
3	е	64	TYR	CB-CG	-5.01	1.44	1.51
3	q	298	PHE	CD1-CE1	5.00	1.49	1.39
5	s	66	GLU	CD-OE2	-5.00	1.20	1.25

All (101) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
4	f	103	ARG	NE-CZ-NH2	-10.80	114.90	120.30
5	s	225	ARG	NE-CZ-NH2	-9.84	115.38	120.30
3	m	154	TYR	CB-CG-CD1	-9.36	115.38	121.00
5	0	225	ARG	NE-CZ-NH2	-9.32	115.64	120.30
4	j	12	ARG	NE-CZ-NH2	-9.21	115.69	120.30
3	q	154	TYR	CB-CG-CD1	-9.17	115.50	121.00
3	a	114	ARG	NE-CZ-NH2	-9.05	115.77	120.30
3	i	124	ARG	NE-CZ-NH2	-9.00	115.80	120.30
3	i	124	ARG	NE-CZ-NH1	8.87	124.73	120.30
3	i	182	TYR	CB-CG-CD2	-8.75	115.75	121.00
5	0	222	ARG	NE-CZ-NH2	-8.62	115.99	120.30
3	i	115	ARG	NE-CZ-NH2	-8.59	116.00	120.30
3	е	146	ARG	NE-CZ-NH2	-8.54	116.03	120.30
4	f	172	ARG	NE-CZ-NH2	-8.36	116.12	120.30
3	i	146	ARG	NE-CZ-NH2	-8.35	116.12	120.30
3	i	253	ARG	NE-CZ-NH1	-8.25	116.17	120.30
5	0	225	ARG	NE-CZ-NH1	8.06	124.33	120.30
4	r	172	ARG	NE-CZ-NH2	-7.80	116.40	120.30
5	s	177	ARG	NE-CZ-NH2	-7.70	116.45	120.30
4	j	100	TYR	CB-CG-CD2	-7.62	116.42	121.00
3	m	114	ARG	NE-CZ-NH2	-7.60	116.50	120.30
5	k	222	ARG	NE-CZ-NH1	7.24	123.92	120.30
4	b	103	ARG	NE-CZ-NH1	7.13	123.87	120.30
6	d	35	ASP	CB-CG-OD1	7.07	124.66	118.30
4	f	103	ARG	NE-CZ-NH1	7.03	123.82	120.30
3	i	67	ARG	NE-CZ-NH1	6.96	123.78	120.30
3	i	115	ARG	NE-CZ-NH1	6.95	123.77	120.30
3	i	114	ARG	NE-CZ-NH2	-6.92	116.84	120.30
3	q	67	ARG	NE-CZ-NH2	-6.92	116.84	120.30
4	j	172	ARG	NE-CZ-NH2	-6.80	116.90	120.30
5	g	105	TYR	CB-CG-CD1	-6.75	116.95	121.00



$\alpha$ $\cdot \cdot$ $\cdot$	C		
Continued	trom	previous	paae
0 0 1 0 0 1 0 0 0 0 0 0	J	r · · · · · · · · · · · · · · · · · · ·	r ~g ~···

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	0	93	ARG	NE-CZ-NH2	-6.67	116.96	120.30
3	q	67	ARG	NE-CZ-NH1	6.66	123.63	120.30
6	1	32	TYR	CB-CG-CD1	-6.64	117.01	121.00
5	с	114	ARG	NE-CZ-NH2	-6.61	117.00	120.30
3	i	238	ARG	NE-CZ-NH2	-6.58	117.01	120.30
5	0	222	ARG	NE-CZ-NH1	6.55	123.58	120.30
5	s	114	ARG	NE-CZ-NH2	-6.46	117.07	120.30
3	q	188	ARG	NE-CZ-NH2	-6.40	117.10	120.30
6	h	35	ASP	CB-CG-OD1	6.38	124.04	118.30
4	f	212	TYR	CB-CG-CD1	-6.33	117.20	121.00
3	i	274	TYR	CB-CA-C	-6.21	97.98	110.40
4	b	201	ARG	NE-CZ-NH1	6.20	123.40	120.30
3	a	122	TYR	CB-CG-CD2	-6.18	117.29	121.00
5	s	93	ARG	NE-CZ-NH2	-6.18	117.21	120.30
5	k	93	ARG	NE-CZ-NH1	6.17	123.39	120.30
6	1	35	ASP	CB-CG-OD1	6.15	123.84	118.30
3	m	67	ARG	NE-CZ-NH1	6.09	123.34	120.30
4	f	145	TYR	CB-CG-CD1	-6.01	117.39	121.00
4	b	264	ARG	NE-CZ-NH2	-5.96	117.32	120.30
3	a	124	ARG	NE-CZ-NH1	5.95	123.28	120.30
5	s	225	ARG	NE-CZ-NH1	5.95	123.28	120.30
5	g	93	ARG	NE-CZ-NH2	-5.91	117.34	120.30
5	k	193	TYR	CB-CG-CD2	-5.89	117.47	121.00
6	l	59	ASP	CB-CG-OD1	5.88	123.59	118.30
6	р	32	TYR	CB-CG-CD1	-5.86	117.48	121.00
5	с	222	ARG	CB-CA-C	-5.79	98.82	110.40
5	0	93	ARG	NE-CZ-NH1	5.78	123.19	120.30
3	m	137	TYR	CB-CG-CD2	-5.77	117.54	121.00
4	b	264	ARG	NE-CZ-NH1	5.74	123.17	120.30
3	е	67	ARG	NE-CZ-NH1	5.73	123.16	120.30
4	j	145	TYR	CB-CG-CD1	-5.69	117.59	121.00
4	b	98	TYR	CB-CG-CD2	-5.61	117.63	121.00
5	S	222	ARG	NE-CZ-NH2	-5.61	117.50	120.30
3	m	124	ARG	NE-CZ-NH1	5.60	123.10	120.30
3	a	238	ARG	NE-CZ-NH2	-5.59	117.50	120.30
3	q	200	TYR	CB-CG-CD1	-5.58	117.65	121.00
3	е	200	TYR	CB-CG-CD1	-5.57	117.66	121.00
3	q	262	ARG	NE-CZ-NH2	-5.57	117.51	120.30
5	k	177	ARG	NE-CZ-NH2	-5.56	117.52	120.30
3	a	275	ARG	NE-CZ-NH1	5.52	123.06	120.30
6	t	32	TYR	CB-CG-CD1	-5.50	117.70	121.00
3	q	267	ARG	NE-CZ-NH2	-5.48	117.56	120.30



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	g	114	ARG	NE-CZ-NH2	-5.47	117.57	120.30
5	k	177	ARG	NE-CZ-NH1	5.46	123.03	120.30
5	0	114	ARG	NE-CZ-NH2	-5.46	117.57	120.30
3	е	124	ARG	NE-CZ-NH2	5.45	123.03	120.30
3	е	146	ARG	NE-CZ-NH1	5.44	123.02	120.30
3	q	107	TYR	CB-CA-C	5.44	121.28	110.40
3	a	137	TYR	CB-CG-CD1	-5.43	117.74	121.00
3	m	270	ARG	NE-CZ-NH1	5.43	123.01	120.30
5	s	192	PHE	CB-CG-CD2	-5.41	117.02	120.80
4	b	142	TYR	CB-CG-CD2	-5.37	117.78	121.00
5	s	144	ARG	NE-CZ-NH2	-5.34	117.63	120.30
5	s	181	ASP	CB-CG-OD1	5.33	123.10	118.30
6	h	32	TYR	CB-CG-CD1	-5.32	117.81	121.00
3	i	107	TYR	CB-CG-CD2	-5.28	117.83	121.00
4	n	98	TYR	CB-CG-CD1	-5.27	117.84	121.00
4	n	12	ARG	NE-CZ-NH2	-5.24	117.68	120.30
6	t	43	ARG	NE-CZ-NH1	5.23	122.92	120.30
3	m	146	ARG	NE-CZ-NH2	-5.21	117.70	120.30
6	t	59	ASP	CB-CG-OD1	5.19	122.97	118.30
3	q	146	ARG	NE-CZ-NH2	-5.18	117.71	120.30
5	s	114	ARG	NE-CZ-NH1	5.18	122.89	120.30
3	a	115	ARG	NE-CZ-NH1	5.17	122.89	120.30
6	d	32	TYR	CB-CG-CD1	-5.17	117.90	121.00
5	k	93	ARG	NE-CZ-NH2	-5.15	117.73	120.30
4	j	12	ARG	NE-CZ-NH1	5.11	122.86	120.30
5	с	114	ARG	NE-CZ-NH1	5.08	122.84	120.30
5	s	91	ASP	CB-CG-OD2	5.03	122.83	118.30
4	b	201	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	490	0	118	3	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Н	540	0	130	1	0
3	a	2213	0	2132	0	0
3	е	2219	0	2137	0	0
3	i	2226	0	2146	0	0
3	m	2176	0	2102	0	0
3	q	2226	0	2146	0	0
4	b	2053	0	1975	0	0
4	f	2008	0	1936	0	0
4	j	2003	0	1931	0	0
4	n	2029	0	1953	0	0
4	r	1975	0	1907	0	0
5	с	1848	0	1836	0	0
5	g	1810	0	1791	0	0
5	k	1801	0	1778	0	0
5	0	1810	0	1791	0	0
5	s	1818	0	1802	0	0
6	d	512	0	493	0	0
6	h	512	0	493	0	0
6	1	512	0	493	0	0
6	р	504	0	487	0	0
6	t	512	0	493	0	0
7	a	16	0	27	0	0
7	е	16	0	27	0	0
7	i	16	0	27	0	0
7	m	16	0	27	0	0
7	q	16	0	27	0	0
All	All	33877	0	32205	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:5:UNK:O	1:L:6:UNK:C	2.54	0.52
1:L:55:UNK:O	1:L:58:UNK:N	2.44	0.50
1:L:55:UNK:O	1:L:56:UNK:C	2.66	0.43
2:H:17:UNK:O	2:H:18:UNK:C	2.70	0.40

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	a	279/298~(94%)	274 (98%)	5 (2%)	0	100 100
3	е	280/298~(94%)	273~(98%)	7 (2%)	0	100 100
3	i	281/298~(94%)	276 (98%)	5 (2%)	0	100 100
3	m	272/298~(91%)	267~(98%)	5 (2%)	0	100 100
3	q	281/298~(94%)	276 (98%)	5 (2%)	0	100 100
4	b	262/272~(96%)	256~(98%)	6(2%)	0	100 100
4	f	254/272~(93%)	247 (97%)	7(3%)	0	100 100
4	j	251/272~(92%)	247 (98%)	4 (2%)	0	100 100
4	n	257/272~(94%)	252 (98%)	5 (2%)	0	100 100
4	r	247/272~(91%)	241 (98%)	6 (2%)	0	100 100
5	с	237/240~(99%)	236 (100%)	1 (0%)	0	100 100
5	g	232/240~(97%)	225 (97%)	7 (3%)	0	100 100
5	k	231/240~(96%)	228 (99%)	3 (1%)	0	100 100
5	О	232/240~(97%)	231 (100%)	1 (0%)	0	100 100
5	S	233/240~(97%)	227 (97%)	6 (3%)	0	100 100
6	d	66/69~(96%)	64 (97%)	2 (3%)	0	100 100
6	h	66/69~(96%)	64 (97%)	2 (3%)	0	100 100
6	1	66/69~(96%)	65~(98%)	1 (2%)	0	100 100
6	р	65/69~(94%)	61 (94%)	4 (6%)	0	100 100
6	t	66/69~(96%)	66 (100%)	0	0	100 100
All	All	4158/4395 (95%)	4076 (98%)	82 (2%)	0	100 100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
3	a	246/260~(95%)	246 (100%)	0	100	100
3	е	247/260~(95%)	247 (100%)	0	100	100
3	i	248/260~(95%)	248 (100%)	0	100	100
3	m	242/260~(93%)	242 (100%)	0	100	100
3	q	248/260~(95%)	248 (100%)	0	100	100
4	b	224/230~(97%)	224 (100%)	0	100	100
4	f	219/230~(95%)	219 (100%)	0	100	100
4	j	219/230~(95%)	218 (100%)	1 (0%)	88	93
4	n	222/230~(96%)	222 (100%)	0	100	100
4	r	216/230~(94%)	216 (100%)	0	100	100
5	с	212/213~(100%)	212 (100%)	0	100	100
5	g	208/213~(98%)	208 (100%)	0	100	100
5	k	207/213~(97%)	207 (100%)	0	100	100
5	О	208/213~(98%)	208 (100%)	0	100	100
5	$\mathbf{S}$	209/213~(98%)	209 (100%)	0	100	100
6	d	57/58~(98%)	57~(100%)	0	100	100
6	h	57/58~(98%)	57~(100%)	0	100	100
6	1	57/58~(98%)	57 (100%)	0	100	100
6	р	56/58~(97%)	56 (100%)	0	100	100
6	t	57/58~(98%)	57 (100%)	0	100	100
All	All	3659/3805~(96%)	3658 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	j	264	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such



sidechains are listed below:

Mol	Chain	Res	Type
4	f	226	ASN
3	m	99	HIS
4	n	111	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tink	Bond lengths			Bond angles		
moi Type	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
7	MYR	m	300	-	15,15,15	1.47	2 (13%)	15,15,15	0.79	0
7	MYR	a	300	-	15,15,15	1.39	2 (13%)	15,15,15	0.99	2 (13%)
7	MYR	q	300	-	15,15,15	1.38	2 (13%)	15,15,15	0.92	1 (6%)
7	MYR	i	300	-	15,15,15	1.40	2 (13%)	15,15,15	0.96	2 (13%)
7	MYR	е	300	-	15,15,15	1.39	2 (13%)	15,15,15	0.77	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



Mol	Type	Chain	$\operatorname{Res}$	Link	Chirals	Torsions	Rings
7	MYR	m	300	-	-	6/13/13/13	-
7	MYR	a	300	-	-	4/13/13/13	-
7	MYR	q	300	-	-	5/13/13/13	-
7	MYR	i	300	-	-	6/13/13/13	-
7	MYR	е	300	-	-	4/13/13/13	_

'-' means no outliers of that kind were identified.

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
7	е	300	MYR	O2-C1	-3.33	1.19	1.30
7	m	300	MYR	O1-C1	3.33	1.33	1.22
7	q	300	MYR	O1-C1	3.30	1.33	1.22
7	i	300	MYR	O2-C1	-3.28	1.19	1.30
7	m	300	MYR	O2-C1	-3.16	1.20	1.30
7	а	300	MYR	O1-C1	3.05	1.32	1.22
7	a	300	MYR	O2-C1	-3.03	1.20	1.30
7	е	300	MYR	O1-C1	3.00	1.32	1.22
7	q	300	MYR	O2-C1	-2.93	1.20	1.30
7	i	300	MYR	01-C1	2.92	1.31	1.22

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
7	i	300	MYR	O2-C1-O1	-2.66	116.68	123.30
7	q	300	MYR	O2-C1-C2	2.62	122.45	114.03
7	a	300	MYR	O2-C1-C2	2.53	122.15	114.03
7	a	300	MYR	O2-C1-O1	-2.44	117.22	123.30
7	i	300	MYR	O2-C1-C2	2.26	121.31	114.03
7	е	300	MYR	O2-C1-O1	-2.26	117.67	123.30

There are no chirality outliers.

All $(25)$	torsion	outliers	are	listed	below:
------------	---------	----------	-----	--------	--------

Mol	Chain	Res	Type	Atoms
7	а	300	MYR	C3-C4-C5-C6
7	i	300	MYR	C4-C5-C6-C7
7	i	300	MYR	C10-C11-C12-C13
7	е	300	MYR	C11-C10-C9-C8
7	е	300	MYR	C5-C6-C7-C8



Mol	Chain	Res	Type	Atoms
7	m	300	MYR	C6-C7-C8-C9
7	q	300	MYR	C11-C10-C9-C8
7	е	300	MYR	C7-C8-C9-C10
7	q	300	MYR	C4-C5-C6-C7
7	m	300	MYR	C11-C12-C13-C14
7	m	300	MYR	O2-C1-C2-C3
7	q	300	MYR	C5-C6-C7-C8
7	а	300	MYR	O1-C1-C2-C3
7	m	300	MYR	O1-C1-C2-C3
7	a	300	MYR	O2-C1-C2-C3
7	i	300	MYR	O2-C1-C2-C3
7	i	300	MYR	O1-C1-C2-C3
7	i	300	MYR	C3-C4-C5-C6
7	a	300	MYR	C11-C10-C9-C8
7	е	300	MYR	C11-C12-C13-C14
7	q	300	MYR	O1-C1-C2-C3
7	m	300	MYR	C5-C6-C7-C8
7	m	300	MYR	C7-C8-C9-C10
7	i	300	MYR	C5-C6-C7-C8
7	q	300	MYR	O2-C1-C2-C3

Continued from previous page...

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-26071. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



6.1.2 Raw map



The images above show the map projected in three orthogonal directions.



### 6.2 Central slices (i)

### 6.2.1 Primary map









Z Index: 140

#### 6.2.2 Raw map



X Index: 140

Y Index: 140



The images above show central slices of the map in three orthogonal directions.



### 6.3 Largest variance slices (i)

### 6.3.1 Primary map



X Index: 125





Z Index: 153

#### 6.3.2 Raw map



X Index: 145

Y Index: 144



The images above show the largest variance slices of the map in three orthogonal directions.



### 6.4 Orthogonal surface views (i)

#### 6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.007. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.



#### Mask visualisation (i) 6.5

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

#### $emd_{26071}_{msk_{1.map}}$ 6.5.1



Υ



# 7 Map analysis (i)

This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 7.2 Volume estimate (i)



The volume at the recommended contour level is  $257 \text{ nm}^3$ ; this corresponds to an approximate mass of 232 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum (i)



\*Reported resolution corresponds to spatial frequency of 0.244  $\rm \AA^{-1}$ 



# 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

#### 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.244  $\mathrm{\AA^{-1}}$ 



### 8.2 Resolution estimates (i)

$\begin{bmatrix} Bosolution ostimato (Å) \end{bmatrix}$	Estimation criterion (FSC cut-off)			
resolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	4.10	-	-	
Author-provided FSC curve	4.10	5.78	4.18	
Unmasked-calculated*	4.53	7.46	4.74	

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.53 differs from the reported value 4.1 by more than 10 %



# 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-26071 and PDB model 7TQT. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.007 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

#### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.007).



### 9.4 Atom inclusion (i)



At the recommended contour level, 64% of all backbone atoms, 43% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

### 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.007) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.4301	0.4030
Н	0.6241	0.3120
L	0.5000	0.3340
a	0.4320	0.4050
b	0.4462	0.4150
с	0.4607	0.4090
d	0.3393	0.4170
е	0.4432	0.4120
f	0.4515	0.4180
g	0.4678	0.4080
h	0.3097	0.4040
i	0.4057	0.4020
j	0.4049	0.3980
k	0.4448	0.4060
1	0.2998	0.3970
m	0.4151	0.4020
n	0.4071	0.3930
0	0.4285	0.3970
р	0.2826	0.3930
q	0.4157	0.3980
r	0.4814	0.4170
S	0.4489	0.4070
t	0.3018	0.3970

