

Full wwPDB X-ray Structure Validation Report (i)

Sep 28, 2024 – 01:08 PM EDT

PDB ID : 7TWL	
Title : Structure of a borosin methyltransferase from Mycena rosella with	peptide A2
(MroMA2) in complex with SAH	
Authors : Zheng, Y.; Ongpipattanakul, C.; Nair, S.K.	
Deposited on : $2022-02-07$	
Resolution : $1.86 \text{ Å}(\text{reported})$	

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
wietric	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359 (1.86 - 1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	400	2% 8 6%	10%	••
1	В	400	2% 8 7%	8%	•••
1	С	400	<u>6%</u> 84%	12%	•••
1	D	400	87%	9%	•••



$7\mathrm{TWL}$

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 12984 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	Δ	200	Total	С	Ν	0	\mathbf{S}	0	2	0
1	A	300	3012	1927	509	563	13	0	5	
1	р	200	Total	С	Ν	0	S	0	1	0
1	D	300	3029	1936	518	562	13	0	4	0
1	C	297	Total	С	Ν	0	S	0	F	0
1		301	3015	1930	511	561	13	0	5	0
1	П	280	Total	С	Ν	0	S	0	9	0
		309	3009	1923	509	564	13	0		

• Molecule 1 is a protein called MroMA2.

• Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
0	Δ	1	Total	С	Ν	0	S	0	0
	A	L	26	14	6	5	1	0	0
0	D	1	Total	С	Ν	0	S	0	0
	D		26	14	6	5	1	0	



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
0	C	1	Total	С	Ν	0	S	0	0	
	U	L	26	14	6	5	1	0		
0	Л	1	Total	С	Ν	0	S	0	0	
	D	L	26	14	6	5	1	0	U	

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	224	Total O 224 224	0	0
3	В	236	Total O 236 236	0	0
3	С	194	Total O 194 194	0	0
3	D	161	Total O 161 161	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: MroMA2



V234 SER 1235 ALM 1235 ALM 2242 ALM 2266 LVS 7270 LVS 7289 DS 7387 DS 7387 DS 7387 LIS 7389 DS 7389 DS 741 DS 741 DS 743 LIS 743 VIS 743 VIS 743 VIS 745 VIS 745</



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	69.11Å 125.23Å 203.31Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution(A)	25.00 - 1.86	Depositor
Resolution (A)	25.00 - 1.86	EDS
% Data completeness	99.9 (25.00-1.86)	Depositor
(in resolution range)	99.9 (25.00-1.86)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.27 (at 1.86 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
B B.	0.188 , 0.223	Depositor
II, II, <i>free</i>	0.191 , 0.225	DCC
R_{free} test set	7287 reflections $(4.91%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	28.7	Xtriage
Anisotropy	0.262	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34, 35.6	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12984	wwPDB-VP
Average B, all atoms $(Å^2)$	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MVA, SAH, MLE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.32	0/3068	0.61	0/4162	
1	В	0.33	0/3088	0.61	0/4186	
1	С	0.31	0/3077	0.60	0/4173	
1	D	0.29	0/3062	0.58	0/4154	
All	All	0.31	0/12295	0.60	0/16675	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3012	0	3029	29	0
1	В	3029	0	3057	35	0
1	С	3015	0	3037	45	0
1	D	3009	0	3018	31	0
2	А	26	0	19	0	0
2	В	26	0	19	0	0
2	С	26	0	19	0	0
2	D	26	0	19	0	0
3	A	224	0	0	2	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes						
3	В	236	0	0	8	0						
3	С	194	0	0	5	0						
3	D	161	0	0	7	0						
All	All	12984	0	12217	119	0						

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:273:GLN:HG3	3:B:778:HOH:O	1.47	1.11
1:C:273:GLN:HG3	3:C:676:HOH:O	1.52	1.06
1:C:357:LYS:HD2	1:C:358:PRO:HD2	1.38	1.01
1:B:250:THR:HB	3:B:743:HOH:O	1.65	0.95
1:B:213:ILE:HG23	3:B:611:HOH:O	1.79	0.81
1:D:203:ASP:OD1	1:D:227:ARG:NH1	2.14	0.80
1:B:135:MET:HG3	1:B:139:LEU:HD22	1.66	0.78
1:C:135:MET:HG3	1:C:139:LEU:HD22	1.69	0.75
1:D:38:LYS:HD3	1:D:87:ASP:OD2	1.88	0.74
1:B:92[B]:ARG:NH2	3:B:601:HOH:O	2.20	0.73
1:B:156:LEU:HD12	1:B:161:LYS:HB2	1.74	0.70
1:C:32[B]:LEU:HD12	3:C:765:HOH:O	1.90	0.69
1:C:305:LYS:HD3	1:D:65:THR:HG22	1.73	0.68
1:B:208:HIS:HD2	3:B:635:HOH:O	1.75	0.68
1:C:161:LYS:HE3	3:D:613:HOH:O	1.92	0.68
1:C:357:LYS:HG3	1:C:360:PRO:HD2	1.77	0.67
1:C:367:GLN:HE21	1:C:368:LEU:H	1.42	0.66
1:C:317:LEU:HD12	1:C:342:ARG:HG3	1.77	0.66
1:C:385:ARG:HG3	1:C:386:ALA:N	2.11	0.65
1:C:397:ILE:HG12	1:D:155:GLU:HB2	1.77	0.65
1:A:156:LEU:HD12	1:A:161:LYS:HB2	1.78	0.64
1:D:219:THR:HG22	3:D:614:HOH:O	1.98	0.64
1:A:135:MET:HG3	1:A:139:LEU:HD22	1.78	0.63
1:C:203:ASP:OD1	1:C:227:ARG:NH2	2.32	0.63
1:C:301:PRO:HA	1:D:68:ASP:OD2	1.99	0.63
1:C:208:HIS:HE1	1:C:242:SER:OG	1.82	0.62
1:C:13:THR:HG21	1:C:32[B]:LEU:HD21	1.81	0.62
1:C:208:HIS:HD2	3:C:696:HOH:O	1.82	0.62
1:B:366[A]:ARG:HD3	3:B:787:HOH:O	2.00	0.61
1:A:132:GLU:HG2	1:A:170:ILE:HD13	1.85	0.59



A		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:D:208:HIS:HD2	3:D:603:HOH:O	1.85	0.59
1:A:271:GLY:HA2	1:A:397:ILE:O	2.02	0.59
1:A:44:THR:HG22	1:B:391:LEU:HD23	1.85	0.58
1:A:155:GLU:HB2	1:B:397:ILE:HG21	1.84	0.58
1:A:208:HIS:HD2	3:A:684:HOH:O	1.87	0.57
1:A:208:HIS:HE1	1:A:242:SER:OG	1.88	0.57
1:B:397:ILE:HD12	1:B:398:VAL:HG13	1.86	0.57
1:C:32[A]:LEU:HD11	1:C:126:LEU:HD11	1.87	0.56
1:A:158:VAL:HG12	1:B:262:LEU:HD13	1.87	0.55
1:D:156:LEU:HD12	1:D:161:LYS:HB2	1.88	0.55
1:C:197:GLU:HG2	1:C:202:LEU:HD22	1.89	0.55
1:C:385:ARG:HG3	1:C:386:ALA:H	1.71	0.55
1:D:208:HIS:HE1	1:D:242:SER:OG	1.89	0.55
1:C:305:LYS:CD	1:D:65:THR:HG22	2.37	0.54
1:C:393:MLE:HN2	1:C:393:MLE:HG	1.88	0.54
1:C:156:LEU:HD12	1:C:161:LYS:HB2	1.89	0.54
1:A:44:THR:HG22	1:B:391:LEU:CD2	2.39	0.53
1:C:357:LYS:HG3	1:C:360:PRO:CD	2.38	0.53
1:A:158:VAL:CG1	1:B:262:LEU:HD13	2.40	0.52
1:C:155:GLU:HB2	1:D:397:ILE:HG12	1.90	0.52
1:B:317:LEU:HD11	1:B:338:PHE:HE2	1.75	0.51
1:D:363:VAL:HG21	1:D:386:ALA:HB2	1.91	0.51
1:B:132:GLU:HG3	1:B:172:TRP:CZ2	2.45	0.51
1:D:290:MET:HE2	3:D:656:HOH:O	2.11	0.51
1:C:196:LEU:HB3	1:C:226:ILE:HD12	1.93	0.51
1:C:394:MVA:HN2	1:C:394:MVA:HG22	1.92	0.51
1:A:158:VAL:HG12	1:B:262:LEU:CD1	2.41	0.51
1:A:206:ILE:CG2	1:A:226:ILE:HD11	2.41	0.50
1:A:241:THR:HA	1:B:391:LEU:HD12	1.94	0.50
1:B:213:ILE:CG2	3:B:611:HOH:O	2.48	0.49
1:B:233:GLU:HA	1:B:236:LYS:HE2	1.93	0.49
1:B:132:GLU:HG2	1:B:170:ILE:HD13	1.93	0.49
3:A:605:HOH:O	1:B:161:LYS:HE3	2.12	0.49
1:B:279:GLN:NE2	3:B:602:HOH:O	2.38	0.48
1:C:13:THR:HG21	1:C:32[B]:LEU:CD2	2.44	0.48
1:A:206:ILE:HG23	1:A:226:ILE:HD11	1.94	0.48
1:C:397:ILE:HG12	1:D:155:GLU:CB	2.41	0.48
1:A:338:PHE:O	1:A:341:THR:HB	2.14	0.48
1:A:351:PHE:CE2	1:A:380:GLN:HG3	2.50	0.47
1:C:392:ALA:HA	1:C:393:MLE:HN1	1.59	0.47
1:D:150:THR:HG23	1:D:172:TRP:NE1	2.30	0.47



		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:D:216:GLN:NE2	1:D:216:GLN:H	2.13	0.47
1:B:208:HIS:HE1	1:B:242:SER:OG	1.97	0.47
3:C:643:HOH:O	1:D:161:LYS:HE3	2.15	0.46
1:C:155:GLU:CB	1:D:397:ILE:HG12	2.46	0.45
1:B:393:MLE:HA	1:B:394:MVA:HN1	1.81	0.45
1:C:357:LYS:HE3	1:C:359:GLY:H	1.82	0.45
1:C:32[A]:LEU:CD1	1:C:126:LEU:HD11	2.45	0.45
1:A:262:LEU:CD1	1:B:158:VAL:HG12	2.47	0.45
1:A:186:ASN:HD22	1:A:186:ASN:HA	1.63	0.44
1:A:393:MLE:HA	1:A:394:MVA:HN1	1.76	0.44
1:B:392:ALA:HA	1:B:393:MLE:HN1	1.67	0.43
1:C:190:HIS:HE1	3:C:647:HOH:O	2.01	0.43
1:D:107:ALA:N	1:D:108:PRO:HD2	2.34	0.43
1:C:12:LEU:O	1:C:123:ALA:HA	2.19	0.43
1:B:317:LEU:CD1	1:B:338:PHE:HE2	2.31	0.43
1:B:363:VAL:HG21	1:B:386:ALA:HB2	2.00	0.43
1:A:12:LEU:O	1:A:123:ALA:HA	2.19	0.42
1:B:216:GLN:H	1:B:216:GLN:HE21	1.67	0.42
1:C:155:GLU:HA	1:C:158:VAL:HG22	2.00	0.42
1:D:219:THR:HG23	3:D:737:HOH:O	2.18	0.42
1:D:270:LYS:HE2	3:D:685:HOH:O	2.19	0.42
1:C:391:LEU:CD2	1:C:393:MLE:HD22	2.49	0.42
1:C:315:GLN:HE21	1:C:315:GLN:HA	1.85	0.42
1:C:202:LEU:HB3	1:C:227:ARG:HB3	2.01	0.42
1:B:397:ILE:HD12	1:B:397:ILE:C	2.39	0.42
1:A:254:ILE:H	1:A:254:ILE:HG12	1.43	0.42
1:D:216:GLN:H	1:D:216:GLN:HE21	1.68	0.42
1:B:235:LEU:C	1:B:235:LEU:HD23	2.40	0.42
1:A:363:VAL:HG21	1:A:386:ALA:HB2	2.01	0.41
1:A:394:MVA:HN2	1:A:394:MVA:HG22	2.02	0.41
1:B:394:MVA:HN2	1:B:394:MVA:HG22	2.03	0.41
1:B:12:LEU:O	1:B:123:ALA:HA	2.20	0.41
1:D:219:THR:CG2	3:D:614:HOH:O	2.61	0.41
1:A:99:TYR:CZ	1:B:391:LEU:HD21	2.56	0.41
1:C:133:ASP:OD2	1:D:133:ASP:OD2	2.37	0.41
1:C:22:ILE:HD13	1:D:319:VAL:HG22	2.02	0.41
1:C:208:HIS:CE1	1:C:242:SER:OG	2.70	0.41
1:A:315:GLN:NE2	1:A:315:GLN:HA	2.36	0.41
1:C:262:LEU:CD1	1:D:158:VAL:HG12	2.51	0.41
1:D:186:ASN:HD22	1:D:186:ASN:HA	1.67	0.41
1:A:132:GLU:OE2	1:A:150:THR:OG1	2.21	0.40



Atom-1	tom-1 Atom-2		Clash overlap (Å)
1:A:107:ALA:N	1:A:108:PRO:HD2	2.36	0.40
1:C:44:THR:HG22	1:D:391:LEU:CD2	2.51	0.40
1:D:394:MVA:HG22	1:D:394:MVA:HN2	2.03	0.40
1:A:203:ASP:OD1	1:A:227:ARG:NH1	2.45	0.40
1:D:40:PHE:O	1:D:96:GLY:HA2	2.21	0.40
1:C:44:THR:CG2	1:D:391:LEU:CD2	2.99	0.40
1:C:186:ASN:HD22	1:C:186:ASN:HA	1.72	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	385/400~(96%)	377~(98%)	8 (2%)	0	100	100
1	В	386/400~(96%)	379~(98%)	7 (2%)	0	100	100
1	С	386/400~(96%)	378~(98%)	8 (2%)	0	100	100
1	D	385/400~(96%)	376~(98%)	9(2%)	0	100	100
All	All	1542/1600~(96%)	1510 (98%)	32(2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	321/327~(98%)	312~(97%)	9~(3%)	38 24
1	В	323/327~(99%)	310~(96%)	13 (4%)	27 12
1	С	321/327~(98%)	307~(96%)	14 (4%)	24 10
1	D	320/327~(98%)	307~(96%)	13 (4%)	26 11
All	All	1285/1308~(98%)	1236~(96%)	49 (4%)	27 13

All (49) residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	92	ARG
1	А	134	TYR
1	А	139	LEU
1	А	151	GLN
1	А	183	VAL
1	А	216	GLN
1	А	254	ILE
1	А	267	THR
1	А	281	VAL
1	В	134	TYR
1	В	139	LEU
1	В	151	GLN
1	В	183	VAL
1	В	216	GLN
1	В	218	VAL
1	В	227	ARG
1	В	236	LYS
1	В	260	GLN
1	В	267	THR
1	В	273	GLN
1	В	380	GLN
1	В	391	LEU
1	С	45	ASP
1	С	134	TYR
1	С	139	LEU
1	С	151	GLN
1	С	202	LEU
1	С	216	GLN
1	С	219	THR
1	С	236	LYS
1	С	250	THR
1	С	260	GLN
1	С	273	GLN



Mol	Chain	Res	Type
1	С	305	LYS
1	С	380	GLN
1	С	385	ARG
1	D	45	ASP
1	D	124	LYS
1	D	134	TYR
1	D	151	GLN
1	D	162	LYS
1	D	216	GLN
1	D	219	THR
1	D	233	GLU
1	D	235	LEU
1	D	260	GLN
1	D	290	MET
1	D	381	GLU
1	D	391	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (31) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	173	GLN
1	А	186	ASN
1	А	208	HIS
1	А	216	GLN
1	А	303	GLN
1	А	309	HIS
1	А	315	GLN
1	А	343	ASN
1	В	173	GLN
1	В	186	ASN
1	В	208	HIS
1	В	216	GLN
1	В	273	GLN
1	В	315	GLN
1	В	343	ASN
1	В	380	GLN
1	С	173	GLN
1	С	186	ASN
1	С	190	HIS
1	С	208	HIS
1	С	216	GLN
1	С	273	GLN



	5	1	1 5
Mol	Chain	Res	Type
1	С	279	GLN
1	С	315	GLN
1	С	367	GLN
1	С	380	GLN
1	D	173	GLN
1	D	186	ASN
1	D	208	HIS
1	D	216	GLN
1	D	343	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	True	Chain	Dec Link		B	ond leng	gths	E	ond ang	gles
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
1	MVA	С	394	1	6,7,8	0.71	0	$6,\!8,\!10$	1.09	0
1	MVA	А	394	1	6,7,8	0.72	0	$6,\!8,\!10$	1.49	1 (16%)
1	MVA	В	394	1	6,7,8	0.57	0	6,8,10	1.44	1 (16%)
1	MLE	В	393	1	7,8,9	0.54	0	7,9,11	1.41	2 (28%)
1	MLE	С	393	1	7,8,9	0.63	0	7,9,11	1.61	1 (14%)
1	MLE	А	393	1	7,8,9	0.53	0	7,9,11	1.85	1 (14%)
1	MVA	D	394	1	6,7,8	0.64	0	6,8,10	1.13	1 (16%)
1	MLE	D	393	1	7,8,9	0.52	0	7,9,11	1.52	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MVA	С	394	1	-	4/6/8/10	-
1	MVA	А	394	1	-	5/6/8/10	-
1	MVA	В	394	1	-	5/6/8/10	-
1	MLE	В	393	1	-	0/5/8/10	-
1	MLE	С	393	1	-	0/5/8/10	-
1	MLE	А	393	1	-	0/5/8/10	-
1	MVA	D	394	1	-	5/6/8/10	-
1	MLE	D	393	1	-	0/5/8/10	-

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	393	MLE	CB-CA-N	4.31	117.14	110.59
1	С	393	MLE	CB-CA-N	3.44	115.81	110.59
1	А	394	MVA	CB-CA-N	3.05	115.14	111.17
1	В	394	MVA	CB-CA-N	2.61	114.57	111.17
1	В	393	MLE	CB-CA-N	2.55	114.47	110.59
1	D	393	MLE	CB-CA-N	2.44	114.30	110.59
1	В	393	MLE	O-C-CA	-2.23	119.03	124.77
1	D	394	MVA	CB-CA-N	2.18	114.01	111.17
1	D	393	MLE	O-C-CA	-2.16	119.21	124.77

There are no chirality outliers.

All (19) torsion outliers are listed below:	All	(19)	torsion	outliers	are	listed	below:
---	-----	------	---------	----------	-----	--------	--------

Mol	Chain	\mathbf{Res}	Type	Atoms
1	А	394	MVA	N-CA-CB-CG1
1	А	394	MVA	N-CA-CB-CG2
1	А	394	MVA	C-CA-CB-CG1
1	А	394	MVA	C-CA-CB-CG2
1	В	394	MVA	N-CA-CB-CG1
1	В	394	MVA	N-CA-CB-CG2
1	В	394	MVA	C-CA-CB-CG1
1	В	394	MVA	C-CA-CB-CG2
1	С	394	MVA	N-CA-CB-CG1
1	С	394	MVA	N-CA-CB-CG2
1	С	394	MVA	C-CA-CB-CG1
1	С	394	MVA	C-CA-CB-CG2
1	D	394	MVA	N-CA-CB-CG1
1	D	394	MVA	N-CA-CB-CG2



Mol	Chain	Res	Type	Atoms						
1	D	394	MVA	C-CA-CB-CG1						
1	D	394	MVA	C-CA-CB-CG2						
1	А	394	MVA	O-C-CA-CB						
1	В	394	MVA	O-C-CA-CB						
1	D	394	MVA	O-C-CA-CB						

There are no ring outliers.

7 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	С	394	MVA	1	0
1	А	394	MVA	2	0
1	В	394	MVA	2	0
1	В	393	MLE	2	0
1	С	393	MLE	3	0
1	А	393	MLE	1	0
1	D	394	MVA	1	0

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mal Turna Chain		Dag	Link	Bo	Bond lengths			Bond angles		
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	SAH	В	500	-	23,28,28	0.79	0	22,40,40	0.96	2 (9%)	
2	SAH	А	500	-	23,28,28	0.78	0	22,40,40	0.94	1 (4%)	
2	SAH	С	500	-	23,28,28	0.66	0	22,40,40	0.92	2 (9%)	
2	SAH	D	500	-	23,28,28	0.76	1 (4%)	22,40,40	1.01	2 (9%)	



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	В	500	-	-	2/11/31/31	0/3/3/3
2	SAH	А	500	-	-	2/11/31/31	0/3/3/3
2	SAH	С	500	-	-	2/11/31/31	0/3/3/3
2	SAH	D	500	-	-	3/11/31/31	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	D	500	SAH	OXT-C	-2.16	1.23	1.30

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	D	500	SAH	C5-C6-N6	2.73	124.47	120.31
2	А	500	SAH	OXT-C-O	-2.71	117.93	124.08
2	В	500	SAH	C5-C6-N6	2.59	124.26	120.31
2	С	500	SAH	C5-C6-N6	2.58	124.25	120.31
2	D	500	SAH	OXT-C-O	-2.41	118.61	124.08
2	С	500	SAH	OXT-C-O	-2.25	118.97	124.08
2	В	500	SAH	OXT-C-O	-2.04	119.45	124.08

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	500	SAH	C3'-C4'-C5'-SD
2	D	500	SAH	O-C-CA-CB
2	В	500	SAH	OXT-C-CA-CB
2	D	500	SAH	OXT-C-CA-CB
2	В	500	SAH	O-C-CA-CB
2	А	500	SAH	O-C-CA-CB
2	А	500	SAH	OXT-C-CA-CB
2	С	500	SAH	OXT-C-CA-CB
2	С	500	SAH	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	386/400~(96%)	-0.06	10 (2%) 57 60	14, 30, 56, 95	3~(0%)
1	В	386/400~(96%)	-0.25	7 (1%) 67 70	15, 28, 46, 75	4 (1%)
1	С	385/400~(96%)	0.12	25 (6%) 26 27	15, 31, 66, 117	5 (1%)
1	D	387/400~(96%)	0.10	9 (2%) 61 63	18, 35, 54, 104	2(0%)
All	All	1544/1600~(96%)	-0.02	51 (3%) 49 53	14, 31, 58, 117	14 (0%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	268	VAL	6.0
1	А	391	LEU	4.8
1	А	281	VAL	4.3
1	С	281	VAL	4.2
1	А	268	VAL	4.2
1	С	391	LEU	4.2
1	В	268	VAL	3.8
1	D	390	ALA	3.8
1	D	281	VAL	3.7
1	В	391	LEU	3.5
1	С	269	THR	3.5
1	В	397	ILE	3.5
1	D	9	PRO	3.4
1	С	397	ILE	3.4
1	В	281	VAL	3.4
1	С	398	VAL	3.4
1	В	398	VAL	3.3
1	А	267	THR	3.3
1	D	386	ALA	3.2
1	С	383	ILE	3.0
1	А	397	ILE	2.9



Mol	Chain	\mathbf{Res}	Type	RSRZ
1	В	269	THR	2.7
1	С	368	LEU	2.7
1	С	305	LYS	2.7
1	D	90	ALA	2.6
1	А	341	THR	2.6
1	А	398	VAL	2.6
1	D	387	ASP	2.6
1	С	371	ALA	2.6
1	С	384	ALA	2.6
1	С	396	VAL	2.5
1	В	267	THR	2.5
1	С	374	SER	2.5
1	С	292	ALA	2.5
1	С	369	PRO	2.4
1	С	380	GLN	2.4
1	С	372	ILE	2.4
1	С	373	ALA	2.4
1	С	386	ALA	2.3
1	С	370	SER	2.3
1	А	269	THR	2.3
1	D	391	LEU	2.3
1	D	266	ALA	2.3
1	С	304	GLU	2.2
1	А	386	ALA	2.2
1	С	293	VAL	2.2
1	С	266	ALA	2.2
1	А	282	SER	2.2
1	С	290	MET	2.1
1	С	267	THR	2.1
1	D	269	THR	2.1

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{\AA}^2)$	Q<0.9
1	MVA	С	394	8/9	0.91	0.12	34,37,39,39	0
1	MVA	В	394	8/9	0.92	0.10	21,25,26,29	0
1	MLE	С	393	9/10	0.92	0.11	$35,\!40,\!42,\!44$	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
1	MLE	D	393	9/10	0.93	0.09	$26,\!28,\!31,\!32$	0
1	MVA	А	394	8/9	0.93	0.08	$25,\!26,\!28,\!29$	0
1	MLE	В	393	9/10	0.94	0.09	$26,\!29,\!36,\!38$	0
1	MLE	А	393	9/10	0.94	0.10	27,30,38,40	0
1	MVA	D	394	8/9	0.97	0.04	23,25,27,27	0

Continued from previous page...

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q < 0.9
2	SAH	D	500	26/26	0.95	0.06	$23,\!26,\!32,\!35$	0
2	SAH	В	500	26/26	0.97	0.06	19,20,22,25	0
2	SAH	С	500	26/26	0.98	0.05	20,23,24,25	0
2	SAH	А	500	26/26	0.98	0.04	18,20,21,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















6.5 Other polymers (i)

There are no such residues in this entry.

