



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2024 – 05:22 PM EST

PDB ID : 8TXF
Title : AvrB bound with RIN4 C-NOI motif
Authors : Peng, W.; Orth, K.
Deposited on : 2023-08-23
Resolution : 1.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	 7% 86% 6% 9%
2	B	30	 10% 77% 7% 17%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5469 atoms, of which 2531 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Avirulence protein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	295	4611	1460	2266	424	454	7	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	expression tag	UNP P13835
A	0	HIS	-	expression tag	UNP P13835

- Molecule 2 is a protein called RIN4.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	B	25	374	126	175	33	40	0	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
4	A	1	20	4	12	1	3	20	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	313	Total	O	0	0
			313	313		
5	B	21	Total	O	0	0
			21	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Avirulence protein B



- Molecule 2: RIN4



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	57.80Å 119.90Å 46.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.71 – 1.29 36.71 – 1.29	Depositor EDS
% Data completeness (in resolution range)	99.0 (36.71-1.29) 99.0 (36.71-1.29)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.11 (at 1.29Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.166 , 0.180 0.165 , 0.179	Depositor DCC
R_{free} test set	1991 reflections (2.45%)	wwPDB-VP
Wilson B-factor (Å ²)	13.6	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5469	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2396	0.67	0/3250
2	B	0.41	0/206	0.54	0/280
All	All	0.48	0/2602	0.66	0/3530

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2345	2266	2265	17	0
2	B	199	175	175	2	0
3	A	52	78	78	4	0
4	A	8	12	12	0	0
5	A	313	0	0	9	1
5	B	21	0	0	1	0
All	All	2938	2531	2530	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:GLN:OE1	5:A:1101:HOH:O	2.00	0.79
1:A:197:GLU:N	1:A:197:GLU:OE1	2.17	0.76
1:A:205:GLN:CG	5:A:1101:HOH:O	2.47	0.62
1:A:142:GLU:HG2	5:A:1317:HOH:O	2.01	0.61
1:A:247:LEU:HD23	1:A:285:ILE:HD11	1.84	0.60
1:A:239:SER:OG	1:A:241:ASP:OD2	2.19	0.59
1:A:188:ARG:HH11	1:A:188:ARG:HG2	1.69	0.56
1:A:205:GLN:CD	5:A:1101:HOH:O	2.40	0.56
1:A:193:HIS:O	1:A:197:GLU:OE1	2.28	0.52
2:B:1:ALA:N	5:B:201:HOH:O	2.42	0.51
1:A:247:LEU:HD23	1:A:285:ILE:CD1	2.41	0.51
3:A:1009:EDO:O2	5:A:1102:HOH:O	2.19	0.50
1:A:232:ASP:OD1	5:A:1104:HOH:O	2.20	0.48
1:A:195:ARG:HD2	2:B:159:PRO:HG2	1.96	0.47
1:A:294:ASN:OD1	3:A:1011:EDO:C1	2.66	0.44
3:A:1004:EDO:H22	5:A:1295:HOH:O	2.17	0.43
1:A:188:ARG:HG2	1:A:188:ARG:NH1	2.34	0.42
1:A:310:GLU:HG2	5:A:1118:HOH:O	2.21	0.41
1:A:154:PHE:CD2	1:A:187:LEU:HD23	2.56	0.41
1:A:243:SER:OG	1:A:246:GLU:HG3	2.21	0.41
3:A:1004:EDO:C2	5:A:1295:HOH:O	2.69	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1378:HOH:O	5:A:1403:HOH:O[4_554]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/323 (91%)	290 (99%)	2 (1%)	1 (0%)	41	17
2	B	23/30 (77%)	22 (96%)	1 (4%)	0	100	100
All	All	316/353 (90%)	312 (99%)	3 (1%)	1 (0%)	41	17

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	53	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/274 (91%)	249 (100%)	0	100	100
2	B	21/26 (81%)	21 (100%)	0	100	100
All	All	270/300 (90%)	270 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	1002	-	3,3,3	0.58	0	2,2,2	0.31	0
3	EDO	A	1006	-	3,3,3	0.39	0	2,2,2	0.17	0
3	EDO	A	1011	-	3,3,3	0.48	0	2,2,2	0.31	0
3	EDO	A	1012	-	3,3,3	0.47	0	2,2,2	0.33	0
3	EDO	A	1013	-	3,3,3	0.49	0	2,2,2	0.27	0
3	EDO	A	1010	-	3,3,3	0.45	0	2,2,2	0.38	0
3	EDO	A	1005	-	3,3,3	0.53	0	2,2,2	0.12	0
3	EDO	A	1003	-	3,3,3	0.57	0	2,2,2	0.05	0
3	EDO	A	1007	-	3,3,3	0.51	0	2,2,2	0.46	0
3	EDO	A	1008	-	3,3,3	0.63	0	2,2,2	0.24	0
3	EDO	A	1004	-	3,3,3	0.45	0	2,2,2	0.16	0
3	EDO	A	1001	-	3,3,3	0.37	0	2,2,2	0.69	0
3	EDO	A	1009	-	3,3,3	0.49	0	2,2,2	0.31	0
4	TRS	A	1014	-	7,7,7	1.37	1 (14%)	9,9,9	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1002	-	-	0/1/1/1	-
3	EDO	A	1006	-	-	0/1/1/1	-
3	EDO	A	1011	-	-	0/1/1/1	-
3	EDO	A	1012	-	-	1/1/1/1	-
3	EDO	A	1013	-	-	0/1/1/1	-
3	EDO	A	1010	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1005	-	-	0/1/1/1	-
3	EDO	A	1003	-	-	0/1/1/1	-
3	EDO	A	1007	-	-	1/1/1/1	-
3	EDO	A	1008	-	-	1/1/1/1	-
3	EDO	A	1004	-	-	1/1/1/1	-
3	EDO	A	1001	-	-	1/1/1/1	-
3	EDO	A	1009	-	-	1/1/1/1	-
4	TRS	A	1014	-	-	0/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1014	TRS	C-N	3.50	1.61	1.49

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1012	EDO	O1-C1-C2-O2
3	A	1004	EDO	O1-C1-C2-O2
3	A	1009	EDO	O1-C1-C2-O2
3	A	1010	EDO	O1-C1-C2-O2
3	A	1007	EDO	O1-C1-C2-O2
3	A	1001	EDO	O1-C1-C2-O2
3	A	1008	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1011	EDO	1	0
3	A	1004	EDO	2	0
3	A	1009	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/323 (91%)	0.39	22 (7%) 14 11	8, 16, 49, 73	0
2	B	25/30 (83%)	0.88	3 (12%) 4 3	12, 28, 43, 61	0
All	All	320/353 (90%)	0.42	25 (7%) 13 10	8, 17, 49, 73	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	320	CYS	8.2
1	A	54	SER	7.4
1	A	26	PHE	6.5
1	A	197	GLU	5.9
1	A	196	THR	5.4
1	A	193	HIS	4.8
1	A	241	ASP	4.7
1	A	40	TYR	4.1
2	B	165	TYR	3.9
1	A	194	LEU	3.7
1	A	198	LEU	3.6
1	A	148	ILE	3.4
1	A	240	SER	3.3
1	A	55	SER	3.3
2	B	156	GLU	3.1
1	A	195	ARG	3.0
1	A	186	TYR	2.8
1	A	56	LYS	2.7
1	A	58	ASN	2.6
1	A	192	ALA	2.3
2	B	155	ASP	2.2
1	A	27	ARG	2.1
1	A	187	LEU	2.1
1	A	190	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	57	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

LIGAND-RSR INFOmissingINFO

6.5 Other polymers [i](#)

There are no such residues in this entry.