



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:48 am BST

PDB ID : 5U02
Title : Crystal structure of S. aureus TarS 217-571
Authors : Worrall, L.J.; Sobhanifar, S.; King, D.T.; Strynadka, N.C.
Deposited on : 2016-11-22
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyl transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	355	2916	1875	489	540	12	0	0	0

- Molecule 2 is IMIDAZOLE (three-letter code: IMD) (formula: C₃H₅N₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
2	A	1	5	3	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	179	179	179	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycosyl transferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.69 Å 105.69 Å 80.50 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	52.83 – 2.30 91.53 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.1 (52.83-2.30) 98.3 (91.53-2.30)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.29 Å)	Xtrriage
Refinement program	PHENIX (1.10_2155)	Depositor
R, R_{free}	0.224 , 0.268 0.224 , 0.271	Depositor DCC
R_{free} test set	1229 reflections (5.34%)	wwPDB-VP
Wilson B-factor (Å ²)	33.5	Xtrriage
Anisotropy	0.019	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3100	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/2976	0.50	0/4013

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2916	0	2911	58	0
2	A	5	0	5	3	0
3	A	179	0	0	7	3
All	All	3100	0	2916	59	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:476:PHE:O	3:A:701:HOH:O	1.86	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:471:ARG:HH12	2:A:601:IMD:H2	1.36	0.91
1:A:380:GLU:OE1	3:A:702:HOH:O	1.97	0.81
1:A:321:GLN:HE21	1:A:323:GLN:HB3	1.46	0.79
1:A:448:ASN:HD22	1:A:482:ASN:HD21	1.33	0.75
1:A:321:GLN:NE2	1:A:323:GLN:HB3	2.03	0.73
1:A:266:VAL:O	1:A:271:GLN:NE2	2.21	0.72
1:A:321:GLN:HE21	1:A:323:GLN:H	1.37	0.71
1:A:513:ARG:NH2	3:A:708:HOH:O	2.22	0.71
1:A:317:GLU:HB3	1:A:320:ARG:HH21	1.56	0.71
1:A:453:TYR:O	3:A:703:HOH:O	2.10	0.69
1:A:317:GLU:O	1:A:320:ARG:HG2	1.95	0.67
1:A:233:LEU:O	1:A:242:LYS:NZ	2.30	0.64
1:A:316:GLU:OE2	3:A:705:HOH:O	2.14	0.64
1:A:321:GLN:HE21	1:A:323:GLN:N	1.95	0.64
1:A:342:GLU:HG3	1:A:343:PRO:HD2	1.81	0.63
1:A:321:GLN:NE2	1:A:323:GLN:H	1.99	0.60
1:A:293:VAL:O	3:A:706:HOH:O	2.16	0.60
1:A:525:VAL:HB	1:A:561:LYS:HB3	1.85	0.59
1:A:448:ASN:HD22	1:A:482:ASN:ND2	2.01	0.58
1:A:318:SER:HA	1:A:321:GLN:HE22	1.68	0.58
1:A:264:LEU:HD11	1:A:316:GLU:HB3	1.85	0.58
1:A:318:SER:HB3	1:A:324:TYR:CD1	2.39	0.57
1:A:321:GLN:HE21	1:A:323:GLN:CB	2.17	0.57
1:A:237:LEU:HG	1:A:238:GLU:H	1.69	0.57
1:A:235:ALA:O	1:A:242:LYS:NZ	2.22	0.56
1:A:227:LEU:O	1:A:231:GLU:HG3	2.06	0.56
1:A:243:ASP:O	1:A:247:ALA:N	2.34	0.55
1:A:313:ARG:HG2	1:A:317:GLU:HG3	1.88	0.55
1:A:336:ILE:HD12	1:A:348:ILE:HD12	1.89	0.54
1:A:303:TYR:OH	1:A:342:GLU:HG2	2.08	0.53
1:A:471:ARG:NH1	2:A:601:IMD:H2	2.16	0.52
2:A:601:IMD:N1	3:A:704:HOH:O	2.11	0.52
1:A:314:THR:O	1:A:318:SER:N	2.40	0.52
1:A:389:LEU:HB2	1:A:398:MET:HB2	1.93	0.51
1:A:351:ALA:H	1:A:377:MET:CE	2.23	0.51
1:A:290:ASP:N	1:A:290:ASP:OD1	2.44	0.50
1:A:387:LEU:HD23	1:A:414:ILE:HD11	1.93	0.49
1:A:226:ARG:O	1:A:230:VAL:HG23	2.12	0.49
1:A:271:GLN:HB3	1:A:309:ILE:CD1	2.44	0.46
1:A:554:ARG:HG3	1:A:555:ASP:OD2	2.16	0.45
1:A:279:GLY:HA3	1:A:307:LYS:HE2	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:SER:HB3	1:A:324:TYR:CE1	2.52	0.45
1:A:378:ILE:HD11	1:A:382:HIS:HB2	1.99	0.44
1:A:237:LEU:CG	1:A:238:GLU:H	2.31	0.44
1:A:334:LEU:HD23	1:A:336:ILE:HD11	2.00	0.44
1:A:321:GLN:HE21	1:A:323:GLN:CA	2.30	0.43
1:A:561:LYS:HB2	1:A:561:LYS:HE3	1.85	0.43
1:A:224:VAL:O	1:A:228:ILE:HG13	2.19	0.43
1:A:325:TYR:O	1:A:326:ARG:HD2	2.18	0.43
1:A:222:TYR:HB2	1:A:277:ALA:HB1	2.01	0.42
1:A:258:ARG:HD3	1:A:274:TRP:HH2	1.83	0.42
1:A:313:ARG:O	1:A:317:GLU:HG3	2.20	0.42
1:A:317:GLU:HA	1:A:320:ARG:HE	1.83	0.42
1:A:243:ASP:HA	1:A:246:LEU:HB3	2.01	0.42
1:A:317:GLU:HB3	1:A:320:ARG:NH2	2.28	0.42
1:A:458:GLU:HB3	1:A:513:ARG:HH11	1.85	0.41
1:A:479:ASP:OD1	1:A:480:PHE:N	2.53	0.41
1:A:324:TYR:CE1	1:A:336:ILE:HD13	2.57	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:772:HOH:O	3:A:856:HOH:O[4_556]	1.76	0.44
3:A:709:HOH:O	3:A:824:HOH:O[3_565]	2.01	0.19
3:A:707:HOH:O	3:A:707:HOH:O[4_556]	2.14	0.06

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	353/356 (99%)	343 (97%)	10 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	319/322 (99%)	311 (98%)	8 (2%)	47 65

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	ARG
1	A	276	ASN
1	A	290	ASP
1	A	296	LYS
1	A	326	ARG
1	A	328	ASP
1	A	376	PHE
1	A	397	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	311	ASN
1	A	321	GLN
1	A	339	ASN
1	A	449	GLN
1	A	482	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IMD	A	601	-	3,5,5	0.40	0	4,5,5	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMD	A	601	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	IMD	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/356 (99%)	0.72	42 (11%) 4 6	21, 42, 105, 134	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	250	LEU	7.3
1	A	289	VAL	6.7
1	A	237	LEU	6.5
1	A	273	GLN	6.1
1	A	247	ALA	5.8
1	A	227	LEU	5.4
1	A	233	LEU	4.9
1	A	274	TRP	4.7
1	A	243	ASP	4.4
1	A	245	ILE	3.7
1	A	229	ALA	3.5
1	A	224	VAL	3.5
1	A	246	LEU	3.4
1	A	239	GLU	3.4
1	A	293	VAL	3.3
1	A	225	MET	3.3
1	A	276	ASN	3.3
1	A	277	ALA	3.2
1	A	218	PRO	3.2
1	A	268	LEU	3.2
1	A	252	ARG	3.1
1	A	241	HIS	3.0
1	A	288	ARG	2.9
1	A	221	PHE	2.7
1	A	258	ARG	2.7
1	A	232	ILE	2.7
1	A	278	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	292	LEU	2.6
1	A	230	VAL	2.5
1	A	226	ARG	2.5
1	A	254	PHE	2.4
1	A	341	GLY	2.3
1	A	228	ILE	2.3
1	A	265	LYS	2.3
1	A	309	ILE	2.3
1	A	264	LEU	2.2
1	A	280	ASP	2.2
1	A	378	ILE	2.2
1	A	329	ILE	2.1
1	A	304	ALA	2.1
1	A	282	ILE	2.1
1	A	321	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	IMD	A	601	5/5	0.91	0.23	49,50,54,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.