



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2023 – 09:14 PM EDT

PDB ID : 7U4Q
Title : Human Synaptotagmin-1 C2A Y181F Ca²⁺ bound
Authors : Dominguez, M.J.; Karmakar, S.; Fuson, K.L.; Sutton, R.B.
Deposited on : 2022-02-28
Resolution : 1.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	 12% 85% 9% • 5%
1	B	133	 13% 86% 8% • 5%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4520 atoms, of which 2132 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Synaptotagmin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	126	2191	715	1091	180	200	5	0	15	0
1	B	126	2093	682	1041	172	195	3	0	6	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	135	GLY	-	expression tag	UNP P21579
A	136	ASP	-	expression tag	UNP P21579
A	137	ILE	-	expression tag	UNP P21579
A	138	THR	-	expression tag	UNP P21579
A	139	HIS	-	expression tag	UNP P21579
A	140	MET	-	expression tag	UNP P21579
A	181	PHE	TYR	engineered mutation	UNP P21579
B	135	GLY	-	expression tag	UNP P21579
B	136	ASP	-	expression tag	UNP P21579
B	137	ILE	-	expression tag	UNP P21579
B	138	THR	-	expression tag	UNP P21579
B	139	HIS	-	expression tag	UNP P21579
B	140	MET	-	expression tag	UNP P21579
B	181	PHE	TYR	engineered mutation	UNP P21579

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	118	Total 118	O 118	0	0
3	B	115	Total 115	O 115	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Synaptotagmin-1

Chain A: 



- Molecule 1: Synaptotagmin-1

Chain B: 



4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	99.81Å 99.81Å 77.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.06 – 1.56 26.06 – 1.56	Depositor EDS
% Data completeness (in resolution range)	99.9 (26.06-1.56) 100.0 (26.06-1.56)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.56Å)	Xtrriage
Refinement program	PHENIX 1.20_4459	Depositor
R, R_{free}	0.164 , 0.189 0.161 , 0.187	Depositor DCC
R_{free} test set	1992 reflections (3.70%)	wwPDB-VP
Wilson B-factor (Å ²)	27.2	Xtrriage
Anisotropy	0.663	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.020 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4520	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/1181	0.72	0/1589
1	B	0.55	0/1109	0.74	0/1498
All	All	0.54	0/2290	0.73	0/3087

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1100	1091	1030	14	0
1	B	1052	1041	1013	11	0
2	A	3	0	0	0	0
3	A	118	0	0	1	0
3	B	115	0	0	0	0
All	All	2388	2132	2043	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:LYS:HG3	1:A:192:LYS:H	1.01	1.15
1:A:191:LYS:HG3	1:A:192:LYS:N	1.83	0.91
1:A:191:LYS:CG	1:A:192:LYS:H	1.84	0.79
1:B:174:MET:HA	1:B:174:MET:HE2	1.70	0.73
1:B:150:LEU:HD21	1:B:225:LEU:HD22	1.89	0.55
1:A:191:LYS:CG	1:A:192:LYS:N	2.52	0.53
1:B:174:MET:HA	1:B:174:MET:CE	2.39	0.53
1:A:189:ASP:HB3	1:A:191:LYS:HG2	1.92	0.52
1:B:180:PRO:HD2	1:B:199:HIS:O	2.11	0.51
1:A:180:PRO:HD2	1:A:199:HIS:O	2.11	0.50
1:A:189:ASP:CB	1:A:191:LYS:HE3	2.44	0.48
1:B:141:GLU:O	1:B:141:GLU:HG2	2.14	0.47
1:A:146:LEU:HD23	1:A:244[B]:PHE:CE1	2.51	0.46
1:B:150:LEU:HD21	1:B:225:LEU:CD2	2.47	0.45
1:B:158:GLN:HG2	1:B:214:LYS:HD2	1.99	0.44
1:A:174[B]:MET:O	1:A:174[B]:MET:HG3	2.16	0.44
1:A:192:LYS:HE3	1:A:194:PHE:CZ	2.53	0.44
1:B:173:ASP:OD1	1:B:173:ASP:N	2.50	0.43
1:A:173[A]:ASP:OD2	3:A:401:HOH:O	2.22	0.42
1:A:189:ASP:HB2	1:A:191:LYS:HE3	2.00	0.42
1:B:146:LEU:HD23	1:B:244:PHE:CE1	2.55	0.42
1:B:174:MET:CE	1:B:174:MET:CA	2.98	0.42
1:A:244[A]:PHE:CE2	1:A:261:ARG:HG3	2.55	0.41
1:A:231:ASP:HB3	1:A:239:ASP:HB2	2.03	0.40
1:B:146:LEU:HD23	1:B:244:PHE:HE1	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	139/133 (104%)	135 (97%)	3 (2%)	1 (1%)	22 5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	130/133 (98%)	125 (96%)	4 (3%)	1 (1%)	19	4
All	All	269/266 (101%)	260 (97%)	7 (3%)	2 (1%)	19	5

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	191	LYS
1	B	172	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/119 (107%)	127 (100%)	0	100	100
1	B	119/119 (100%)	118 (99%)	1 (1%)	81	66
All	All	246/238 (103%)	245 (100%)	1 (0%)	91	82

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	173	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	126/133 (94%)	0.68	16 (12%) 3 3	23, 36, 62, 88	0
1	B	126/133 (94%)	0.95	17 (13%) 3 2	22, 34, 66, 116	0
All	All	252/266 (94%)	0.82	33 (13%) 3 2	22, 35, 69, 116	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	174	MET	12.8
1	B	175	GLY	9.7
1	B	172	LEU	8.9
1	A	235	PHE	7.6
1	B	173	ASP	7.3
1	B	171	ALA	6.9
1	B	266	ALA	6.5
1	B	191	LYS	6.3
1	A	191	LYS	5.3
1	A	255[A]	HIS	4.5
1	B	256[A]	VAL	4.3
1	A	234	ARG	4.2
1	B	257	THR	4.0
1	B	190	LYS	3.8
1	A	237	LYS	3.8
1	A	189	ASP	3.7
1	B	244	PHE	3.7
1	B	142	LYS	3.5
1	A	229	VAL	3.3
1	A	190	LYS	3.1
1	B	176	GLY	3.1
1	B	255[A]	HIS	3.0
1	B	141	GLU	3.0
1	A	236	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	163	ILE	2.7
1	A	266	ALA	2.7
1	A	182	VAL	2.6
1	B	182	VAL	2.6
1	A	164	ILE	2.4
1	A	253	PHE	2.3
1	A	188	PRO	2.2
1	B	207	PHE	2.1
1	A	244[A]	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	A	303	1/1	0.97	0.05	59,59,59,59	1
2	CA	A	302	1/1	0.99	0.10	36,36,36,36	1
2	CA	A	301	1/1	1.00	0.06	35,35,35,35	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CA A 303:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.