



wwPDB X-ray Structure Validation Summary Report

Aug 7, 2020 – 01:18 AM BST

PDB ID : 5U9C
Title : 1.9 Angstrom Resolution Crystal Structure of dTDP-4-dehydrorhamnose Reductase from *Yersinia enterocolitica*
Authors : Minasov, G.; Shuvalova, L.; Flores, K.; Dubrovska, I.; Olphie, A.; Grimshaw, S.; Kwon, K.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2016-12-15
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	
1	B	292	
1	C	292	
1	D	292	
1	E	292	
1	F	292	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMD	A	303	-	-	-	X
5	CIT	D	302	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 15720 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called dTDP-4-dehydrorhamnose Reductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	288	2324	1481	395	434	5	9	0	15	0
1	B	287	2365	1507	403	440	5	10	0	21	0
1	C	288	2313	1474	396	431	5	7	0	13	0
1	D	291	2357	1501	401	441	5	9	0	17	0
1	E	291	2405	1526	411	453	5	10	0	23	0
1	F	287	2342	1490	400	439	5	8	0	18	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q56905
A	-1	ASN	-	expression tag	UNP Q56905
A	0	ALA	-	expression tag	UNP Q56905
B	-2	SER	-	expression tag	UNP Q56905
B	-1	ASN	-	expression tag	UNP Q56905
B	0	ALA	-	expression tag	UNP Q56905
C	-2	SER	-	expression tag	UNP Q56905
C	-1	ASN	-	expression tag	UNP Q56905
C	0	ALA	-	expression tag	UNP Q56905
D	-2	SER	-	expression tag	UNP Q56905
D	-1	ASN	-	expression tag	UNP Q56905
D	0	ALA	-	expression tag	UNP Q56905
E	-2	SER	-	expression tag	UNP Q56905
E	-1	ASN	-	expression tag	UNP Q56905
E	0	ALA	-	expression tag	UNP Q56905
F	-2	SER	-	expression tag	UNP Q56905
F	-1	ASN	-	expression tag	UNP Q56905

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	0	ALA	-	expression tag	UNP Q56905

- Molecule 2 is beta-D-fructopyranose (three-letter code: BDF) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	12	6	6	0	0
2	C	1	12	6	6	0	0
2	C	1	12	6	6	0	0

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N 5 3 2	0	0
3	A	1	Total C N 5 3 2	0	0

- Molecule 4 is D-MALATE (three-letter code: MLT) (formula: $C_4H_6O_5$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 9 4 5	0	0

- Molecule 5 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total C O 13 6 7	0	0
5	D	1	Total C O 13 6 7	0	0
5	E	1	Total C O 13 6 7	0	0
5	F	1	Total C O 13 6 7	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	D	1	Total Cl 2 2	0	1

- Molecule 7 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	F	1	8	4	1	3	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	265	Total	O	0	10
			271	271		
8	B	188	Total	O	0	9
			194	194		
8	C	252	Total	O	0	12
			263	263		
8	D	292	Total	O	0	17
			306	306		
8	E	223	Total	O	0	10
			231	231		
8	F	226	Total	O	0	7
			232	232		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: dTDP-4-dehydrorhamnose Reductase




- Molecule 1: dTDP-4-dehydrorhamnose Reductase



- Molecule 1: dTDP-4-dehydrorhamnose Reductase



- Molecule 1: dTDP-4-dehydrorhamnose Reductase




- Molecule 1: dTDP-4-dehydrorhamnose Reductase





- Molecule 1: dTDP-4-dehydrorhamnose Reductase

Chain F: 4% 95%



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.44Å 184.87Å 187.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.94 – 1.90 29.94 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.94-1.90) 99.6 (29.94-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.160 , 0.191 0.167 , 0.197	Depositor DCC
R_{free} test set	8981 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtrriage
Anisotropy	0.109	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.001 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15720	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BDF, IMD, CL, MLT, CIT, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.49	0/2367	0.70	1/3195 (0.0%)
1	B	0.47	0/2408	0.70	0/3248
1	C	0.48	0/2356	0.69	0/3181
1	D	0.51	0/2399	0.71	0/3237
1	E	0.49	0/2447	0.72	3/3300 (0.1%)
1	F	0.46	0/2385	0.70	0/3220
All	All	0.49	0/14362	0.70	4/19381 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	E	129	ARG	NE-CZ-NH2	-5.65	117.48	120.30
1	A	129	ARG	NE-CZ-NH1	5.59	123.10	120.30
1	E	264[A]	MSE	CG-SE-CE	-5.26	87.32	98.90
1	E	264[B]	MSE	CG-SE-CE	-5.26	87.32	98.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2324	0	2334	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2365	0	2379	16	0
1	C	2313	0	2326	6	0
1	D	2357	0	2369	12	0
1	E	2405	0	2403	8	0
1	F	2342	0	2344	7	0
2	A	12	0	12	0	0
2	C	24	0	24	0	0
3	A	10	0	10	0	0
4	B	9	0	4	0	0
5	D	26	0	10	1	0
5	E	13	0	5	1	0
5	F	13	0	5	0	0
6	D	2	0	0	0	0
7	F	8	0	12	0	0
8	A	271	0	0	0	0
8	B	194	0	0	1	0
8	C	263	0	0	0	0
8	D	306	0	0	2	0
8	E	231	0	0	0	0
8	F	232	0	0	0	0
All	All	15720	0	14237	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 55 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:212[A]:LEU:HD21	1:E:264[A]:MSE:SE	2.23	0.89
1:B:13[B]:MSE:HA	1:B:13[B]:MSE:HE3	1.58	0.83
1:B:71[B]:ARG:NH2	8:B:401:HOH:O	2.12	0.81
1:D:234[A]:ASN:OD1	1:D:236[A]:LYS:HE2	1.85	0.76
1:B:13[B]:MSE:HA	1:B:13[B]:MSE:CE	2.18	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/292 (103%)	294 (98%)	7 (2%)	0	100	100
1	B	306/292 (105%)	296 (97%)	9 (3%)	1 (0%)	41	31
1	C	299/292 (102%)	288 (96%)	11 (4%)	0	100	100
1	D	306/292 (105%)	299 (98%)	7 (2%)	0	100	100
1	E	312/292 (107%)	304 (97%)	7 (2%)	1 (0%)	41	31
1	F	303/292 (104%)	293 (97%)	10 (3%)	0	100	100
All	All	1827/1752 (104%)	1774 (97%)	51 (3%)	2 (0%)	51	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	143	THR
1	E	268	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/236 (108%)	253 (100%)	1 (0%)	91	91
1	B	259/236 (110%)	256 (99%)	3 (1%)	71	70
1	C	251/236 (106%)	251 (100%)	0	100	100
1	D	258/236 (109%)	258 (100%)	0	100	100
1	E	264/236 (112%)	259 (98%)	5 (2%)	57	53

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	256/236 (108%)	256 (100%)	0	100	100
All	All	1542/1416 (109%)	1533 (99%)	9 (1%)	88	87

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	124[A]	MSE
1	E	264[B]	MSE
1	E	220	ILE
1	B	33	GLU
1	E	124[B]	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	35	ASN
1	D	81	HIS
1	E	81	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	CIT	D	302	-	3,12,12	0.36	0	3,17,17	4.84	3 (100%)
2	BDF	C	302	-	12,12,12	1.00	1 (8%)	18,18,18	1.58	5 (27%)
2	BDF	A	301	-	12,12,12	0.99	1 (8%)	18,18,18	0.99	1 (5%)
2	BDF	C	301	-	12,12,12	0.92	1 (8%)	18,18,18	0.66	0
3	IMD	A	303	-	3,5,5	0.35	0	4,5,5	0.59	0
4	MLT	B	301	-	2,8,8	0.41	0	3,10,10	0.32	0
7	TRS	F	302	-	7,7,7	0.33	0	9,9,9	0.38	0
3	IMD	A	302	-	3,5,5	0.33	0	4,5,5	0.58	0
5	CIT	E	301	-	3,12,12	0.69	0	3,17,17	1.27	1 (33%)
5	CIT	F	301	-	3,12,12	0.43	0	3,17,17	0.79	0
5	CIT	D	301	-	3,12,12	0.66	0	3,17,17	1.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CIT	D	302	-	-	4/6/16/16	-
2	BDF	C	302	-	-	0/3/23/23	0/1/1/1
2	BDF	A	301	-	-	3/3/23/23	0/1/1/1
2	BDF	C	301	-	-	0/3/23/23	0/1/1/1
3	IMD	A	303	-	-	-	0/1/1/1
4	MLT	B	301	-	-	2/2/8/8	-
7	TRS	F	302	-	-	2/9/9/9	-
3	IMD	A	302	-	-	-	0/1/1/1
5	CIT	E	301	-	-	0/6/16/16	-
5	CIT	F	301	-	-	4/6/16/16	-
5	CIT	D	301	-	-	0/6/16/16	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	BDF	O6-C2	2.45	1.44	1.42
2	C	302	BDF	O6-C2	2.33	1.44	1.42
2	C	301	BDF	O6-C2	2.06	1.44	1.42

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	302	CIT	C4-C3-C2	6.70	127.24	109.33
5	D	302	CIT	C3-C4-C5	-4.34	108.03	114.98
2	C	302	BDF	O6-C2-C3	3.34	113.10	109.76
2	C	302	BDF	C5-C4-C3	2.74	113.55	110.48
5	D	302	CIT	C3-C2-C1	-2.56	110.89	114.98

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	302	CIT	C1-C2-C3-O7
5	D	302	CIT	C1-C2-C3-C6
2	A	301	BDF	O1-C1-C2-O2
2	A	301	BDF	O1-C1-C2-O6
7	F	302	TRS	N-C-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	302	CIT	1	0
5	E	301	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/292 (96%)	0.02	16 (5%) 23 26	22, 35, 73, 98	0
1	B	282/292 (96%)	0.26	19 (6%) 17 20	25, 42, 79, 105	0
1	C	283/292 (96%)	0.16	18 (6%) 19 22	24, 37, 71, 109	0
1	D	285/292 (97%)	-0.06	8 (2%) 53 56	23, 32, 55, 78	0
1	E	285/292 (97%)	0.04	13 (4%) 32 35	24, 36, 66, 108	0
1	F	282/292 (96%)	0.01	13 (4%) 32 35	25, 37, 61, 97	0
All	All	1699/1752 (96%)	0.07	87 (5%) 28 31	22, 36, 72, 109	0

The worst 5 of 87 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	289	VAL	11.2
1	C	288	LEU	8.8
1	E	-2	SER	8.6
1	A	288	LEU	7.8
1	B	288	LEU	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CIT	D	301	13/13	0.63	0.35	68,88,101,105	0
5	CIT	F	301	13/13	0.65	0.31	73,98,105,107	0
4	MLT	B	301	9/9	0.65	0.20	66,78,89,90	0
5	CIT	D	302	13/13	0.66	0.53	80,90,95,96	0
3	IMD	A	302	5/5	0.67	0.18	95,95,95,95	0
7	TRS	F	302	8/8	0.69	0.21	73,84,86,87	0
2	BDF	C	302	12/12	0.76	0.28	37,67,71,81	0
3	IMD	A	303	5/5	0.79	0.46	97,97,98,100	0
2	BDF	A	301	12/12	0.80	0.23	64,72,74,75	0
6	CL	D	303[B]	1/1	0.88	0.20	58,58,58,58	1
6	CL	D	303[A]	1/1	0.88	0.20	57,57,57,57	1
2	BDF	C	301	12/12	0.89	0.17	50,57,61,61	0
5	CIT	E	301	13/13	0.92	0.12	45,51,56,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.