



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2023 – 11:39 AM EDT

PDB ID : 7U9E
Title : Pfs230 D1 domain in complex with 230AL-26
Authors : Tang, W.K.; Tolia, N.H.
Deposited on : 2022-03-10
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	 12% 92% 6%
2	B	261	 9% 81% 6% 13%
3	H	257	 15% 86% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9700 atoms, of which 4782 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gametocyte surface protein P230.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	180	2875	917	1440	223	291	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	585	GLN	ASN	conflict	UNP P68874

- Molecule 2 is a protein called LMIV230-01.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	227	3470	1106	1723	297	337	7	0	0	0

- Molecule 3 is a protein called 230AL-26.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	H	226	3294	1044	1619	291	334	6	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		
4	B	18	Total	O	0	0
			18	18		
4	H	13	Total	O	0	0
			13	13		

4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	106.59Å 150.94Å 103.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.81 – 2.39 19.81 – 2.39	Depositor EDS
% Data completeness (in resolution range)	95.0 (19.81-2.39) 95.0 (19.81-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 2.38Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.225 , 0.269 0.225 , 0.270	Depositor DCC
R_{free} test set	1588 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	56.6	Xtrriage
Anisotropy	0.346	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9700	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1461	0.49	0/1976
2	B	0.27	0/1786	0.51	0/2418
3	H	0.26	0/1711	0.51	0/2327
All	All	0.26	0/4958	0.50	0/6721

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1435	1440	1440	4	0
2	B	1747	1723	1724	10	0
3	H	1675	1619	1621	3	0
4	A	30	0	0	0	0
4	B	18	0	0	0	0
4	H	13	0	0	0	0
All	All	4918	4782	4785	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:67:ARG:NH2	2:B:90:ASP:OD2	2.28	0.67
3:H:139:SER:HA	3:H:233:LEU:HD11	1.83	0.59
2:B:180:GLN:HB2	2:B:190:LEU:HD11	1.86	0.56
1:A:663:LEU:HG	1:A:670:LEU:HD22	1.88	0.55
2:B:35:ILE:HD12	2:B:105:ALA:HB2	1.92	0.50
2:B:35:ILE:HD13	2:B:239:TRP:CZ3	2.47	0.49
1:A:554:VAL:HG12	1:A:555:ASP:H	1.78	0.49
1:A:554:VAL:O	1:A:555:ASP:HB2	2.12	0.49
2:B:154:LEU:HD23	2:B:247:LEU:HD13	1.95	0.48
1:A:554:VAL:HG12	1:A:555:ASP:N	2.30	0.46
3:H:105:ASP:HA	3:H:184:LEU:HD22	1.99	0.43
2:B:193:LYS:O	2:B:194:ALA:HB3	2.19	0.42
3:H:6:GLU:HG3	3:H:96:CYS:SG	2.60	0.42
2:B:104:VAL:CG1	2:B:234:TYR:CZ	3.04	0.41
2:B:250:LYS:O	2:B:251:ARG:C	2.59	0.40
2:B:35:ILE:CD1	2:B:105:ALA:HB2	2.51	0.40
2:B:221:LEU:HD21	2:B:247:LEU:HD21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/191 (93%)	171 (96%)	7 (4%)	0	100	100
2	B	223/261 (85%)	213 (96%)	10 (4%)	0	100	100
3	H	222/257 (86%)	213 (96%)	9 (4%)	0	100	100
All	All	623/709 (88%)	597 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/177 (95%)	167 (99%)	1 (1%)	86	94
2	B	188/200 (94%)	188 (100%)	0	100	100
3	H	182/195 (93%)	182 (100%)	0	100	100
All	All	538/572 (94%)	537 (100%)	1 (0%)	93	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	726	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/191 (94%)	0.86	23 (12%) 3 3	48, 59, 95, 211	0
2	B	227/261 (86%)	0.86	23 (10%) 7 6	47, 59, 89, 113	0
3	H	226/257 (87%)	0.93	38 (16%) 1 1	47, 65, 102, 129	0
All	All	633/709 (89%)	0.88	84 (13%) 3 3	47, 61, 97, 211	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	224	TYR	7.1
1	A	554	VAL	5.6
2	B	118	SER	5.5
2	B	36	TRP	5.4
2	B	251	ARG	5.1
3	H	213	ILE	5.1
3	H	139	SER	4.8
1	A	599	LEU	4.8
2	B	143	SER	4.7
1	A	556	GLU	4.3
3	H	206	GLY	4.2
3	H	247	LEU	4.2
2	B	48	MET	4.1
3	H	88	VAL	3.9
1	A	557	LEU	3.8
3	H	173	TRP	3.7
3	H	178	PRO	3.7
3	H	225	TYR	3.7
2	B	57	ARG	3.5
1	A	555	ASP	3.4
1	A	668	TYR	3.3
2	B	177	ALA	3.3
2	B	87	ARG	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	183	PRO	3.3
1	A	634	LYS	3.3
3	H	205	SER	3.2
1	A	588	ASN	3.2
3	H	11	LEU	3.2
3	H	198	ASP	3.2
1	A	627	PRO	3.1
1	A	717	LYS	3.1
1	A	715	ASN	3.1
3	H	41	PRO	3.1
1	A	714	ASP	3.1
1	A	559	LYS	3.1
1	A	721	GLY	3.1
3	H	226	CYS	3.0
2	B	56	ASP	3.0
3	H	36	TRP	3.0
1	A	706	CYS	2.9
2	B	62	GLN	2.9
2	B	116	VAL	2.9
3	H	180	THR	2.9
3	H	179	GLY	2.9
2	B	35	ILE	2.9
2	B	115	THR	2.8
3	H	155	VAL	2.8
1	A	703	TYR	2.8
3	H	1	GLU	2.8
3	H	208	SER	2.8
2	B	247	LEU	2.7
3	H	160	SER	2.7
3	H	227	ALA	2.7
3	H	211	LEU	2.7
2	B	184	GLY	2.6
2	B	83	LEU	2.6
2	B	240	THR	2.6
3	H	34	MET	2.6
1	A	604	SER	2.5
2	B	178	TRP	2.5
2	B	81	MET	2.5
3	H	154	ARG	2.4
3	H	10	GLY	2.4
1	A	702	PHE	2.4
2	B	100	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	H	13	LYS	2.4
3	H	200	PHE	2.4
1	A	722	ILE	2.4
3	H	153	GLN	2.3
2	B	229	TYR	2.3
3	H	117	SER	2.3
1	A	662	LEU	2.3
3	H	2	VAL	2.2
2	B	94	TYR	2.1
1	A	704	PHE	2.1
3	H	65	LYS	2.1
3	H	177	VAL	2.1
1	A	728	GLU	2.1
3	H	223	ASP	2.1
3	H	150	THR	2.1
3	H	195	GLY	2.1
3	H	216	LEU	2.1
3	H	232	ASN	2.0
1	A	595	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.