



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2023 – 10:52 AM EDT

PDB ID : 5UCP  
Title : Class II fructose-1,6-bisphosphate aldolase E142A variant of *Helicobacter pylori* with FBP and cleavage products  
Authors : Jacques, B.; Sygusch, J.  
Deposited on : 2016-12-22  
Resolution : 1.44 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2021 (1.46-1.42)
Clashscore	141614	2086 (1.46-1.42)
Ramachandran outliers	138981	2047 (1.46-1.42)
Sidechain outliers	138945	2047 (1.46-1.42)
RSRZ outliers	127900	1993 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	
1	B	307	

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 9696 atoms, of which 4477 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructose-bisphosphate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	292	4542	1471	2230	401	428	12	0	4	0
1	B	293	4539	1474	2222	402	429	12	0	4	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	48	ALA	THR	conflict	UNP P56109
A	67	ILE	THR	conflict	UNP P56109
A	142	ALA	GLU	engineered mutation	UNP P56109
B	48	ALA	THR	conflict	UNP P56109
B	67	ILE	THR	conflict	UNP P56109
B	142	ALA	GLU	engineered mutation	UNP P56109


- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Zn	0	3
			3	3		
3	B	3	Total	Zn	0	3
			3	3		

- Molecule 4 is 1,6-di-O-phosphono-D-fructose (three-letter code: P6F) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>12</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	P		
4	A	1	30	6	10	12	2	0	1

- Molecule 5 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula:  $C_3H_7O_6P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	P		
5	B	1	15	3	5	6	1	0	0

- Molecule 6 is GLYCERALDEHYDE-3-PHOSPHATE (three-letter code: G3H) (formula:  $C_3H_7O_6P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	P		
6	B	1	30	6	10	12	2	0	1


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	271	Total	O	0	1
			271	271		
7	B	261	Total	O	0	0
			261	261		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fructose-bisphosphate aldolase



- Molecule 1: Fructose-bisphosphate aldolase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	39.48Å 64.63Å 62.84Å 82.15° 74.39° 75.78°	Depositor
Resolution (Å)	29.21 – 1.44 29.21 – 1.44	Depositor EDS
% Data completeness (in resolution range)	82.2 (29.21-1.44) 82.2 (29.21-1.44)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.35 (at 1.44Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575)	Depositor
R, $R_{free}$	0.162 , 0.196 0.162 , 0.196	Depositor DCC
$R_{free}$ test set	4503 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.6	Xtrriage
Anisotropy	0.148	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 46.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.196 for -h,-l,-k	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9696	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 13P, G3H, NA, P6F, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/2356	0.51	0/3166
1	B	0.33	0/2361	0.51	0/3173
All	All	0.34	0/4717	0.51	0/6339

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2312	2230	2320	9	0
1	B	2317	2222	2325	12	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	A	20	10	10	1	0
5	B	10	5	4	3	0
6	B	20	10	10	2	0
7	A	271	0	0	4	3
7	B	261	0	0	2	2
All	All	5219	4477	4669	23	4



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:185:PHE:CD2	1:B:189:PRO:HA	2.30	0.66
1:A:104:ASP:OD2	7:A:501:HOH:O	2.15	0.64
1:A:7:GLU:OE2	7:A:502:HOH:O	2.15	0.63
5:B:405:13P:O3	6:B:406[C]:G3H:O1	2.22	0.58
1:A:306:LYS:NZ	7:A:504:HOH:O	2.36	0.57
1:B:137:ARG:O	1:B:152:ALA:HB1	2.07	0.54
1:A:170:ASP:OD1	7:A:503:HOH:O	2.19	0.53
1:B:137:ARG:C	1:B:152:ALA:HB1	2.30	0.52
1:B:242:GLN:NE2	7:B:503:HOH:O	2.21	0.52
1:B:182:ALA:HB1	1:B:233:SER:HB2	1.95	0.47
1:B:104:ASP:OD2	7:B:502:HOH:O	2.21	0.46
1:B:253:ASN:HB3	5:B:405:13P:H31	1.97	0.46
1:A:28:GLU:HG2	1:A:283:PHE:HB3	1.98	0.45
1:B:185:PHE:CG	1:B:189:PRO:HA	2.51	0.45
5:B:405:13P:H32	6:B:406[C]:G3H:O2	2.18	0.44
1:B:28:GLU:HG2	1:B:283:PHE:HB3	1.98	0.44
1:B:183:PHE:CE1	1:B:231:LYS:HB2	2.52	0.44
1:B:137:ARG:HB2	1:B:153:VAL:HG12	2.00	0.44
1:A:180:HIS:HD2	4:A:405[A]:P6F:H4	1.83	0.43
1:B:205:ILE:HB	1:B:206:PRO:HD2	2.02	0.42
1:A:206:PRO:HA	1:A:250:ASN:OD1	2.19	0.42
1:A:17:TYR:CE2	1:A:306:LYS:HE2	2.55	0.41
1:A:205:ILE:HB	1:A:206:PRO:HD2	2.02	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:504:HOH:O	7:A:508:HOH:O[1_655]	1.99	0.21
7:A:513:HOH:O	7:B:510:HOH:O[1_565]	2.01	0.19
7:B:552:HOH:O	7:B:710:HOH:O[1_455]	2.09	0.11
7:A:504:HOH:O	7:A:698:HOH:O[1_655]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/307 (96%)	291 (99%)	2 (1%)	1 (0%)	41	19
1	B	295/307 (96%)	290 (98%)	5 (2%)	0	100	100
All	All	589/614 (96%)	581 (99%)	7 (1%)	1 (0%)	47	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	231	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/253 (98%)	243 (98%)	4 (2%)	62	31
1	B	247/253 (98%)	244 (99%)	3 (1%)	71	43
All	All	494/506 (98%)	487 (99%)	7 (1%)	65	37

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	PHE
1	A	102	MET
1	A	165	LYS
1	A	171	TYR
1	B	45	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	102	MET
1	B	171	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	G3H	B	406[C]	3	8,9,9	0.56	0	10,12,12	1.25	0
5	13P	B	405	3	9,9,9	0.60	0	10,12,12	1.64	4 (40%)
4	P6F	A	405[A]	-	18,19,19	0.68	0	21,28,28	0.96	0
6	G3H	B	406[A]	-	8,9,9	0.60	0	10,12,12	1.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	G3H	B	406[C]	3	-	5/7/8/8	-
5	13P	B	405	3	-	5/7/8/8	-
4	P6F	A	405[A]	-	-	6/24/24/24	-
6	G3H	B	406[A]	-	-	3/7/8/8	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	405	13P	O2-C2-C3	-2.69	116.67	120.77
5	B	405	13P	O3-C3-C2	-2.66	103.56	112.43
5	B	405	13P	O3P-P-O2P	2.10	115.66	107.64
5	B	405	13P	O3P-P-O1	-2.10	101.16	106.73

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	405[A]	P6F	C2-C3-C4-O4
4	A	405[A]	P6F	C2-C3-C4-C5
4	A	405[A]	P6F	O3-C3-C4-O4
4	A	405[A]	P6F	O3-C3-C4-C5
4	A	405[A]	P6F	C6-O6-P2-O4P
5	B	405	13P	C1-O1-P-O2P
5	B	405	13P	C1-O1-P-O3P
6	B	406[A]	G3H	C1-C2-C3-O1P
6	B	406[A]	G3H	O2-C2-C3-O1P
6	B	406[C]	G3H	C3-O1P-P-O2P
6	B	406[C]	G3H	C3-O1P-P-O4P
5	B	405	13P	O1-C1-C2-O2
5	B	405	13P	C1-O1-P-O1P
5	B	405	13P	O1-C1-C2-C3
4	A	405[A]	P6F	C6-O6-P2-O5P
6	B	406[A]	G3H	C3-O1P-P-O2P
6	B	406[C]	G3H	C1-C2-C3-O1P
6	B	406[C]	G3H	C3-O1P-P-O3P
6	B	406[C]	G3H	O2-C2-C3-O1P

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	406[C]	G3H	2	0
5	B	405	13P	3	0
4	A	405[A]	P6F	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/307 (95%)	-0.09	16 (5%) 25 25	10, 18, 54, 154	0
1	B	293/307 (95%)	0.00	18 (6%) 21 20	11, 18, 62, 142	0
All	All	585/614 (95%)	-0.05	34 (5%) 23 23	10, 18, 56, 154	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	152	ALA	13.3
1	B	183	PHE	10.1
1	B	153	VAL	5.9
1	B	187	GLY	5.5
1	A	183	PHE	5.3
1	A	153	VAL	5.3
1	A	187	GLY	5.1
1	B	182	ALA	4.8
1	B	1	MET	4.5
1	B	186	LYS	4.3
1	A	186	LYS	4.3
1	A	185	PHE	4.2
1	B	190	LYS	4.0
1	B	189	PRO	3.7
1	A	1	MET	3.4
1	A	231	LYS	3.1
1	B	137	ARG	3.0
1	A	188	GLU	3.0
1	B	191	LEU	3.0
1	B	185	PHE	2.9
1	A	191	LEU	2.9
1	B	181	GLY	2.8
1	A	45	PHE	2.8
1	A	180	HIS	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	180	HIS	2.6
1	B	188	GLU	2.6
1	A	137	ARG	2.5
1	A	182	ALA	2.5
1	B	184	LYS	2.4
1	A	181	GLY	2.3
1	B	179	SER	2.3
1	B	45	PHE	2.2
1	A	190	LYS	2.2
1	A	240	PHE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	G3H	B	406[A]	10/10	0.90	0.19	27,30,35,35	15
6	G3H	B	406[C]	10/10	0.90	0.19	20,26,32,34	15
5	13P	B	405	10/10	0.92	0.13	21,29,35,35	15
4	P6F	A	405[A]	20/20	0.94	0.12	17,26,33,36	30
3	ZN	A	404[C]	1/1	0.96	0.16	21,21,21,21	1
3	ZN	B	404[C]	1/1	0.97	0.08	23,23,23,23	1
2	NA	B	401	1/1	0.98	0.09	24,24,24,24	0
3	ZN	A	402[A]	1/1	0.98	0.03	29,29,29,29	1
2	NA	A	401	1/1	0.98	0.12	27,27,27,27	0
3	ZN	B	402[A]	1/1	0.98	0.03	25,25,25,25	1
3	ZN	B	403[B]	1/1	0.98	0.06	29,29,29,29	1
3	ZN	A	403[B]	1/1	0.99	0.03	20,20,20,20	1



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.