

Full wwPDB X-ray Structure Validation Report (i)

Sep 29, 2024 – 09:45 AM EDT

PDB ID	:	3UDX	
Title	:	Crystal structure of Acinetobacter baumannii PBP1a in complex with	
		Imipenem	
Authors	:	Han, S.	
Deposited on	:	2011-10-28	
Resolution	:	2.50 Å(reported)	

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))		
Rfree	164625	5504 (2.50-2.50)		
Clashscore	180529	6282 (2.50-2.50)		
Ramachandran outliers	177936	6191 (2.50-2.50)		
Sidechain outliers	177891	6193 (2.50-2.50)		
RSRZ outliers	164620	5504 (2.50-2.50)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	А	731	^{2%} 70%	11%	•	18%			
1	В	731	73%	9%		19%			



3UDX

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9771 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Penicillin-binding protein 1a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	508	Total	С	Ν	0	\mathbf{S}	0	0	0
1	Л	090	4715	3006	832	861	16	0	0	0
1	р	505	Total	С	Ν	0	S	0	1	0
	D	090	4691	2990	829	856	16	0		

Chain	Residue	Modelled	Actual	Comment	Reference
А	9	MET	-	expression tag	UNP G1C794
А	10	HIS	-	expression tag	UNP G1C794
А	11	HIS	-	expression tag	UNP G1C794
А	12	HIS	-	expression tag	UNP G1C794
А	13	HIS	-	expression tag	UNP G1C794
А	14	HIS	-	expression tag	UNP G1C794
А	15	HIS	-	expression tag	UNP G1C794
А	16	GLU	-	expression tag	UNP G1C794
А	17	ASN	-	expression tag	UNP G1C794
А	18	LEU	-	expression tag	UNP G1C794
А	19	TYR	-	expression tag	UNP G1C794
А	20	PHE	-	expression tag	UNP G1C794
А	21	GLN	-	expression tag	UNP G1C794
А	22	SER	-	expression tag	UNP G1C794
А	23	HIS	-	expression tag	UNP G1C794
А	24	MET	-	expression tag	UNP G1C794
В	9	MET	-	expression tag	UNP G1C794
В	10	HIS	-	expression tag	UNP G1C794
В	11	HIS	-	expression tag	UNP G1C794
В	12	HIS	-	expression tag	UNP G1C794
В	13	HIS	-	expression tag	UNP G1C794
В	14	HIS	-	expression tag	UNP G1C794
В	15	HIS	-	expression tag	UNP G1C794
В	16	GLU	-	expression tag	UNP G1C794
В	17	ASN	-	expression tag	UNP G1C794

There are 32 discrepancies between the modelled and reference sequences:



Contentia	continuad from proto de pagem									
Chain	Residue	Modelled Actual		Comment	Reference					
В	18	LEU	-	expression tag	UNP G1C794					
В	19	TYR	-	expression tag	UNP G1C794					
В	20	PHE	-	expression tag	UNP G1C794					
В	21	GLN	-	expression tag	UNP G1C794					
В	22	SER	-	expression tag	UNP G1C794					
В	23	HIS	-	expression tag	UNP G1C794					
В	24	MET	-	expression tag	UNP G1C794					

• Molecule 2 is (5R)-5-[(1S,2R)-1-formyl-2-hydroxypropyl]-3-[(2-{[(E)-iminomethyl]amino}e thyl)sulfanyl]-4,5-dihydro-1H-pyrrole-2-carbox ylic acid (three-letter code: IM2) (formula: $C_{12}H_{19}N_3O_4S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
9	Λ	1	Total	С	Ν	0	\mathbf{S}	0	0	
	A	1	20	12	3	4	1	0	0	
0	В	1	Total	С	Ν	0	S	0	0	
	D	L	20	12	3	4	1	0	U	

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	133	Total O 133 133	0	0
3	В	192	Total O 192 192	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Penicillin-binding protein 1a



F511 ● F512 ● F520 F





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	119.27Å 242.93Å 49.12Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	48.02 - 2.50	Depositor
	48.02 - 2.50	EDS
% Data completeness	93.1 (48.02-2.50)	Depositor
(in resolution range)	93.5(48.02-2.50)	EDS
R_{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.55 (at 2.51 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.9.6	Depositor
R R.	0.187 , 0.238	Depositor
II, II, <i>free</i>	0.188 , 0.245	DCC
R_{free} test set	2376 reflections $(5.04%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	43.8	Xtriage
Anisotropy	0.343	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 66.9	EDS
L-test for $twinning^2$	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9771	wwPDB-VP
Average B, all atoms $(Å^2)$	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $\mathrm{IM}2$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
			# Z > 5	RMSZ	# Z > 5	
1	А	0.49	0/4824	0.70	0/6541	
1	В	0.53	0/4803	0.73	0/6514	
All	All	0.51	0/9627	0.71	0/13055	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4715	0	4701	35	0
1	В	4691	0	4673	31	0
2	А	20	0	16	0	0
2	В	20	0	16	0	0
3	А	133	0	0	0	0
3	В	192	0	0	0	0
All	All	9771	0	9406	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:B:298:ARG:H	1:B:390:GLN:HE22	1.17	0.88
1:A:298:ARG:H	1:A:390:GLN:HE22	1.23	0.83
1:B:356:ASN:HD21	1:B:704:ARG:H	1.38	0.71
1:A:323:LYS:HE3	1:A:325:ASN:HD21	1.58	0.68
1:A:505:ARG:HH22	1:A:515:GLU:HG3	1.58	0.68
1:A:513:LEU:HB3	1:A:518:ILE:HD11	1.79	0.65
1:B:406:GLY:O	1:B:552:PRO:HA	1.97	0.64
1:B:704:ARG:O	1:B:705:ARG:HB2	1.96	0.64
1:B:528:THR:N	1:B:529:PRO:HD2	2.13	0.63
1:B:356:ASN:ND2	1:B:704:ARG:HG2	2.18	0.59
1:A:288:LEU:HD22	1:A:394:VAL:HG13	1.85	0.59
1:A:556:GLN:HE21	1:A:557:ARG:HH11	1.50	0.58
1:B:376:ARG:HD2	1:B:391:VAL:HG23	1.86	0.57
1:B:471:ASP:HB2	1:B:473:ARG:HH11	1.71	0.56
1:A:496:LEU:HD22	1:A:525:ALA:HB2	1.87	0.56
1:A:71:PHE:O	1:B:553:HIS:HA	2.08	0.54
1:A:695:GLY:HA2	1:A:702:LEU:HD21	1.90	0.53
1:A:661:ILE:HD11	1:A:717:ILE:HA	1.92	0.52
1:A:351:PRO:HG2	1:A:359:GLY:HA3	1.92	0.52
1:B:431:GLN:HG2	1:B:528:THR:O	2.10	0.51
1:A:31:VAL:HB	1:A:40:ALA:HB3	1.93	0.51
1:A:261:ALA:HA	1:B:144:TYR:CE1	2.45	0.50
1:A:528:THR:N	1:A:529:PRO:HD2	2.26	0.50
1:B:528:THR:H	1:B:529:PRO:HD2	1.75	0.49
1:A:513:LEU:HD21	1:A:531:VAL:HG12	1.93	0.49
1:A:461:THR:HG22	1:A:466:THR:OG1	2.12	0.49
1:A:313:TYR:CE1	1:A:419:GLN:HB2	2.48	0.49
1:B:510:ASP:O	1:B:549:ARG:HD3	2.13	0.48
1:B:473:ARG:O	1:B:488:ARG:NH2	2.46	0.48
1:A:148:ILE:HG21	1:A:184:PRO:HB3	1.94	0.47
1:A:514:GLN:HB2	1:A:517:GLN:HG2	1.96	0.47
1:A:640:TYR:HD1	1:A:732:VAL:HG23	1.81	0.46
1:B:72:PHE:HB3	1:B:140:ILE:HD12	1.96	0.46
1:B:298:ARG:H	1:B:390:GLN:NE2	1.99	0.46
1:A:298:ARG:H	1:A:390:GLN:NE2	2.03	0.46
1:A:137:LYS:NZ	1:B:256:HIS:HD2	2.14	0.46
1:A:558:ILE:HB	1:A:567:TYR:HB3	1.98	0.46
1:B:229:ILE:HD12	1:B:231:LEU:H	1.80	0.45
1:B:371:VAL:O	1:B:372:LYS:HB2	2.17	0.45
1:A:144:TYR:O	1:A:148:ILE:HG13	2.18	0.44
1:A:29:LEU:HD22	1:A:266:TYR:O	2.18	0.44



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:406:GLY:O	1:A:552:PRO:HA	2.18	0.44
1:B:528:THR:N	1:B:529:PRO:CD	2.81	0.43
1:B:310:PHE:O	1:B:334:GLN:NE2	2.52	0.42
1:A:303:HIS:HD2	1:A:373:ASP:OD1	2.02	0.42
1:B:483:ALA:HB1	1:B:491:VAL:HG11	2.01	0.42
1:B:191:ASN:HB3	1:B:194:VAL:HG12	2.02	0.42
1:B:145:VAL:HG22	1:B:157:ILE:HG12	2.01	0.42
1:A:483:ALA:HA	1:A:488:ARG:HG2	2.01	0.42
1:B:356:ASN:HD21	1:B:704:ARG:N	2.12	0.42
1:B:704:ARG:O	1:B:705:ARG:CB	2.65	0.42
1:B:323:LYS:HB3	1:B:330:GLU:HB2	2.02	0.41
1:A:72:PHE:HB3	1:A:140:ILE:HD12	2.02	0.41
1:A:316:THR:HG22	1:A:389:VAL:HG21	2.02	0.41
1:A:484:LEU:HD23	1:A:645:ILE:HG21	2.01	0.41
1:A:383:LYS:HB3	1:A:384:THR:H	1.66	0.41
1:B:496:LEU:HD23	1:B:522:TYR:HD1	1.86	0.41
1:A:356:ASN:OD1	1:A:704:ARG:HB2	2.21	0.40
1:A:380:ASN:HB2	1:A:383:LYS:O	2.21	0.40
1:A:319:ALA:HB1	1:A:331:ALA:HB1	2.03	0.40
1:B:166:ASN:HB2	1:B:234:PRO:HD3	2.03	0.40
1:B:195:ASN:HB3	1:B:198:ARG:HB3	2.03	0.40
1:B:210:MET:HB3	1:B:216:ILE:HG12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	590/731~(81%)	564 (96%)	23~(4%)	3~(0%)	25	44
1	В	588/731~(80%)	562~(96%)	24~(4%)	2~(0%)	37	56
All	All	1178/1462~(81%)	1126 (96%)	47 (4%)	5~(0%)	30	49



All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	349	ALA
1	А	383	LYS
1	В	349	ALA
1	В	705	ARG
1	А	379	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	490/608 (81%)	465~(95%)	25~(5%)	20 40		
1	В	487/608~(80%)	467 (96%)	20 (4%)	26 50		
All	All	977/1216 (80%)	932~(95%)	45 (5%)	23 45		

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	59	PHE
1	А	133	GLN
1	А	139	ASP
1	А	195	ASN
1	А	198	ARG
1	А	240	ASN
1	А	263	ASP
1	А	275	LYS
1	А	311	ARG
1	А	352	TYR
1	А	394	VAL
1	А	401	ILE
1	А	421	LYS
1	А	447	ARG
1	А	470	SER
1	А	481	ARG
1	А	511	PHE
1	А	556	GLN



Mol	Chain	Res	Type
1	А	568	GLU
1	А	579	ILE
1	А	635	LYS
1	А	650	ILE
1	А	675	ASP
1	А	685	ASN
1	А	732	VAL
1	В	27	LYS
1	В	35	ASP
1	В	53	LYS
1	В	59	PHE
1	В	68	ASP
1	В	135	LEU
1	В	142	SER
1	В	189	LYS
1	В	209	ARG
1	В	218	GLN
1	В	225	VAL
1	В	240	ASN
1	В	520	ARG
1	В	556	GLN
1	В	626	GLN
1	В	627	TYR
1	В	650	ILE
1	В	700	THR
1	В	710	ILE
1	В	733	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	235	ASN
1	А	295	HIS
1	А	303	HIS
1	А	325	ASN
1	А	390	GLN
1	А	551	GLN
1	А	556	GLN
1	А	629	GLN
1	А	652	HIS
1	В	212	GLN
1	В	256	HIS



Mol	Chain	Res	Type
1	В	356	ASN
1	В	390	GLN
1	В	469	ASN
1	В	514	GLN
1	В	551	GLN
1	В	556	GLN
1	В	629	GLN
1	В	674	ASN
1	В	722	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tink	Bo	ond leng	$_{\rm sths}$	B	ond ang	gles
	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IM2	А	998	1	15,20,20	0.90	0	8,26,26	1.92	2 (25%)
2	IM2	В	999	1	15,20,20	0.85	0	8,26,26	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IM2	А	998	1	-	5/18/32/32	0/1/1/1
2	IM2	В	999	1	-	7/18/32/32	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	998	IM2	O32-C31-C3	3.40	122.13	116.73
2	А	998	IM2	C3-C2-S21	2.71	128.19	124.37

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
2	А	998	IM2	C1-C5-C6-C7
2	А	998	IM2	C1-C5-C6-C61
2	А	998	IM2	C7-C6-C61-C62
2	А	998	IM2	C23-C22-S21-C2
2	А	998	IM2	N26-C25-N24-C23
2	В	999	IM2	C1-C5-C6-C7
2	В	999	IM2	C1-C5-C6-C61
2	В	999	IM2	N4-C3-C31-O31
2	В	999	IM2	N4-C3-C31-O32
2	В	999	IM2	N26-C25-N24-C23
2	В	999	IM2	C2-C3-C31-O31
2	В	999	IM2	C2-C3-C31-O32

All (12) torsion outliers are listed below:

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the



average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	598/731~(81%)	0.01	12 (2%) 64 62	31, 53, 88, 122	0
1	В	595/731~(81%)	-0.11	20 (3%) 48 45	21, 46, 93, 120	1 (0%)
All	All	1193/1462~(81%)	-0.05	32 (2%) 56 52	21, 50, 93, 122	1 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	В	511	PHE	4.6	
1	А	579	ILE	4.2	
1	А	654	THR	3.9	
1	В	627	TYR	3.7	
1	В	194	VAL	3.5	
1	В	653	GLY	3.4	
1	В	707	TYR	3.3	
1	В	581	ALA	3.3	
1	В	470	SER	3.2	
1	А	709	GLY	3.1	
1	А	513	LEU	3.0	
1	В	462	ILE	2.9	
1	В	661	ILE	2.8	
1	А	512	GLY	2.8	
1	В	465	TRP	2.7	
1	В	626	GLN	2.7	
1	В	507	LEU	2.7	
1	А	736	LYS	2.7	
1	В	135	LEU	2.7	
1	В	460	ILE	2.6	
1	В	733	ARG	2.6	
1	В	229	ILE	2.5	
1	A	627	TYR	2.5	
1	В	640	TYR	2.3	



Mol	Chain	Res	Type	RSRZ
1	А	626	GLN	2.3
1	А	383	LYS	2.3
1	В	190	TYR	2.3
1	А	733	ARG	2.2
1	А	578	CYS	2.2
1	В	189	LYS	2.2
1	В	513	LEU	2.1
1	А	734	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	IM2	В	999	20/20	0.88	0.13	46,70,98,99	0
2	IM2	А	998	20/20	0.90	0.13	52,76,95,95	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers (i)

There are no such residues in this entry.

