



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2021 – 04:33 AM EDT

PDB ID : 1ULZ
Title : Crystal structure of the biotin carboxylase subunit of pyruvate carboxylase
Authors : Kondo, S.; Nakajima, Y.; Sugio, S.; Yong-Biao, J.; Sueda, S.; Kondo, H.
Deposited on : 2003-09-18
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3840 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called pyruvate carboxylase n-terminal domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	451	3565	2273	608	668	16	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	VAL	PHE	engineered mutation	UNP O67483
A	352	ARG	GLY	SEE REMARK 999	UNP O67483
A	405	ALA	VAL	SEE REMARK 999	UNP O67483

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	275	Total	O	0	0
			275	275		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.42Å 122.14Å 59.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20 28.68 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.20) 99.6 (28.68-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.00 (at 2.20Å)	Xtrriage
Refinement program	CNX	Depositor
R, R_{free}	0.199 , 0.236 0.198 , 0.229	Depositor DCC
R_{free} test set	1715 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtrriage
Anisotropy	0.091	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 54.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3840	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/3634	0.57	0/4906

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3565	0	3593	73	0
2	A	275	0	0	2	0
All	All	3840	0	3593	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:438:LYS:HB3	1:A:438:LYS:NZ	1.98	0.78
1:A:177:VAL:O	1:A:181:GLU:HG3	1.91	0.71
1:A:258:VAL:HA	1:A:273:MET:HE1	1.70	0.71
1:A:105:HIS:CD2	1:A:107:LYS:H	2.10	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:HIS:HD2	1:A:107:LYS:H	1.40	0.69
1:A:297:PRO:O	1:A:301:MET:HG2	1.94	0.67
1:A:258:VAL:HG22	1:A:273:MET:CE	2.26	0.65
1:A:352:ARG:HB2	1:A:414:VAL:HG12	1.81	0.63
1:A:258:VAL:HG22	1:A:273:MET:HE2	1.80	0.62
1:A:150:ILE:O	1:A:200:LYS:HE2	1.98	0.62
1:A:111:LEU:HD21	1:A:121:VAL:HG21	1.81	0.62
1:A:438:LYS:O	1:A:442:GLU:HG3	2.03	0.59
1:A:174:GLU:OE2	1:A:178:LYS:HE3	2.02	0.58
1:A:1:MET:O	1:A:1:MET:HG3	2.03	0.58
1:A:111:LEU:HD22	1:A:112:MET:HE2	1.85	0.58
1:A:451:GLU:HG2	1:A:451:GLU:OXT	2.03	0.57
1:A:184:SER:O	1:A:188:GLU:HG3	2.07	0.55
1:A:249:GLU:CD	1:A:249:GLU:H	2.11	0.54
1:A:438:LYS:HB3	1:A:438:LYS:HZ1	1.72	0.54
1:A:438:LYS:NZ	1:A:438:LYS:CB	2.70	0.54
1:A:229:SER:O	1:A:437:THR:HA	2.08	0.54
1:A:258:VAL:HG13	1:A:273:MET:CE	2.39	0.53
1:A:204:ASN:ND2	1:A:438:LYS:HD3	2.24	0.52
1:A:19:ARG:HG3	1:A:19:ARG:HH11	1.75	0.52
1:A:336:CYS:HB3	1:A:421:LEU:HD13	1.91	0.52
1:A:250:LYS:HD3	2:A:626:HOH:O	2.10	0.51
1:A:443:HIS:O	1:A:446:VAL:HG22	2.11	0.51
1:A:296:HIS:CG	1:A:297:PRO:HD3	2.45	0.51
1:A:450:GLU:O	1:A:451:GLU:HB2	2.10	0.51
1:A:394:THR:OG1	1:A:397:GLU:HG3	2.11	0.51
1:A:353:VAL:HG22	1:A:376:GLU:HG3	1.93	0.50
1:A:105:HIS:HD2	1:A:107:LYS:HB2	1.77	0.50
1:A:341:GLU:O	1:A:343:PRO:HD3	2.12	0.49
1:A:257:ILE:HA	1:A:260:LYS:HE3	1.95	0.49
1:A:233:ARG:O	1:A:234:ASN:HB2	2.13	0.48
1:A:438:LYS:HB3	1:A:438:LYS:HZ2	1.78	0.48
1:A:111:LEU:HD22	1:A:112:MET:CE	2.42	0.48
1:A:104:PRO:HG3	1:A:290:THR:HB	1.95	0.48
1:A:111:LEU:HD13	1:A:112:MET:CE	2.44	0.48
1:A:403:ARG:O	1:A:407:GLU:HG3	2.13	0.47
1:A:434:LYS:HE3	1:A:434:LYS:HA	1.95	0.47
1:A:300:GLU:HG2	1:A:305:ILE:O	2.15	0.47
1:A:296:HIS:N	1:A:297:PRO:CD	2.78	0.47
1:A:168:ARG:NH1	2:A:571:HOH:O	2.48	0.46
1:A:335:GLU:HG3	1:A:389:ILE:HG13	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:395:TRP:O	1:A:399:VAL:HG23	2.15	0.46
1:A:191:PHE:CD1	1:A:191:PHE:N	2.84	0.46
1:A:436:THR:OG1	1:A:438:LYS:HG2	2.16	0.46
1:A:92:LYS:O	1:A:96:GLU:HG3	2.17	0.45
1:A:207:HIS:CD2	1:A:228:CYS:HB2	2.52	0.45
1:A:347:PHE:CD2	1:A:347:PHE:N	2.86	0.44
1:A:204:ASN:ND2	1:A:438:LYS:CD	2.81	0.44
1:A:277:ALA:HB2	1:A:283:LEU:CD2	2.47	0.44
1:A:63:GLN:NE2	1:A:67:ASN:OD1	2.50	0.44
1:A:378:THR:HB	1:A:379:PRO:HD2	2.00	0.43
1:A:418:ILE:O	1:A:422:ILE:HG13	2.18	0.43
1:A:431:LYS:HE2	1:A:431:LYS:HB3	1.81	0.43
1:A:140:LEU:HD23	1:A:140:LEU:O	2.19	0.43
1:A:6:LEU:C	1:A:6:LEU:HD23	2.40	0.43
1:A:254:TYR:O	1:A:258:VAL:HG23	2.19	0.43
1:A:232:ARG:C	1:A:232:ARG:HD2	2.39	0.42
1:A:233:ARG:HD3	1:A:233:ARG:HA	1.93	0.42
1:A:227:ASP:HB3	1:A:240:ILE:HG12	2.02	0.41
1:A:250:LYS:HE2	1:A:250:LYS:HB3	1.68	0.41
1:A:342:ASP:OD1	1:A:344:LYS:HB2	2.20	0.41
1:A:108:VAL:HG11	1:A:267:TYR:CD2	2.56	0.41
1:A:363:PHE:C	1:A:363:PHE:CD1	2.95	0.41
1:A:390:THR:HB	1:A:401:ARG:HG2	2.04	0.40
1:A:204:ASN:N	1:A:205:PRO:CD	2.84	0.40
1:A:258:VAL:HG13	1:A:273:MET:HE2	2.03	0.40
1:A:232:ARG:O	1:A:235:GLN:HB2	2.22	0.40
1:A:258:VAL:CA	1:A:273:MET:HE1	2.45	0.40
1:A:258:VAL:HG13	1:A:273:MET:HE1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	449/451 (100%)	433 (96%)	15 (3%)	1 (0%)	47 55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	269	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	372/372 (100%)	364 (98%)	8 (2%)	52 65

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	LEU
1	A	117	ARG
1	A	176	LEU
1	A	369	HIS
1	A	409	TYR
1	A	434	LYS
1	A	438	LYS
1	A	445	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	HIS
1	A	204	ASN
1	A	235	GLN
1	A	279	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	451/451 (100%)	-0.07	23 (5%) 28 26	16, 27, 48, 58	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	8.1
1	A	348	ALA	6.1
1	A	349	PRO	5.8
1	A	191	PHE	5.2
1	A	351	THR	5.0
1	A	346	ASN	4.7
1	A	350	SER	4.5
1	A	451	GLU	4.4
1	A	193	ARG	4.0
1	A	233	ARG	3.7
1	A	345	LYS	3.6
1	A	344	LYS	3.2
1	A	450	GLU	3.0
1	A	280	GLU	2.6
1	A	192	GLY	2.6
1	A	188	GLU	2.4
1	A	161	GLY	2.3
1	A	117	ARG	2.3
1	A	234	ASN	2.3
1	A	342	ASP	2.0
1	A	116	ALA	2.0
1	A	347	PHE	2.0
1	A	138	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.