



## wwPDB EM Validation Summary Report ⓘ

Nov 2, 2024 – 05:26 PM EDT

PDB ID : 8UOQ  
EMDB ID : EMD-42437  
Title : Composite map of PIC\_delta\_TFIIK form2  
Authors : Yang, C.; Murakami, K.  
Deposited on : 2023-10-20  
Resolution : 3.80 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113  
Mogul : 2022.3.0, CSD as543be (2022)  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

























Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	M	345	
2	A	1733	
3	B	1224	
4	C	318	
5	E	215	
6	F	155	
7	H	146	
8	I	122	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	J	70	
10	K	120	
11	L	70	
12	Q	735	
13	P	400	
14	S	309	
15	O	240	
16	U	286	
17	V	122	
18	W	482	
19	X	328	
20	D	221	
21	G	171	
22	0	778	
23	1	642	
24	4	338	
25	6	461	
26	7	843	
27	2	513	
28	5	72	
29	N	64	
30	T	64	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
33	SF4	0	801	-	-	X	-

## 2 Entry composition [i](#)

There are 33 unique types of molecules in this entry. The entry contains 70481 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription initiation factor IIB.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	M	279	2175	1382	373	403	17	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	A	1425	11167	7036	1948	2121	62	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	B	1166	9227	5823	1619	1729	56	0	0

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	C	265	2086	1312	347	414	13	0	0

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	214	1752	1111	309	321	11	0	0

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	87	705	451	119	132	3	0	0

- Molecule 7 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	H	135	Total	C	N	O	S	0	0
			1080	679	182	214	5		

- Molecule 8 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	I	114	Total	C	N	O	S	0	0
			927	571	168	178	10		

- Molecule 9 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	J	66	Total	C	N	O	S	0	0
			540	345	94	95	6		

- Molecule 10 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	K	115	Total	C	N	O	S	0	0
			924	593	157	172	2		

- Molecule 11 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	L	44	Total	C	N	O	S	0	0
			352	217	70	61	4		

- Molecule 12 is a protein called Transcription initiation factor IIF subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	Q	214	Total	C	N	O	S	0	0
			1619	1017	297	299	6		

- Molecule 13 is a protein called Transcription initiation factor IIF subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	P	179	Total	C	N	O	S	0	0
			1484	941	258	279	6		

- Molecule 14 is a protein called Transcription elongation factor S-II.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	S	164	Total	C	N	O	S	0	0
			1294	809	230	247	8		

- Molecule 15 is a protein called TATA-box-binding protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	181	Total	C	N	O	S	0	0
			1422	925	243	248	6		

- Molecule 16 is a protein called Transcription initiation factor IIA large subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	U	107	Total	C	N	O	S	0	0
			885	559	147	176	3		

- Molecule 17 is a protein called Transcription initiation factor IIA subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	V	104	Total	C	N	O	S	0	0
			815	511	136	164	4		

- Molecule 18 is a protein called Transcription initiation factor IIE subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	W	247	Total	C	N	O	S	0	0
			2010	1275	347	381	7		

- Molecule 19 is a protein called Transcription initiation factor IIE subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	X	160	Total	C	N	O	S	0	0
			1288	826	212	245	5		

- Molecule 20 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	D	168	Total	C	N	O	S	0	0
			1331	822	237	270	2		

- Molecule 21 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	G	171	Total	C	N	O	S	0	0
			1335	858	221	248	8		

- Molecule 22 is a protein called General transcription and DNA repair factor IIIH helicase subunit XPD/RAD3.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	0	752	Total	C	N	O	S	0	0
			6091	3882	1029	1142	38		

- Molecule 23 is a protein called General transcription and DNA repair factor IIIH subunit TFB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	1	417	Total	C	N	O	S	0	0
			3382	2139	587	640	16		

- Molecule 24 is a protein called General transcription and DNA repair factor IIIH subunit TFB4.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	4	292	Total	C	N	O	S	0	0
			2267	1449	376	428	14		

- Molecule 25 is a protein called General transcription and DNA repair factor IIIH subunit SSL1.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	6	355	Total	C	N	O	S	0	0
			2786	1765	481	512	28		

- Molecule 26 is a protein called General transcription and DNA repair factor IIIH helicase subunit XPB.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	7	608	Total	C	N	O	S	0	0
			4889	3110	847	906	26		

- Molecule 27 is a protein called General transcription and DNA repair factor IIIH subunit TFB2.



Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
27	2	445	3546	2291	585	654	16	0	0

- Molecule 28 is a protein called General transcription and DNA repair factor IIIH subunit TFB5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
28	5	66	498	314	89	93	2	0	0

- Molecule 29 is a DNA chain called non-template strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
29	N	63	1288	621	225	380	62	0	0

- Molecule 30 is a DNA chain called template strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
30	T	63	1291	619	236	373	63	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
T	-10	DC	DT	conflict	GB 2567904391

- Molecule 31 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
31	M	1	Total	Zn	0
			1	1	
31	A	2	Total	Zn	0
			2	2	
31	B	1	Total	Zn	0
			1	1	
31	C	1	Total	Zn	0
			1	1	
31	I	2	Total	Zn	0
			2	2	
31	J	1	Total	Zn	0
			1	1	

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
31	L	1	Total	Zn	0
			1	1	
31	S	1	Total	Zn	0
			1	1	
31	4	1	Total	Zn	0
			1	1	
31	6	4	Total	Zn	0
			4	4	

- Molecule 32 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
32	A	1	Total	Mg	0
			1	1	
32	7	1	Total	Mg	0
			1	1	

- Molecule 33 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).




Mol	Chain	Residues	Atoms			AltConf
33	0	1	Total	Fe	S	0
			8	4	4	


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcription initiation factor IIB




- Molecule 2: DNA-directed RNA polymerase II subunit RPB1








Chain F: 




- Molecule 7: DNA-directed RNA polymerases I, II, and III subunit RPABC3

Chain H: 




- Molecule 8: DNA-directed RNA polymerase II subunit RPB9

Chain I: 





- Molecule 9: DNA-directed RNA polymerases I, II, and III subunit RPABC5

Chain J: 




- Molecule 10: DNA-directed RNA polymerase II subunit RPB11

Chain K: 



- Molecule 11: DNA-directed RNA polymerases I, II, and III subunit RPABC4

Chain L: 



- Molecule 12: Transcription initiation factor IIF subunit alpha



















- Molecule 24: General transcription and DNA repair factor IIH subunit TFB4




- Molecule 25: General transcription and DNA repair factor IIH subunit SSL1




- Molecule 26: General transcription and DNA repair factor IIH helicase subunit XPB









• Molecule 28: General transcription and DNA repair factor IIIH subunit TFB5



• Molecule 29: non-template strand



• Molecule 30: template strand



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	90136	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.25	Depositor
Minimum defocus (nm)	750	Depositor
Maximum defocus (nm)	1750	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.054	Depositor
Minimum map value	0.000	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.01	Depositor
Map size ( $\text{\AA}$ )	414.72003, 414.72003, 414.72003	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.08, 1.08, 1.08	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	M	0.46	1/2204 (0.0%)	0.84	7/2963 (0.2%)
2	A	0.48	2/11368 (0.0%)	0.83	29/15383 (0.2%)
3	B	0.50	4/9402 (0.0%)	0.85	30/12680 (0.2%)
4	C	0.46	0/2124	0.82	4/2879 (0.1%)
5	E	0.45	0/1788	0.75	1/2406 (0.0%)
6	F	0.48	0/717	0.83	1/967 (0.1%)
7	H	0.58	1/1097 (0.1%)	0.88	2/1484 (0.1%)
8	I	0.45	0/945	0.79	2/1273 (0.2%)
9	J	0.83	3/549 (0.5%)	1.65	11/738 (1.5%)
10	K	0.44	0/942	0.74	0/1272
11	L	0.56	0/354	1.03	2/468 (0.4%)
12	Q	0.47	0/1648	0.85	3/2226 (0.1%)
13	P	0.45	0/1511	0.84	6/2035 (0.3%)
14	S	0.47	1/1317 (0.1%)	0.76	3/1778 (0.2%)
15	O	0.39	0/1449	0.72	1/1952 (0.1%)
16	U	0.71	2/898 (0.2%)	0.82	1/1212 (0.1%)
17	V	0.54	0/822	0.95	6/1109 (0.5%)
18	W	0.39	0/2045	0.74	6/2757 (0.2%)
19	X	0.38	0/1312	0.63	1/1767 (0.1%)
20	D	0.40	0/1339	0.69	2/1793 (0.1%)
21	G	0.50	2/1363 (0.1%)	0.78	2/1840 (0.1%)
22	0	0.43	2/6209 (0.0%)	0.72	10/8384 (0.1%)
23	1	0.45	1/3434 (0.0%)	0.75	7/4624 (0.2%)
24	4	0.40	0/2305	0.66	0/3117
25	6	0.50	5/2843 (0.2%)	0.72	5/3845 (0.1%)
26	7	0.48	4/4992 (0.1%)	0.73	7/6754 (0.1%)
27	2	0.45	0/3611	0.76	11/4881 (0.2%)
28	5	0.47	1/502 (0.2%)	0.89	3/677 (0.4%)
29	N	0.88	0/1443	1.22	13/2226 (0.6%)
30	T	0.89	1/1449 (0.1%)	1.21	4/2233 (0.2%)
All	All	0.50	30/71982 (0.0%)	0.82	180/97723 (0.2%)



Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	M	0	5
2	A	0	8
3	B	0	6
7	H	0	1
9	J	0	1
10	K	0	1
11	L	0	1
12	Q	0	1
13	P	0	1
14	S	0	2
15	O	0	1
17	V	0	1
20	D	0	1
21	G	0	1
22	0	0	2
27	2	0	1
All	All	0	34

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1156	ASP	CG-OD2	-11.08	0.99	1.25
16	U	258	TRP	CE3-CZ3	-11.03	1.19	1.38
26	7	575	ARG	CG-CD	-10.55	1.25	1.51
26	7	677	TYR	CD1-CE1	-10.33	1.23	1.39
25	6	347	TYR	CE1-CZ	-8.88	1.27	1.38

The worst 5 of 180 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	J	24	LEU	CB-CG-CD2	-27.19	64.77	111.00
12	Q	114	MET	CG-SD-CE	-14.42	77.13	100.20
1	M	172	MET	CG-SD-CE	-12.56	80.10	100.20
18	W	227	MET	CG-SD-CE	-12.44	80.30	100.20
9	J	38	ARG	CG-CD-NE	11.35	135.63	111.80

There are no chirality outliers.

5 of 34 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	M	269	ILE	Peptide
1	M	271	GLY	Peptide
1	M	272	LYS	Peptide
1	M	30	TYR	Peptide
1	M	31	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	2175	0	2283	37	0
2	A	11167	0	11189	222	0
3	B	9227	0	9200	180	0
4	C	2086	0	2045	36	0
5	E	1752	0	1776	21	0
6	F	705	0	731	8	0
7	H	1080	0	1049	30	0
8	I	927	0	881	17	0
9	J	540	0	553	16	0
10	K	924	0	934	10	0
11	L	352	0	375	8	0
12	Q	1619	0	1452	34	0
13	P	1484	0	1480	32	0
14	S	1294	0	1289	27	0
15	O	1422	0	1500	27	0
16	U	885	0	866	27	0
17	V	815	0	822	27	0
18	W	2010	0	2026	37	0
19	X	1288	0	1307	18	0
20	D	1331	0	1345	21	0
21	G	1335	0	1346	25	0
22	0	6091	0	6155	142	0
23	1	3382	0	3436	52	0
24	4	2267	0	2323	37	0
25	6	2786	0	2804	58	0
26	7	4889	0	4876	102	0
27	2	3546	0	3593	58	0
28	5	498	0	506	12	0
29	N	1288	0	719	17	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
30	T	1291	0	713	12	0
31	4	1	0	0	0	0
31	6	4	0	0	0	0
31	A	2	0	0	0	0
31	B	1	0	0	0	0
31	C	1	0	0	0	0
31	I	2	0	0	0	0
31	J	1	0	0	0	0
31	L	1	0	0	0	0
31	M	1	0	0	0	0
31	S	1	0	0	0	0
32	7	1	0	0	0	0
32	A	1	0	0	0	0
33	0	8	0	0	2	0
All	All	70481	0	69574	1257	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 1257 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:J:24:LEU:CG	9:J:24:LEU:CD1	1.74	1.56
2:A:107:CYS:HB3	2:A:110:CYS:SG	2.01	1.00
25:6:406:CYS:HB3	25:6:440:CYS:SG	2.04	0.96
22:0:134:ARG:O	22:0:138:ASN:HB2	1.68	0.91
11:L:27:LEU:N	11:L:39:SER:HG	1.73	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	M	273/345 (79%)	238 (87%)	35 (13%)	0	100	100
2	A	1417/1733 (82%)	1249 (88%)	168 (12%)	0	100	100
3	B	1150/1224 (94%)	1000 (87%)	148 (13%)	2 (0%)	44	74
4	C	263/318 (83%)	231 (88%)	32 (12%)	0	100	100
5	E	212/215 (99%)	193 (91%)	19 (9%)	0	100	100
6	F	85/155 (55%)	78 (92%)	7 (8%)	0	100	100
7	H	129/146 (88%)	109 (84%)	20 (16%)	0	100	100
8	I	112/122 (92%)	98 (88%)	14 (12%)	0	100	100
9	J	64/70 (91%)	55 (86%)	9 (14%)	0	100	100
10	K	113/120 (94%)	107 (95%)	6 (5%)	0	100	100
11	L	42/70 (60%)	30 (71%)	12 (29%)	0	100	100
12	Q	208/735 (28%)	198 (95%)	10 (5%)	0	100	100
13	P	173/400 (43%)	161 (93%)	12 (7%)	0	100	100
14	S	162/309 (52%)	143 (88%)	18 (11%)	1 (1%)	22	55
15	O	179/240 (75%)	168 (94%)	11 (6%)	0	100	100
16	U	101/286 (35%)	96 (95%)	5 (5%)	0	100	100
17	V	100/122 (82%)	97 (97%)	3 (3%)	0	100	100
18	W	241/482 (50%)	230 (95%)	11 (5%)	0	100	100
19	X	158/328 (48%)	145 (92%)	13 (8%)	0	100	100
20	D	164/221 (74%)	159 (97%)	5 (3%)	0	100	100
21	G	169/171 (99%)	154 (91%)	14 (8%)	1 (1%)	22	55
22	0	750/778 (96%)	715 (95%)	35 (5%)	0	100	100
23	1	407/642 (63%)	394 (97%)	13 (3%)	0	100	100
24	4	286/338 (85%)	276 (96%)	10 (4%)	0	100	100
25	6	351/461 (76%)	334 (95%)	17 (5%)	0	100	100
26	7	604/843 (72%)	564 (93%)	40 (7%)	0	100	100
27	2	435/513 (85%)	417 (96%)	18 (4%)	0	100	100
28	5	64/72 (89%)	55 (86%)	9 (14%)	0	100	100
All	All	8412/11459 (73%)	7694 (92%)	714 (8%)	4 (0%)	100	100

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	364	ILE
3	B	363	HIS
21	G	57	GLN
14	S	167	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	245/299 (82%)	245 (100%)	0	100	100
2	A	1235/1520 (81%)	1232 (100%)	3 (0%)	92	94
3	B	1000/1061 (94%)	997 (100%)	3 (0%)	91	92
4	C	233/274 (85%)	231 (99%)	2 (1%)	75	82
5	E	196/197 (100%)	196 (100%)	0	100	100
6	F	77/137 (56%)	77 (100%)	0	100	100
7	H	118/128 (92%)	116 (98%)	2 (2%)	56	72
8	I	108/116 (93%)	106 (98%)	2 (2%)	52	69
9	J	61/65 (94%)	61 (100%)	0	100	100
10	K	99/102 (97%)	99 (100%)	0	100	100
11	L	39/57 (68%)	39 (100%)	0	100	100
12	Q	147/641 (23%)	144 (98%)	3 (2%)	50	68
13	P	166/363 (46%)	165 (99%)	1 (1%)	84	88
14	S	141/274 (52%)	140 (99%)	1 (1%)	81	86
15	O	153/205 (75%)	153 (100%)	0	100	100
16	U	99/260 (38%)	96 (97%)	3 (3%)	36	58
17	V	94/108 (87%)	93 (99%)	1 (1%)	70	79
18	W	224/429 (52%)	223 (100%)	1 (0%)	89	91
19	X	144/295 (49%)	142 (99%)	2 (1%)	62	75
20	D	146/200 (73%)	146 (100%)	0	100	100
21	G	151/152 (99%)	151 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
22	0	684/707 (97%)	684 (100%)	0	100	100
23	1	389/589 (66%)	387 (100%)	2 (0%)	86	90
24	4	259/300 (86%)	259 (100%)	0	100	100
25	6	322/418 (77%)	322 (100%)	0	100	100
26	7	540/737 (73%)	537 (99%)	3 (1%)	84	88
27	2	394/468 (84%)	392 (100%)	2 (0%)	86	90
28	5	53/66 (80%)	53 (100%)	0	100	100
All	All	7517/10168 (74%)	7486 (100%)	31 (0%)	88	91

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
12	Q	333	LYS
26	7	348	ARG
16	U	44	LYS
27	2	419	LYS
23	1	306	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
23	1	610	ASN
26	7	740	HIS
23	1	639	ASN
26	7	584	ASN
27	2	500	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 17 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
33	SF4	0	801	-	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
33	SF4	0	801	-	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
33	0	801	SF4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.




## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-42437. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y



Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)

### 6.3.1 Primary map



X Index: 216



Y Index: 232




Z Index: 210


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1031 nm<sup>3</sup>; this corresponds to an approximate mass of 932 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)



\*Reported resolution corresponds to spatial frequency of 0.263 Å<sup>-1</sup>


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-42437 and PDB model 8UOQ. Per-residue inclusion information can be found in section [3](#) on page [11](#).


### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8280	 0.2390
0	 0.6950	 0.1780
1	 0.6570	 0.1870
2	 0.7450	 0.1480
4	 0.7750	 0.1900
5	 0.6180	 0.1540
6	 0.7370	 0.1850
7	 0.7490	 0.1710
A	 0.9580	 0.3280
B	 0.9520	 0.3380
C	 0.9020	 0.2690
D	 0.3700	 0.1540
E	 0.9640	 0.3190
F	 0.9300	 0.3100
G	 0.6530	 0.1970
H	 0.9270	 0.3000
I	 0.9470	 0.3070
J	 0.9010	 0.2700
K	 0.8330	 0.2730
L	 0.7940	 0.1800
M	 0.7590	 0.2010
N	 0.9490	 0.2210
O	 0.9000	 0.1860
P	 0.9170	 0.2080
Q	 0.8310	 0.2520
S	 0.9020	 0.1940
T	 0.9480	 0.2290
U	 0.5180	 0.1400
V	 0.6040	 0.1810
W	 0.8160	 0.1690
X	 0.8750	 0.1570

