



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:12 AM UTC

PDB ID : 3UPC / pdb\_00003upc  
Title : A general strategy for the generation of human antibody variable domains with increased aggregation resistance  
Authors : Dudgeon, K.; Rouet, R.; Kokmeijer, I.; Langley, D.B.; Christ, D.  
Deposited on : 2011-11-17  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	113	<div><div>%</div><div>88%9%</div><div>.</div></div>
1	B	113	<div><div>%</div><div>94%</div><div>..</div></div>
1	C	113	<div><div>90%7%</div><div>..</div></div>
1	D	113	<div><div>%</div><div>93%5%</div><div>.</div></div>
1	E	113	<div><div>14%</div><div>93%</div><div>...</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	113	<div><div></div><div>4%</div><div>85%</div><div>12%</div><div>..</div></div>
1	G	113	<div><div></div><div>2%</div><div>94%</div><div></div><div>..</div></div>
1	H	113	<div><div></div><div>%</div><div>90%</div><div>7%</div><div>..</div></div>
1	I	113	<div><div></div><div>7%</div><div>91%</div><div>6%</div><div>..</div></div>
1	J	113	<div><div></div><div>3%</div><div>93%</div><div>5%</div><div>..</div></div>

## 2 Entry composition


There are 5 unique types of molecules in this entry. The entry contains 8217 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called heavy chain variable domain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	113	Total	C	N	O	S	0	0	0
			822	516	138	164	4			
1	B	113	Total	C	N	O	S	0	0	0
			826	520	140	162	4			
1	C	113	Total	C	N	O	S	0	0	0
			827	520	140	163	4			
1	D	113	Total	C	N	O	S	0	0	0
			825	518	139	164	4			
1	E	113	Total	C	N	O	S	0	0	0
			773	487	135	147	4			
1	F	113	Total	C	N	O	S	0	0	0
			823	518	139	162	4			
1	G	113	Total	C	N	O	S	0	0	0
			820	518	139	159	4			
1	H	113	Total	C	N	O	S	0	0	0
			817	514	136	163	4			
1	I	113	Total	C	N	O	S	0	0	0
			805	508	136	157	4			
1	J	113	Total	C	N	O	S	0	0	0
			791	498	133	156	4			

- Molecule 2 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: C<sub>10</sub>H<sub>22</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			16	10	6		
2	D	1	Total	C	O	0	0
			16	10	6		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula:  $C_6H_{14}O_4$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula:  $C_8H_{18}O_5$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	G	1	Total	C	O	0	0
			13	8	5		
5	H	1	Total	C	O	0	0
			13	8	5		
5	I	1	Total	C	O	0	0
			13	8	5		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: heavy chain variable domain





- Molecule 1: heavy chain variable domain



- Molecule 1: heavy chain variable domain





- Molecule 1: heavy chain variable domain




- Molecule 1: heavy chain variable domain





- Molecule 1: heavy chain variable domain





- Molecule 1: heavy chain variable domain




- Molecule 1: heavy chain variable domain



- Molecule 1: heavy chain variable domain



- Molecule 1: heavy chain variable domain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.61Å 143.12Å 145.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	145.61 – 2.80 145.61 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (145.61-2.80) 99.9 (145.61-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.15	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.61 (at 2.81Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.209 , 0.247 0.210 , 0.243	Depositor DCC
$R_{free}$ test set	2180 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.6	Xtriage
Anisotropy	0.597	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 41.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.023 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8217	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.72 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7543e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, PEG, 1PE, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	0/840	1.01	3/1142 (0.3%)
1	B	0.82	0/844	0.84	1/1145 (0.1%)
1	C	0.78	0/845	0.95	3/1146 (0.3%)
1	D	0.77	0/843	0.91	3/1145 (0.3%)
1	E	0.74	0/790	0.78	1/1077 (0.1%)
1	F	0.78	0/841	0.87	2/1141 (0.2%)
1	G	0.76	0/838	0.86	1/1139 (0.1%)
1	H	0.83	0/835	0.82	0/1135
1	I	0.78	0/823	0.80	0/1120
1	J	0.71	0/808	1.00	3/1101 (0.3%)
All	All	0.78	0/8307	0.89	17/11291 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	1
1	G	0	1
1	H	0	1
1	I	0	1
1	J	0	1
All	All	0	10

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	103	PHE	N-CA-C	18.50	135.87	110.35
1	A	103	PHE	N-CA-C	14.89	132.94	109.86
1	C	103	PHE	N-CA-C	14.71	133.03	109.94
1	D	103	PHE	N-CA-C	13.87	131.12	110.17
1	F	103	PHE	N-CA-C	11.86	123.87	108.34
1	A	102	ALA	N-CA-C	9.27	125.16	111.34
1	G	103	PHE	N-CA-C	6.30	124.22	110.80
1	D	102	ALA	N-CA-C	6.24	124.08	110.80
1	J	103	PHE	N-CA-CB	-6.07	102.01	110.38
1	C	102	ALA	N-CA-C	5.88	123.32	110.80
1	F	103	PHE	N-CA-CB	-5.59	100.81	110.87
1	C	103	PHE	N-CA-CB	-5.49	100.14	110.07
1	J	102	ALA	N-CA-C	5.22	121.92	110.80
1	D	103	PHE	N-CA-CB	-5.18	102.18	110.51
1	B	103	PHE	N-CA-C	5.16	121.80	110.80
1	E	103	PHE	N-CA-C	5.01	121.47	110.80
1	A	105	TYR	N-CA-C	5.00	116.73	111.28

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	102	ALA	Peptide
1	B	102	ALA	Peptide
1	C	102	ALA	Peptide
1	D	102	ALA	Peptide
1	E	102	ALA	Peptide
1	F	102	ALA	Peptide
1	G	102	ALA	Peptide
1	H	102	ALA	Peptide
1	I	102	ALA	Peptide
1	J	102	ALA	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	822	0	731	8	0
1	B	826	0	749	10	0
1	C	827	0	752	7	0
1	D	825	0	740	7	0
1	E	773	0	646	10	0
1	F	823	0	746	18	0
1	G	820	0	740	8	0
1	H	817	0	730	13	0
1	I	805	0	703	16	0
1	J	791	0	674	7	0
2	A	16	0	22	0	0
2	D	16	0	22	0	0
3	B	10	0	14	0	0
4	C	7	0	10	0	0
5	G	13	0	18	0	0
5	H	13	0	18	0	0
5	I	13	0	18	0	0
All	All	8217	0	7333	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:102:ALA:HB1	1:I:103:PHE:HB3	1.43	1.00
1:H:102:ALA:HB1	1:H:103:PHE:HB3	1.50	0.93
1:E:83:MET:HE1	1:I:112:VAL:HG21	1.56	0.87
1:G:102:ALA:HB3	1:G:103:PHE:HD2	1.39	0.86
1:G:102:ALA:HB3	1:G:103:PHE:CD2	2.12	0.84
1:I:102:ALA:CB	1:I:103:PHE:HB3	2.10	0.79
1:E:102:ALA:HB3	1:E:103:PHE:HD2	1.48	0.79
1:A:83:MET:HE1	1:B:112:VAL:HG21	1.66	0.76
1:I:102:ALA:CB	1:I:103:PHE:HD1	1.97	0.76
1:B:102:ALA:HB3	1:B:103:PHE:HD1	1.52	0.73
1:H:102:ALA:CB	1:H:103:PHE:HB3	2.20	0.71
1:E:12:VAL:HG11	1:E:86:LEU:HD22	1.71	0.71
1:F:83:MET:HE1	1:J:112:VAL:HG21	1.72	0.71
1:G:83:MET:HE1	1:H:112:VAL:HG21	1.74	0.68
1:C:83:MET:HE1	1:D:112:VAL:HG21	1.76	0.67
1:F:102:ALA:CB	1:J:102:ALA:HB2	2.25	0.66
1:I:102:ALA:CB	1:I:103:PHE:CD1	2.77	0.66

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:102:ALA:HB1	1:I:103:PHE:CB	2.24	0.64
1:A:112:VAL:HG21	1:B:83:MET:HE1	1.80	0.63
1:H:102:ALA:HB1	1:H:103:PHE:CB	2.26	0.63
1:E:102:ALA:HB3	1:E:103:PHE:CD2	2.34	0.63
1:H:102:ALA:HB1	1:H:103:PHE:HD2	1.65	0.62
1:H:20:LEU:HG	1:H:83:MET:HE2	1.82	0.61
1:H:102:ALA:HB1	1:H:103:PHE:CD2	2.35	0.61
1:F:103:PHE:H	1:F:103:PHE:HD2	1.47	0.60
1:B:102:ALA:HB3	1:B:103:PHE:CD1	2.36	0.59
1:I:102:ALA:CB	1:I:103:PHE:CB	2.79	0.59
1:I:102:ALA:HB1	1:I:103:PHE:CD1	2.38	0.58
1:E:112:VAL:HG21	1:I:83:MET:HE1	1.84	0.58
1:C:16:GLY:O	1:C:86:LEU:HD12	2.05	0.56
1:F:12:VAL:HG21	1:F:86:LEU:HD12	1.87	0.56
1:A:20:LEU:HG	1:A:83:MET:HE2	1.87	0.56
1:F:114:VAL:HG11	1:J:87:ARG:C	2.31	0.55
1:F:87:ARG:HG3	1:F:89:GLU:HG2	1.89	0.55
1:E:102:ALA:CB	1:E:103:PHE:HD2	2.17	0.55
1:E:102:ALA:CB	1:E:103:PHE:CD2	2.90	0.55
1:F:102:ALA:HB3	1:J:102:ALA:HB2	1.88	0.54
1:F:12:VAL:HG11	1:F:86:LEU:HD13	1.89	0.54
1:C:112:VAL:HG21	1:D:83:MET:HE1	1.90	0.53
1:B:20:LEU:HG	1:B:83:MET:HE2	1.91	0.53
1:I:20:LEU:HG	1:I:83:MET:HE2	1.91	0.53
1:C:20:LEU:HG	1:C:83:MET:HE2	1.90	0.52
1:E:83:MET:HB3	1:E:86:LEU:HD11	1.91	0.52
1:G:102:ALA:CB	1:G:103:PHE:CD2	2.90	0.52
1:B:102:ALA:CB	1:B:103:PHE:HB3	2.40	0.51
1:F:20:LEU:HG	1:F:83:MET:HE2	1.92	0.51
1:B:102:ALA:CB	1:B:103:PHE:HD1	2.20	0.51
1:F:83:MET:HB3	1:F:86:LEU:HD21	1.93	0.51
1:B:102:ALA:HB1	1:B:103:PHE:HB3	1.93	0.51
1:H:102:ALA:CB	1:H:103:PHE:CB	2.87	0.50
1:H:60:TYR:HB2	1:H:65:LYS:HG2	1.92	0.50
1:D:89:GLU:H	1:D:89:GLU:CD	2.20	0.50
1:F:103:PHE:N	1:F:103:PHE:CD2	2.78	0.49
1:A:67:ARG:NH1	1:A:87:ARG:HD2	2.29	0.48
1:F:6:GLU:OE1	1:J:108:GLN:NE2	2.38	0.48
1:F:65:LYS:NZ	1:H:105:TYR:O	2.47	0.48
1:I:102:ALA:HB3	1:I:103:PHE:CD1	2.48	0.48
1:A:18:LEU:HB3	1:A:83:MET:HE3	1.97	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:18:LEU:HB3	1:C:83:MET:HE3	1.96	0.46
1:F:18:LEU:HB3	1:F:83:MET:HE3	1.98	0.46
1:F:111:LEU:HD12	1:F:112:VAL:N	2.31	0.46
1:C:102:ALA:HB2	1:D:102:ALA:HB2	1.97	0.45
1:B:95:TYR:OH	1:I:65:LYS:HE2	2.17	0.45
1:G:20:LEU:HG	1:G:83:MET:HE2	2.00	0.44
1:H:102:ALA:CB	1:H:103:PHE:HD2	2.30	0.44
1:B:102:ALA:CB	1:B:103:PHE:CD1	3.00	0.44
1:E:83:MET:CB	1:E:86:LEU:HD11	2.48	0.44
1:G:112:VAL:HG21	1:H:83:MET:HE1	2.00	0.44
1:A:105:TYR:O	1:I:65:LYS:HE3	2.18	0.43
1:A:83:MET:HB3	1:A:86:LEU:HD21	2.00	0.43
1:F:102:ALA:HB2	1:J:102:ALA:HB2	1.97	0.43
1:G:18:LEU:HB3	1:G:83:MET:HE3	2.01	0.43
1:A:39:GLN:HB2	1:A:45:LEU:HD23	2.00	0.43
1:I:102:ALA:HB3	1:I:103:PHE:HD1	1.80	0.42
1:E:39:GLN:HB2	1:E:45:LEU:HD23	2.01	0.42
1:D:103:PHE:HB2	1:D:108:GLN:OE1	2.19	0.42
1:D:45:LEU:HD23	1:D:45:LEU:HA	1.92	0.42
1:D:20:LEU:HG	1:D:83:MET:HE2	2.01	0.42
1:F:45:LEU:HD23	1:F:45:LEU:HA	1.92	0.42
1:G:73:ASP:OD2	1:G:73:ASP:C	2.62	0.41
1:I:36:TRP:HD1	1:I:70:ILE:HD12	1.84	0.41
1:F:18:LEU:HD23	1:F:83:MET:CE	2.51	0.41
1:J:20:LEU:HG	1:J:83:MET:HE2	2.02	0.41
1:H:45:LEU:HD23	1:H:45:LEU:HA	1.94	0.41
1:C:63:SER:HB2	1:C:89:GLU:OE2	2.21	0.41
1:I:18:LEU:HD23	1:I:83:MET:CE	2.51	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	111/113 (98%)	107 (96%)	4 (4%)	0	100	100
1	B	111/113 (98%)	108 (97%)	2 (2%)	1 (1%)	14	41
1	C	111/113 (98%)	108 (97%)	3 (3%)	0	100	100
1	D	111/113 (98%)	109 (98%)	2 (2%)	0	100	100
1	E	111/113 (98%)	108 (97%)	2 (2%)	1 (1%)	14	41
1	F	111/113 (98%)	108 (97%)	2 (2%)	1 (1%)	14	41
1	G	111/113 (98%)	108 (97%)	2 (2%)	1 (1%)	14	41
1	H	111/113 (98%)	107 (96%)	2 (2%)	2 (2%)	6	23
1	I	111/113 (98%)	106 (96%)	3 (3%)	2 (2%)	6	23
1	J	111/113 (98%)	108 (97%)	3 (3%)	0	100	100
All	All	1110/1130 (98%)	1077 (97%)	25 (2%)	8 (1%)	18	47

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	103	PHE
1	E	103	PHE
1	G	103	PHE
1	H	103	PHE
1	I	103	PHE
1	F	102	ALA
1	H	102	ALA
1	I	102	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/89 (86%)	75 (97%)	2 (3%)	40	75
1	B	78/89 (88%)	76 (97%)	2 (3%)	40	75
1	C	79/89 (89%)	76 (96%)	3 (4%)	29	64
1	D	78/89 (88%)	77 (99%)	1 (1%)	61	86

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	60/89 (67%)	58 (97%)	2 (3%)	33	69
1	F	78/89 (88%)	74 (95%)	4 (5%)	21	54
1	G	76/89 (85%)	75 (99%)	1 (1%)	61	86
1	H	77/89 (86%)	74 (96%)	3 (4%)	28	64
1	I	71/89 (80%)	69 (97%)	2 (3%)	38	73
1	J	67/89 (75%)	65 (97%)	2 (3%)	36	72
All	All	741/890 (83%)	719 (97%)	22 (3%)	36	72

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	GLU
1	A	103	PHE
1	B	49	SER
1	B	103	PHE
1	C	65	LYS
1	C	89	GLU
1	C	103	PHE
1	D	103	PHE
1	E	86	LEU
1	E	103	PHE
1	F	12	VAL
1	F	21	SER
1	F	33	ASP
1	F	103	PHE
1	G	103	PHE
1	H	25	SER
1	H	28	THR
1	H	65	LYS
1	I	33	ASP
1	I	65	LYS
1	J	103	PHE
1	J	110	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	77	ASN
1	D	77	ASN

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	E	82	GLN
1	G	39	GLN
1	H	82	GLN
1	I	77	ASN
1	I	82	GLN
1	J	77	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	1PE	D	116	-	15,15,15	0.86	0	14,14,14	0.61	0
3	PGE	B	116	-	9,9,9	0.83	0	8,8,8	0.84	0
2	1PE	A	2	-	15,15,15	0.83	0	14,14,14	0.92	0
5	PG4	G	116	-	12,12,12	0.72	0	11,11,11	0.63	0
4	PEG	C	116	-	6,6,6	0.69	0	5,5,5	0.28	0
5	PG4	H	1	-	12,12,12	0.76	0	11,11,11	0.77	0
5	PG4	I	116	-	12,12,12	0.72	0	11,11,11	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PE	D	116	-	-	8/13/13/13	-
3	PGE	B	116	-	-	3/7/7/7	-
2	1PE	A	2	-	-	7/13/13/13	-
5	PG4	G	116	-	-	3/10/10/10	-
4	PEG	C	116	-	-	2/4/4/4	-
5	PG4	H	1	-	-	4/10/10/10	-
5	PG4	I	116	-	-	4/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	116	PG4	C8-C7-O4-C6
2	A	2	1PE	C15-C25-OH5-C14
2	D	116	1PE	C15-C25-OH5-C14
2	D	116	1PE	OH4-C13-C23-OH3
2	A	2	1PE	OH5-C14-C24-OH4
2	A	2	1PE	OH6-C15-C25-OH5
5	G	116	PG4	O1-C1-C2-O2
3	B	116	PGE	O2-C3-C4-O3
5	G	116	PG4	O3-C5-C6-O4
3	B	116	PGE	O1-C1-C2-O2
4	C	116	PEG	O2-C3-C4-O4
5	I	116	PG4	O3-C5-C6-O4
5	H	1	PG4	O4-C7-C8-O5
4	C	116	PEG	O1-C1-C2-O2
5	I	116	PG4	O2-C3-C4-O3
2	D	116	1PE	C12-C22-OH3-C23
2	D	116	1PE	C14-C24-OH4-C13
2	D	116	1PE	C23-C13-OH4-C24
2	A	2	1PE	C13-C23-OH3-C22
5	I	116	PG4	C4-C3-O2-C2
5	H	1	PG4	C4-C3-O2-C2
2	A	2	1PE	OH2-C12-C22-OH3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	116	PGE	C6-C5-O3-C4
5	H	1	PG4	C1-C2-O2-C3
2	D	116	1PE	OH5-C14-C24-OH4
2	D	116	1PE	OH6-C15-C25-OH5
5	I	116	PG4	C8-C7-O4-C6
2	A	2	1PE	C14-C24-OH4-C13
5	H	1	PG4	O3-C5-C6-O4
2	A	2	1PE	OH4-C13-C23-OH3
2	D	116	1PE	OH7-C16-C26-OH6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	113/113 (100%)	-0.23	1 (0%) 81 74	20, 35, 48, 61	0
1	B	113/113 (100%)	-0.27	1 (0%) 81 74	23, 34, 48, 70	0
1	C	113/113 (100%)	-0.35	0 100 100	15, 31, 51, 61	0
1	D	113/113 (100%)	-0.37	1 (0%) 81 74	21, 34, 53, 60	0
1	E	113/113 (100%)	0.77	16 (14%) 6 5	32, 59, 100, 127	0
1	F	113/113 (100%)	-0.22	5 (4%) 39 31	21, 36, 51, 80	0
1	G	113/113 (100%)	-0.16	2 (1%) 67 58	25, 38, 54, 69	0
1	H	113/113 (100%)	-0.24	1 (0%) 81 74	22, 32, 55, 78	0
1	I	113/113 (100%)	0.31	8 (7%) 22 16	28, 50, 70, 93	0
1	J	113/113 (100%)	0.02	3 (2%) 56 46	27, 50, 69, 80	0
All	All	1130/1130 (100%)	-0.07	38 (3%) 48 39	15, 39, 69, 127	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	77	ASN	5.1
1	E	75	SER	5.1
1	E	115	SER	5.0
1	I	115	SER	4.1
1	I	43	LYS	4.1
1	E	73	ASP	4.0
1	E	30	SER	3.8
1	I	3	GLN	3.7
1	E	7	SER	3.7
1	D	115	SER	3.6
1	I	102	ALA	3.5
1	E	3	GLN	3.4
1	J	115	SER	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	I	42	GLY	3.0
1	E	31	ASP	2.9
1	G	115	SER	2.8
1	E	100	TYR	2.8
1	F	115	SER	2.8
1	E	70	ILE	2.7
1	F	3	GLN	2.6
1	J	57	SER	2.6
1	F	42	GLY	2.5
1	H	115	SER	2.4
1	F	44	GLY	2.3
1	E	57	SER	2.3
1	I	26	GLY	2.3
1	F	43	LYS	2.2
1	E	5	LEU	2.2
1	I	41	PRO	2.2
1	E	33	ASP	2.2
1	A	115	SER	2.1
1	E	102	ALA	2.1
1	G	31	ASP	2.1
1	B	30	SER	2.1
1	E	54	SER	2.1
1	I	107	GLY	2.1
1	J	28	THR	2.1
1	E	76	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PEG	C	116	7/7	0.77	0.26	36,59,64,64	0
5	PG4	I	116	13/13	0.82	0.21	45,58,70,71	0
3	PGE	B	116	10/10	0.84	0.20	29,42,52,57	0
5	PG4	H	1	13/13	0.86	0.16	41,51,55,58	0
2	1PE	A	2	16/16	0.86	0.18	27,49,61,68	0
5	PG4	G	116	13/13	0.87	0.20	45,57,97,101	0
2	1PE	D	116	16/16	0.90	0.16	40,54,68,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.