



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 11:40 pm BST

PDB ID : 4UP2
Title : Crystal structure of Escherichia coli tryptophanase purified from alkaline stressed bacterial culture.
Authors : Rety, S.; Deschamps, P.; Leulliot, N.
Deposited on : 2014-06-11
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	
1	B	471	
1	C	471	
1	D	471	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	1473	-	-	-	X
2	SO4	B	1459	-	-	-	X
2	SO4	D	1459	-	-	X	-
2	SO4	D	1460	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 25834 atoms, of which 12736 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRYPTOPHANASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	420	Total	C	H	N	O	S	0	0	0
			6612	2119	3297	558	618	20			
1	B	408	Total	C	H	N	O	S	0	0	0
			6381	2047	3171	539	605	19			
1	C	404	Total	C	H	N	O	S	0	0	0
			6336	2034	3155	534	595	18			
1	D	397	Total	C	H	N	O	S	0	0	0
			6242	2000	3107	528	589	18			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	D	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	D	1	5	4	1	0	0
2	D	1	5	4	1	0	0
2	D	1	5	4	1	0	0
2	D	1	5	4	1	0	0
2	D	1	5	4	1	0	0

- Molecule 3 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	S		
3	B	1	10	2	6	1	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	39	39	39	0	0
4	B	32	32	32	0	0
4	C	25	25	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	17	Total	O	0	0
			17	17		



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	158.24Å 158.24Å 387.77Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.01 – 2.78 47.01 – 2.78	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.01-2.78) 99.9 (47.01-2.78)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 2.77Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.215 , 0.246 0.218 , 0.247	Depositor DCC
R_{free} test set	3672 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	56.3	Xtrriage
Anisotropy	0.029	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	25834	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/3382	0.41	0/4562
1	B	0.22	0/3273	0.40	0/4415
1	C	0.22	0/3245	0.39	0/4377
1	D	0.22	0/3196	0.39	0/4311
All	All	0.22	0/13096	0.40	0/17665

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	462	ARG	Peptide
1	A	463	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3315	3297	3297	5	0
1	B	3210	3171	3171	14	0
1	C	3181	3155	3155	9	0
1	D	3135	3107	3107	13	0
2	A	30	0	0	0	0
2	B	40	0	0	1	0
2	C	40	0	0	0	0
2	D	30	0	0	3	0
3	B	4	6	6	0	0
4	A	39	0	0	0	0
4	B	32	0	0	0	0
4	C	25	0	0	0	0
4	D	17	0	0	0	0
All	All	13098	12736	12736	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:310:LEU:N	2:D:1459:SO4:S	2.65	0.70
1:A:177:GLU:OE2	1:A:219:LYS:NZ	2.27	0.68
1:A:466:ALA:HB2	1:B:84:ALA:HB1	1.81	0.63
1:D:118:GLU:OE1	1:D:125:ARG:NH2	2.31	0.63
1:D:310:LEU:N	2:D:1459:SO4:O1	2.34	0.61
1:B:228:SER:HG	1:B:258:TYR:HH	1.51	0.56
1:B:117:ARG:NH2	1:B:221:ASP:O	2.39	0.55
1:B:22:THR:O	1:B:27:ARG:NH1	2.40	0.55
1:C:27:ARG:NH2	1:C:388:VAL:O	2.42	0.53
1:B:2:GLU:OE2	1:D:441:HIS:ND1	2.42	0.53
1:C:195:ILE:HD13	1:C:226:MET:SD	2.51	0.51
1:B:279:LEU:HD21	1:B:318:LEU:HD21	1.93	0.49
1:D:451:GLY:C	1:D:452:LEU:HD12	2.33	0.49
1:A:228:SER:HB2	1:A:231:PHE:HB3	1.94	0.49
1:C:379:GLN:NE2	1:C:395:GLU:OE2	2.46	0.49
1:D:264:LEU:HD23	1:D:264:LEU:C	2.33	0.48
1:C:264:LEU:HD23	1:C:264:LEU:C	2.32	0.48
1:C:196:THR:HG21	1:C:230:ARG:HD2	1.95	0.48
1:B:264:LEU:C	1:B:264:LEU:HD23	2.34	0.48
1:D:110:ILE:HB	1:D:111:PRO:HD3	1.95	0.47
1:B:196:THR:HG21	1:B:230:ARG:HD2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:133:ASN:N	1:D:133:ASN:OD1	2.48	0.46
1:C:228:SER:HB2	1:C:231:PHE:HB3	1.97	0.46
1:B:24:ARG:N	2:B:1460:SO4:O2	2.49	0.45
1:D:22:THR:O	1:D:27:ARG:NH1	2.48	0.45
1:A:462:ARG:O	1:A:463:HIS:HB2	2.16	0.44
1:A:14:ARG:HA	1:A:14:ARG:NE	2.34	0.43
1:B:288:PHE:HA	1:B:289:ASP:C	2.40	0.43
1:D:394:VAL:HG22	1:D:395:GLU:N	2.34	0.42
1:D:310:LEU:N	2:D:1459:SO4:O4	2.52	0.42
1:B:133:ASN:OD1	1:B:133:ASN:N	2.50	0.42
1:D:177:GLU:OE2	1:D:219:LYS:NZ	2.53	0.42
1:C:195:ILE:CG2	1:C:231:PHE:HA	2.50	0.42
1:D:228:SER:HB2	1:D:231:PHE:HB3	2.02	0.41
1:B:197:SER:O	1:B:202:GLY:HA2	2.19	0.41
1:B:258:TYR:CE2	1:B:264:LEU:HD12	2.55	0.41
1:C:396:ILE:O	1:C:396:ILE:HG22	2.21	0.41
1:B:228:SER:HB2	1:B:231:PHE:HB3	2.02	0.41
1:C:118:GLU:HA	1:C:123:LEU:HB3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	412/471 (88%)	399 (97%)	11 (3%)	2 (0%)	29	58
1	B	400/471 (85%)	387 (97%)	13 (3%)	0	100	100
1	C	396/471 (84%)	383 (97%)	12 (3%)	1 (0%)	41	70
1	D	389/471 (83%)	376 (97%)	13 (3%)	0	100	100
All	All	1597/1884 (85%)	1545 (97%)	49 (3%)	3 (0%)	47	76

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	464	PHE
1	A	463	HIS
1	C	396	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/391 (88%)	342 (99%)	4 (1%)	71	90
1	B	334/391 (85%)	331 (99%)	3 (1%)	78	92
1	C	331/391 (85%)	328 (99%)	3 (1%)	78	92
1	D	327/391 (84%)	325 (99%)	2 (1%)	86	95
All	All	1338/1564 (86%)	1326 (99%)	12 (1%)	78	92

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207	LEU
1	A	351	VAL
1	A	463	HIS
1	A	468	LEU
1	B	164	ARG
1	B	267	SER
1	B	287	PHE
1	C	96	THR
1	C	195	ILE
1	C	207	LEU
1	D	224	VAL
1	D	452	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

29 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	C	1461	-	4,4,4	0.13	0	6,6,6	0.06	0
2	SO4	B	1456	-	4,4,4	0.13	0	6,6,6	0.04	0
2	SO4	D	1459	-	4,4,4	0.14	0	6,6,6	0.04	0
2	SO4	C	1457	-	4,4,4	0.15	0	6,6,6	0.05	0
2	SO4	A	1469	-	4,4,4	0.14	0	6,6,6	0.06	0
2	SO4	A	1472	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	D	1457	-	4,4,4	0.13	0	6,6,6	0.07	0
2	SO4	B	1460	-	4,4,4	0.14	0	6,6,6	0.06	0
2	SO4	C	1460	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	C	1456	-	4,4,4	0.14	0	6,6,6	0.08	0
2	SO4	D	1455	-	4,4,4	0.15	0	6,6,6	0.04	0
2	SO4	B	1462	-	4,4,4	0.14	0	6,6,6	0.06	0
2	SO4	C	1458	-	4,4,4	0.14	0	6,6,6	0.04	0
2	SO4	A	1473	-	4,4,4	0.13	0	6,6,6	0.06	0
2	SO4	C	1462	-	4,4,4	0.27	0	6,6,6	0.28	0
2	SO4	A	1474	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	D	1456	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	D	1460	-	4,4,4	0.14	0	6,6,6	0.06	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BME	B	1461	-	3,3,3	0.30	0	1,2,2	0.12	0
2	SO4	B	1455	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	A	1471	-	4,4,4	0.13	0	6,6,6	0.06	0
2	SO4	B	1454	-	4,4,4	0.13	0	6,6,6	0.07	0
2	SO4	B	1457	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	C	1455	-	4,4,4	0.14	0	6,6,6	0.04	0
2	SO4	B	1459	-	4,4,4	0.14	0	6,6,6	0.07	0
2	SO4	C	1459	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	A	1470	-	4,4,4	0.15	0	6,6,6	0.04	0
2	SO4	D	1458	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	B	1458	-	4,4,4	0.14	0	6,6,6	0.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BME	B	1461	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1461	BME	O1-C1-C2-S2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1459	SO4	3	0
2	B	1460	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	420/471 (89%)	0.30	15 (3%) 42 37	27, 49, 85, 114	0
1	B	408/471 (86%)	0.32	14 (3%) 45 39	27, 52, 88, 114	0
1	C	404/471 (85%)	0.33	5 (1%) 79 76	28, 53, 90, 111	0
1	D	397/471 (84%)	0.76	51 (12%) 3 2	34, 73, 118, 128	0
All	All	1629/1884 (86%)	0.43	85 (5%) 27 22	27, 55, 100, 128	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	131	PHE	6.3
1	A	287	PHE	5.2
1	B	289	ASP	5.2
1	D	188	VAL	4.7
1	D	452	LEU	4.7
1	C	452	LEU	4.6
1	D	128	MET	4.5
1	D	129	VAL	4.5
1	D	185	PRO	4.4
1	A	1	MET	4.4
1	D	127	LYS	3.9
1	A	288	PHE	3.9
1	D	120	GLU	3.7
1	D	373	ALA	3.7
1	D	162	GLY	3.6
1	B	449	ILE	3.5
1	D	183	VAL	3.5
1	D	151	ARG	3.4
1	D	130	ALA	3.4
1	A	397	GLY	3.4
1	A	3	ASN	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	218	LYS	3.3
1	D	453	THR	3.2
1	D	376	PHE	3.2
1	D	417	LEU	3.2
1	B	396	ILE	3.1
1	D	368	LEU	3.0
1	A	461	LEU	3.0
1	D	418	LEU	3.0
1	C	396	ILE	2.9
1	D	226	MET	2.9
1	D	351	VAL	2.8
1	D	284	ASP	2.8
1	D	377	PRO	2.8
1	B	2	GLU	2.8
1	D	126	SER	2.8
1	D	176	LEU	2.7
1	D	367	LEU	2.7
1	A	180	ILE	2.7
1	D	125	ARG	2.7
1	B	399	PHE	2.7
1	C	32	ILE	2.6
1	A	4	PHE	2.6
1	D	190	TYR	2.6
1	B	103	ARG	2.6
1	D	180	ILE	2.6
1	D	189	PRO	2.6
1	D	113	LEU	2.6
1	D	114	ILE	2.6
1	D	395	GLU	2.5
1	B	308	GLY	2.5
1	B	452	LEU	2.5
1	D	121	LYS	2.5
1	B	397	GLY	2.5
1	A	462	ARG	2.5
1	D	3	ASN	2.5
1	A	2	GLU	2.5
1	D	150	VAL	2.4
1	D	454	PHE	2.4
1	A	396	ILE	2.4
1	D	187	ASN	2.4
1	B	288	PHE	2.4
1	A	153	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	148	CYS	2.3
1	D	365	GLY	2.2
1	D	369	PRO	2.2
1	D	164	ARG	2.2
1	B	447	SER	2.2
1	A	454	PHE	2.2
1	D	220	TYR	2.2
1	A	181	GLU	2.2
1	D	116	LYS	2.1
1	D	119	GLN	2.1
1	D	135	PHE	2.1
1	B	450	LYS	2.1
1	D	416	GLU	2.1
1	C	453	THR	2.1
1	D	352	CYS	2.1
1	B	40	LEU	2.1
1	C	195	ILE	2.1
1	D	224	VAL	2.1
1	B	398	SER	2.1
1	D	394	VAL	2.0
1	D	57	GLY	2.0
1	D	108	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	B	1459	5/5	0.60	0.51	123,132,156,271	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	D	1460	5/5	0.64	0.43	176,177,241,283	0
2	SO4	B	1457	5/5	0.65	0.23	102,113,117,137	0
2	SO4	D	1459	5/5	0.69	0.27	93,111,116,141	0
2	SO4	B	1456	5/5	0.71	0.23	113,113,132,141	0
2	SO4	A	1472	5/5	0.73	0.29	67,83,120,123	0
2	SO4	B	1460	5/5	0.74	0.20	120,121,184,253	0
2	SO4	A	1473	5/5	0.75	0.54	90,92,118,129	0
2	SO4	C	1457	5/5	0.78	0.27	84,95,120,123	0
2	SO4	B	1455	5/5	0.80	0.28	61,81,97,118	0
2	SO4	C	1462	5/5	0.83	0.27	69,74,81,151	0
2	SO4	C	1459	5/5	0.83	0.34	94,95,119,127	0
2	SO4	C	1460	5/5	0.86	0.23	121,124,129,132	0
2	SO4	C	1461	5/5	0.86	0.50	84,88,107,125	0
2	SO4	C	1458	5/5	0.87	0.27	83,87,110,111	0
2	SO4	A	1474	5/5	0.88	0.20	91,99,111,123	0
2	SO4	D	1457	5/5	0.88	0.27	69,81,110,112	0
2	SO4	D	1458	5/5	0.90	0.20	80,92,94,99	0
2	SO4	B	1462	5/5	0.91	0.32	120,137,146,194	0
2	SO4	D	1455	5/5	0.92	0.50	82,88,114,116	0
2	SO4	A	1471	5/5	0.92	0.27	56,62,87,100	0
2	SO4	C	1456	5/5	0.93	0.19	57,80,88,91	0
3	BME	B	1461	4/4	0.93	0.12	46,60,70,74	0
2	SO4	C	1455	5/5	0.94	0.55	84,86,104,110	0
2	SO4	A	1470	5/5	0.94	0.14	60,64,83,109	0
2	SO4	D	1456	5/5	0.94	0.16	69,81,105,117	0
2	SO4	B	1458	5/5	0.95	0.16	67,79,81,85	0
2	SO4	B	1454	5/5	0.96	0.12	52,58,65,95	0
2	SO4	A	1469	5/5	0.97	0.17	64,68,70,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.