



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2024 – 12:15 PM EDT

PDB ID : 8URF
Title : Crystal Structure of human ASGR2 CRD (Carbohydrate Recognition Domain)
bound to 8G8 Fab
Authors : Sampathumar, P.; Li, Y.
Deposited on : 2023-10-25
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	214	 5% 86% 13%
2	H	232	 3% 84% 7% 9%
3	A	159	 % 73% 7% 19%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	CL	A	407	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4649 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 8G8 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	211	1646	1031	279	331	5	0	2	0

- Molecule 2 is a protein called 8G8 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	212	1627	1030	272	318	7	0	2	0

- Molecule 3 is a protein called Asialoglycoprotein receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	128	1083	686	190	201	6	0	1	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	312	SER	-	expression tag	UNP P07307
A	313	GLY	-	expression tag	UNP P07307
A	314	ARG	-	expression tag	UNP P07307
A	315	GLY	-	expression tag	UNP P07307
A	316	LEU	-	expression tag	UNP P07307
A	317	ASN	-	expression tag	UNP P07307
A	318	ASP	-	expression tag	UNP P07307
A	319	ILE	-	expression tag	UNP P07307
A	320	PHE	-	expression tag	UNP P07307
A	321	GLU	-	expression tag	UNP P07307
A	322	ALA	-	expression tag	UNP P07307
A	323	GLN	-	expression tag	UNP P07307
A	324	LYS	-	expression tag	UNP P07307
A	325	ILE	-	expression tag	UNP P07307

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	326	GLU	-	expression tag	UNP P07307
A	327	TRP	-	expression tag	UNP P07307
A	328	HIS	-	expression tag	UNP P07307
A	329	GLU	-	expression tag	UNP P07307
A	330	HIS	-	expression tag	UNP P07307
A	331	HIS	-	expression tag	UNP P07307
A	332	HIS	-	expression tag	UNP P07307
A	333	HIS	-	expression tag	UNP P07307
A	334	HIS	-	expression tag	UNP P07307
A	335	HIS	-	expression tag	UNP P07307

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total Ca 3 3	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	L	96	Total O 97 97	0	1
7	H	74	Total O 74 74	0	0
7	A	88	Total O 88 88	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	102.41Å 102.41Å 358.99Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.54 – 1.90 33.54 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.54-1.90) 100.0 (33.54-1.90)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.89Å)	Xtrriage
Refinement program	PHENIX (1.14rc1_3177: ???)	Depositor
R, R_{free}	0.166 , 0.203 0.166 , 0.204	Depositor DCC
R_{free} test set	2844 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtrriage
Anisotropy	0.195	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4649	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL, CA, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.64	0/1689	0.74	0/2292
2	H	0.58	0/1667	0.73	1/2275 (0.0%)
3	A	0.75	0/1127	0.70	0/1539
All	All	0.65	0/4483	0.72	1/6106 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	106	ASP	CB-CG-OD1	5.17	122.95	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1646	0	1596	20	0
2	H	1627	0	1588	10	0
3	A	1083	0	955	9	0
4	A	18	0	19	2	0
4	H	6	0	7	0	0
4	L	6	0	8	0	0
5	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	1	0	0	2	0
7	A	88	0	0	0	0
7	H	74	0	0	1	0
7	L	97	0	0	0	0
All	All	4649	0	4173	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:270:HIS:NE2	6:A:407:CL:CL	2.28	1.03
2:H:40[B]:ARG:NH1	7:H:401:HOH:O	2.05	0.88
1:L:185:ASP:OD2	1:L:185:ASP:N	2.30	0.64
1:L:182:SER:OG	1:L:185:ASP:OD2	2.09	0.64
1:L:105:GLU:HG2	1:L:166:GLN:OE1	2.05	0.57
1:L:190:LYS:CD	1:L:210:ASN:HB2	2.35	0.57
3:A:218:TRP:HH2	4:A:405:GOL:H31	1.73	0.54
1:L:190:LYS:HD2	1:L:210:ASN:HB2	1.91	0.52
1:L:169:LYS:N	1:L:169:LYS:HD3	2.25	0.51
2:H:30:LEU:HD21	2:H:74:THR:HG23	1.93	0.51
1:L:89:GLN:HB2	1:L:98:PHE:CD2	2.48	0.49
1:L:190:LYS:O	1:L:210:ASN:HA	2.13	0.49
2:H:40[A]:ARG:HH21	2:H:40[A]:ARG:HG2	1.78	0.48
1:L:105:GLU:OE1	1:L:173:TYR:OH	2.23	0.47
2:H:3:GLN:HE21	2:H:5[B]:GLN:HG3	1.79	0.47
1:L:187:GLU:OE2	1:L:187:GLU:HA	2.12	0.47
1:L:61:ARG:HB2	1:L:76:ASN:O	2.14	0.47
1:L:186:TYR:O	1:L:192:TYR:OH	2.27	0.47
2:H:40[A]:ARG:HG2	2:H:40[A]:ARG:NH2	2.29	0.46
3:A:272:LEU:O	3:A:291:PHE:HB3	2.15	0.46
3:A:218:TRP:CH2	4:A:405:GOL:H31	2.52	0.45
2:H:71:ASN:OD1	2:H:80:TYR:HB2	2.17	0.45
3:A:177:CYS:SG	3:A:183:GLU:HB2	2.58	0.44
1:L:168:SER:C	1:L:169:LYS:HD3	2.39	0.43
1:L:75:ILE:HG21	1:L:78:LEU:HD23	2.01	0.43
3:A:239:ASP:OD1	3:A:277:ASP:HA	2.18	0.43
1:L:47:LEU:HA	1:L:58:VAL:HG21	2.01	0.42
1:L:37:GLN:HB2	1:L:47:LEU:HD11	2.02	0.42
1:L:78:LEU:HD22	1:L:104:LEU:HD21	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:270:HIS:CD2	6:A:407:CL:CL	3.08	0.42
2:H:99:SER:HB3	2:H:105:LEU:HD12	2.02	0.42
2:H:13:ARG:NH2	2:H:118:SER:O	2.52	0.42
3:A:241:ASP:OD2	3:A:245:LYS:HE2	2.20	0.42
1:L:28:ASN:OD1	1:L:68:GLY:HA2	2.21	0.41
2:H:65:LYS:HG3	2:H:66:ASP:N	2.36	0.41
1:L:159:SER:HA	1:L:178:THR:O	2.20	0.41
3:A:257:LYS:HA	3:A:287:TRP:CZ3	2.56	0.40
1:L:154:LEU:HD12	1:L:154:LEU:H	1.86	0.40
2:H:38:LYS:HE2	2:H:40[A]:ARG:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	211/214 (99%)	203 (96%)	8 (4%)	0	100	100
2	H	210/232 (90%)	209 (100%)	1 (0%)	0	100	100
3	A	127/159 (80%)	126 (99%)	1 (1%)	0	100	100
All	All	548/605 (91%)	538 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	187/187 (100%)	186 (100%)	1 (0%)	88	89
2	H	183/197 (93%)	180 (98%)	3 (2%)	62	60
3	A	115/139 (83%)	114 (99%)	1 (1%)	78	79
All	All	485/523 (93%)	480 (99%)	5 (1%)	81	76

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	52	LYS
2	H	40[A]	ARG
2	H	40[B]	ARG
2	H	77	ASN
3	A	241	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	3	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCA	H	1	2	7,8,9	1.91	1 (14%)	9,10,12	2.31	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	1	PCA	CD-N	4.81	1.47	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	PCA	CA-N-CD	-3.39	101.97	113.58
2	H	1	PCA	OE-CD-CG	-3.32	120.97	126.76
2	H	1	PCA	CB-CA-N	2.93	111.71	103.30
2	H	1	PCA	CB-CA-C	-2.88	108.74	112.70
2	H	1	PCA	CG-CD-N	2.37	114.52	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	406	-	5,5,5	1.38	1 (20%)	5,5,5	0.86	0
4	GOL	H	301	-	5,5,5	1.43	1 (20%)	5,5,5	0.73	0
4	GOL	A	405	-	5,5,5	1.57	1 (20%)	5,5,5	0.71	0
4	GOL	A	404	5	5,5,5	1.49	1 (20%)	5,5,5	0.75	0
4	GOL	L	301	-	5,5,5	1.44	1 (20%)	5,5,5	1.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	406	-	-	4/4/4/4	-
4	GOL	H	301	-	-	0/4/4/4	-
4	GOL	A	405	-	-	0/4/4/4	-
4	GOL	A	404	5	-	0/4/4/4	-
4	GOL	L	301	-	-	2/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	L	301	GOL	C3-C2	2.79	1.63	1.51
4	H	301	GOL	O2-C2	-2.33	1.36	1.43
4	A	404	GOL	C1-C2	2.26	1.61	1.51
4	A	405	GOL	O2-C2	-2.20	1.36	1.43
4	A	406	GOL	O2-C2	-2.11	1.37	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	301	GOL	C1-C2-C3-O3
4	A	406	GOL	O1-C1-C2-C3
4	A	406	GOL	C1-C2-C3-O3
4	L	301	GOL	O2-C2-C3-O3
4	A	406	GOL	O1-C1-C2-O2
4	A	406	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	405	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	211/214 (98%)	0.06	10 (4%) 31 34	33, 45, 85, 102	0
2	H	211/232 (90%)	0.18	7 (3%) 46 49	34, 49, 71, 96	0
3	A	128/159 (80%)	-0.02	1 (0%) 86 87	31, 40, 62, 91	0
All	All	550/605 (90%)	0.09	18 (3%) 46 49	31, 45, 78, 102	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	195	GLY	3.8
1	L	190	LYS	3.6
1	L	184	ALA	3.6
1	L	182	SER	3.4
1	L	1	ASP	3.3
1	L	185	ASP	3.0
2	H	33	TRP	2.8
1	L	152	ASN	2.7
2	H	196	THR	2.7
1	L	188	LYS	2.7
2	H	105	LEU	2.5
1	L	169	LYS	2.4
3	A	272	LEU	2.3
2	H	163	ALA	2.2
1	L	189	HIS	2.1
1	L	151	ASP	2.1
2	H	57	SER	2.0
2	H	217	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	H	1	8/9	0.94	0.15	49,64,74,81	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	L	301	6/6	0.71	0.18	58,65,75,75	0
4	GOL	A	406	6/6	0.73	0.20	67,71,77,80	0
4	GOL	H	301	6/6	0.86	0.17	59,61,65,68	0
4	GOL	A	405	6/6	0.94	0.15	39,56,65,73	0
4	GOL	A	404	6/6	0.98	0.09	33,35,37,43	0
5	CA	A	401	1/1	0.99	0.07	39,39,39,39	0
5	CA	A	402	1/1	0.99	0.11	33,33,33,33	0
6	CL	A	407	1/1	0.99	0.04	55,55,55,55	1
5	CA	A	403	1/1	1.00	0.09	33,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CA A 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.