



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2023 – 02:47 AM EDT

PDB ID : 6UTE
Title : Crystal structure of Z032 Fab in complex with WNV EDIII
Authors : Esswein, S.R.; Gristick, H.B.; Keeffe, J.R.; Bjorkman, P.J.
Deposited on : 2019-10-29
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 17255 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Z032 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	219	Total 1621	C 1023	N 270	O 322	S 6	0	0	0
1	A	223	Total 1651	C 1041	N 276	O 328	S 6	0	0	0
1	E	220	Total 1627	C 1026	N 271	O 324	S 6	0	0	0
1	G	223	Total 1651	C 1041	N 276	O 328	S 6	0	0	0
1	I	224	Total 1657	C 1044	N 277	O 330	S 6	0	0	0

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	221	HIS	-	expression tag	UNP S6B291
C	222	HIS	-	expression tag	UNP S6B291
C	223	HIS	-	expression tag	UNP S6B291
C	224	HIS	-	expression tag	UNP S6B291
C	225	HIS	-	expression tag	UNP S6B291
A	221	HIS	-	expression tag	UNP S6B291
A	222	HIS	-	expression tag	UNP S6B291
A	223	HIS	-	expression tag	UNP S6B291
A	224	HIS	-	expression tag	UNP S6B291
A	225	HIS	-	expression tag	UNP S6B291
E	221	HIS	-	expression tag	UNP S6B291
E	222	HIS	-	expression tag	UNP S6B291
E	223	HIS	-	expression tag	UNP S6B291
E	224	HIS	-	expression tag	UNP S6B291
E	225	HIS	-	expression tag	UNP S6B291
G	221	HIS	-	expression tag	UNP S6B291
G	222	HIS	-	expression tag	UNP S6B291
G	223	HIS	-	expression tag	UNP S6B291
G	224	HIS	-	expression tag	UNP S6B291

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	225	HIS	-	expression tag	UNP S6B291
I	221	HIS	-	expression tag	UNP S6B291
I	222	HIS	-	expression tag	UNP S6B291
I	223	HIS	-	expression tag	UNP S6B291
I	224	HIS	-	expression tag	UNP S6B291
I	225	HIS	-	expression tag	UNP S6B291

- Molecule 2 is a protein called Z032 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	213	1662	1047	277	332	6	0	0	0
2	B	213	1662	1047	277	332	6	0	0	0
2	F	213	1662	1047	277	332	6	0	0	0
2	H	210	1638	1029	274	329	6	0	0	0
2	J	213	1662	1047	277	332	6	0	0	0

- Molecule 3 is a protein called Envelope domain III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	S	98	732	467	125	138	2	0	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	F	1	Total	C	O	0	0
			6	3	3		
4	J	1	Total	C	O	0	0
			6	3	3		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.23Å 114.02Å 127.26Å 90.00° 109.50° 90.00°	Depositor
Resolution (Å)	39.99 – 2.90	Depositor
% Data completeness (in resolution range)	99.7 (39.99-2.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.90Å)	Xtrriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.223 , 0.264	Depositor
Wilson B-factor (Å ²)	54.3	Xtrriage
Anisotropy	0.407	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtrriage
Total number of atoms	17255	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	B	501	-	5,5,5	0.91	0	5,5,5	1.00	0
4	GOL	F	601	-	5,5,5	0.88	0	5,5,5	1.04	0
4	GOL	J	301	-	5,5,5	0.88	0	5,5,5	1.00	0
4	GOL	D	302	-	5,5,5	0.88	0	5,5,5	1.00	0
4	GOL	D	301	-	5,5,5	0.91	0	5,5,5	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	501	-	-	2/4/4/4	-
4	GOL	F	601	-	-	2/4/4/4	-
4	GOL	J	301	-	-	0/4/4/4	-
4	GOL	D	302	-	-	0/4/4/4	-
4	GOL	D	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	601	GOL	O1-C1-C2-C3
4	F	601	GOL	O1-C1-C2-O2
4	B	501	GOL	O1-C1-C2-C3
4	B	501	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

4.8 Polymer linkage issues

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.