



## Full wwPDB EM Validation Report ⓘ

Feb 27, 2024 – 08:51 AM EST

PDB ID : 6UTG  
EMDB ID : EMD-20878  
Title : Allosteric coupling between alpha-rings of the 20S proteasome, 20S singly capped with a PA26/V230F  
Authors : Yu, Z.; Cheng, Y.  
Deposited on : 2019-10-29  
Resolution : 3.40 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	203	6% 98% .
1	2	203	5% 98% .
1	H	203	6% 98% .
1	I	203	6% 98% .
1	J	203	6% 98% .
1	K	203	6% 98% .
1	L	203	7% 98% .
1	M	203	7% 98% .
1	N	203	7% 98% .


*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	V	203	6% 98%
1	W	203	5% 98%
1	X	203	7% 98%
1	Y	203	5% 98%
1	Z	203	6% 98%
2	A	227	• 99%
2	B	227	• 98%
2	C	227	• 98%
2	D	227	• 98%
2	E	227	• 99%
2	F	227	• 99%
2	G	227	5% 98%
3	O	228	7% 95%
3	P	228	7% 95%
3	Q	228	7% 95%
3	R	228	7% 95%
3	S	228	7% 95%
3	T	228	7% 95%
3	U	228	7% 95%
4	a	224	9% 98%
4	b	224	10% 98%
4	c	224	8% 98%
4	d	224	8% 98%
4	e	224	8% 98%
4	f	224	7% 98%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	g	224	 8% 98%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 58149 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	1	203	1558	985	264	298	11	0	0
1	2	203	1558	985	264	298	11	0	0
1	H	203	1558	985	264	298	11	0	0
1	I	203	1558	985	264	298	11	0	0
1	J	203	1558	985	264	298	11	0	0
1	K	203	1558	985	264	298	11	0	0
1	L	203	1558	985	264	298	11	0	0
1	M	203	1558	985	264	298	11	0	0
1	N	203	1558	985	264	298	11	0	0
1	V	203	1558	985	264	298	11	0	0
1	W	203	1558	985	264	298	11	0	0
1	X	203	1558	985	264	298	11	0	0
1	Y	203	1558	985	264	298	11	0	0
1	Z	203	1558	985	264	298	11	0	0

- Molecule 2 is a protein called Proteasome subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	A	227	1768	1123	299	343	3	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	227	Total	C	N	O	S	0	0
			1768	1123	299	343	3		
2	C	227	Total	C	N	O	S	0	0
			1768	1123	299	343	3		
2	D	227	Total	C	N	O	S	0	0
			1768	1123	299	343	3		
2	E	227	Total	C	N	O	S	0	0
			1768	1123	299	343	3		
2	F	227	Total	C	N	O	S	0	0
			1768	1123	299	343	3		
2	G	227	Total	C	N	O	S	0	0
			1768	1123	299	343	3		

- Molecule 3 is a protein called Proteasome activator protein PA26.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	O	218	Total	C	N	O	S	0	0
			1683	1056	296	325	6		
3	P	218	Total	C	N	O	S	0	0
			1683	1056	296	325	6		
3	Q	218	Total	C	N	O	S	0	0
			1683	1056	296	325	6		
3	R	218	Total	C	N	O	S	0	0
			1683	1056	296	325	6		
3	S	218	Total	C	N	O	S	0	0
			1683	1056	296	325	6		
3	T	218	Total	C	N	O	S	0	0
			1683	1056	296	325	6		
3	U	218	Total	C	N	O	S	0	0
			1683	1056	296	325	6		

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	49	VAL	THR	conflict	UNP Q9U8G2
O	226	THR	SER	conflict	UNP Q9U8G2
O	230	PHE	VAL	engineered mutation	UNP Q9U8G2
P	49	VAL	THR	conflict	UNP Q9U8G2
P	226	THR	SER	conflict	UNP Q9U8G2
P	230	PHE	VAL	engineered mutation	UNP Q9U8G2
Q	49	VAL	THR	conflict	UNP Q9U8G2
Q	226	THR	SER	conflict	UNP Q9U8G2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Q	230	PHE	VAL	engineered mutation	UNP Q9U8G2
R	49	VAL	THR	conflict	UNP Q9U8G2
R	226	THR	SER	conflict	UNP Q9U8G2
R	230	PHE	VAL	engineered mutation	UNP Q9U8G2
S	49	VAL	THR	conflict	UNP Q9U8G2
S	226	THR	SER	conflict	UNP Q9U8G2
S	230	PHE	VAL	engineered mutation	UNP Q9U8G2
T	49	VAL	THR	conflict	UNP Q9U8G2
T	226	THR	SER	conflict	UNP Q9U8G2
T	230	PHE	VAL	engineered mutation	UNP Q9U8G2
U	49	VAL	THR	conflict	UNP Q9U8G2
U	226	THR	SER	conflict	UNP Q9U8G2
U	230	PHE	VAL	engineered mutation	UNP Q9U8G2

- Molecule 4 is a protein called Proteasome subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	a	224	1740	1104	295	338	3	0	0
4	b	224	1740	1104	295	338	3	0	0
4	c	224	1740	1104	295	338	3	0	0
4	d	224	1740	1104	295	338	3	0	0
4	e	224	1740	1104	295	338	3	0	0
4	f	224	1740	1104	295	338	3	0	0
4	g	224	1740	1104	295	338	3	0	0


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	66	ALA	LYS	engineered mutation	UNP P25156
b	66	ALA	LYS	engineered mutation	UNP P25156
c	66	ALA	LYS	engineered mutation	UNP P25156
d	66	ALA	LYS	engineered mutation	UNP P25156
e	66	ALA	LYS	engineered mutation	UNP P25156
f	66	ALA	LYS	engineered mutation	UNP P25156
g	66	ALA	LYS	engineered mutation	UNP P25156


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta



- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta







- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 1: Proteasome subunit beta




- Molecule 2: Proteasome subunit alpha




- Molecule 2: Proteasome subunit alpha



- Molecule 2: Proteasome subunit alpha




- Molecule 2: Proteasome subunit alpha




- Molecule 2: Proteasome subunit alpha






- Molecule 2: Proteasome subunit alpha




- Molecule 2: Proteasome subunit alpha




- Molecule 3: Proteasome activator protein PA26




- Molecule 3: Proteasome activator protein PA26



- Molecule 3: Proteasome activator protein PA26




- Molecule 3: Proteasome activator protein PA26




- Molecule 3: Proteasome activator protein PA26

Chain S:  7% 95%




• Molecule 3: Proteasome activator protein PA26

Chain T:  7% 95%




• Molecule 3: Proteasome activator protein PA26

Chain U:  7% 95%




• Molecule 4: Proteasome subunit alpha

Chain a:  9% 98%




• Molecule 4: Proteasome subunit alpha

Chain b:  10% 98%




• Molecule 4: Proteasome subunit alpha

Chain c:  8% 98%




• Molecule 4: Proteasome subunit alpha


Chain d:  8% 98%




## • Molecule 4: Proteasome subunit alpha



## • Molecule 4: Proteasome subunit alpha



## • Molecule 4: Proteasome subunit alpha



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	50000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TALOS ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	60	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	3.313	Depositor
Minimum map value	-2.116	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.114	Depositor
Recommended contour level	0.472	Depositor
Map size ( $\text{\AA}$ )	394.88, 394.88, 394.88	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.234, 1.234, 1.234	Depositor

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	1	0.47	0/1577	0.67	2/2129 (0.1%)
1	2	0.47	0/1577	0.67	2/2129 (0.1%)
1	H	0.47	0/1577	0.67	2/2129 (0.1%)
1	I	0.47	0/1577	0.67	2/2129 (0.1%)
1	J	0.47	0/1577	0.67	2/2129 (0.1%)
1	K	0.47	0/1577	0.67	2/2129 (0.1%)
1	L	0.47	0/1577	0.67	2/2129 (0.1%)
1	M	0.47	0/1577	0.67	2/2129 (0.1%)
1	N	0.47	0/1577	0.67	2/2129 (0.1%)
1	V	0.47	0/1577	0.67	2/2129 (0.1%)
1	W	0.47	0/1577	0.67	2/2129 (0.1%)
1	X	0.47	0/1577	0.67	2/2129 (0.1%)
1	Y	0.47	0/1577	0.67	2/2129 (0.1%)
1	Z	0.47	0/1577	0.67	2/2129 (0.1%)
2	A	0.56	0/1792	0.71	0/2416
2	B	0.57	1/1792 (0.1%)	0.71	0/2416
2	C	0.56	0/1792	0.71	0/2416
2	D	0.56	0/1792	0.71	1/2416 (0.0%)
2	E	0.55	0/1792	0.71	0/2416
2	F	0.56	0/1792	0.71	0/2416
2	G	0.54	0/1792	0.71	0/2416
3	O	0.39	0/1707	0.51	0/2307
3	P	0.38	0/1707	0.51	0/2307
3	Q	0.38	0/1707	0.51	0/2307
3	R	0.38	0/1707	0.51	0/2307
3	S	0.39	0/1707	0.51	0/2307
3	T	0.38	0/1707	0.51	0/2307
3	U	0.39	0/1707	0.51	0/2307
4	a	0.46	0/1763	0.77	2/2376 (0.1%)
4	b	0.46	0/1763	0.76	2/2376 (0.1%)
4	c	0.46	0/1763	0.77	2/2376 (0.1%)
4	d	0.46	0/1763	0.76	2/2376 (0.1%)
4	e	0.46	0/1763	0.77	2/2376 (0.1%)
4	f	0.46	0/1763	0.76	2/2376 (0.1%)

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
4	g	0.46	0/1763	0.76	2/2376 (0.1%)
All	All	0.47	1/58912 (0.0%)	0.67	43/79499 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	1	0	1
1	2	0	1
1	H	0	1
1	I	0	1
1	J	0	1
1	K	0	1
1	L	0	1
1	M	0	1
1	N	0	1
1	V	0	1
1	W	0	1
1	X	0	1
1	Y	0	1
1	Z	0	1
2	A	0	2
2	B	0	2
2	C	0	2
2	D	0	2
2	E	0	2
2	F	0	2
2	G	0	2
4	a	0	1
4	b	0	1
4	c	0	1
4	d	0	1
4	e	0	1
4	f	0	1
4	g	0	1
All	All	0	35

All (1) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	21	LEU	CA-CB	-5.13	1.42	1.53

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	2	21	THR	C-N-CA	7.40	140.21	121.70
1	K	21	THR	C-N-CA	7.39	140.17	121.70
1	1	21	THR	C-N-CA	7.39	140.17	121.70
1	N	21	THR	C-N-CA	7.39	140.17	121.70
1	H	21	THR	C-N-CA	7.38	140.16	121.70
1	Y	21	THR	C-N-CA	7.38	140.16	121.70
1	J	21	THR	C-N-CA	7.38	140.15	121.70
1	X	21	THR	C-N-CA	7.38	140.15	121.70
1	V	21	THR	C-N-CA	7.38	140.14	121.70
1	L	21	THR	C-N-CA	7.37	140.13	121.70
1	I	21	THR	C-N-CA	7.37	140.12	121.70
1	W	21	THR	C-N-CA	7.36	140.11	121.70
1	M	21	THR	C-N-CA	7.36	140.10	121.70
1	Z	21	THR	C-N-CA	7.35	140.09	121.70
1	I	68	LEU	CB-CG-CD2	-6.84	99.37	111.00
1	1	68	LEU	CB-CG-CD2	-6.83	99.39	111.00
1	J	68	LEU	CB-CG-CD2	-6.83	99.39	111.00
1	V	68	LEU	CB-CG-CD2	-6.83	99.39	111.00
1	K	68	LEU	CB-CG-CD2	-6.83	99.39	111.00
1	N	68	LEU	CB-CG-CD2	-6.83	99.39	111.00
1	Y	68	LEU	CB-CG-CD2	-6.83	99.39	111.00
1	M	68	LEU	CB-CG-CD2	-6.82	99.40	111.00
1	L	68	LEU	CB-CG-CD2	-6.82	99.40	111.00
1	2	68	LEU	CB-CG-CD2	-6.82	99.41	111.00
1	W	68	LEU	CB-CG-CD2	-6.82	99.41	111.00
1	Z	68	LEU	CB-CG-CD2	-6.82	99.41	111.00
1	H	68	LEU	CB-CG-CD2	-6.81	99.42	111.00
1	X	68	LEU	CB-CG-CD2	-6.80	99.43	111.00
4	e	152	ASP	CB-CG-OD1	6.36	124.02	118.30
4	c	152	ASP	CB-CG-OD1	6.29	123.96	118.30
4	f	152	ASP	CB-CG-OD1	6.25	123.93	118.30
4	d	152	ASP	CB-CG-OD1	6.24	123.92	118.30
4	g	152	ASP	CB-CG-OD1	6.24	123.92	118.30
4	a	152	ASP	CB-CG-OD1	6.21	123.89	118.30
4	b	152	ASP	CB-CG-OD1	6.21	123.88	118.30
4	f	207	LEU	C-N-CA	6.07	136.87	121.70
4	a	207	LEU	C-N-CA	6.07	136.86	121.70
4	c	207	LEU	C-N-CA	6.06	136.86	121.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	e	207	LEU	C-N-CA	6.06	136.85	121.70
4	d	207	LEU	C-N-CA	6.05	136.84	121.70
4	b	207	LEU	C-N-CA	6.05	136.83	121.70
4	g	207	LEU	C-N-CA	6.05	136.83	121.70
2	D	221	TYR	CA-CB-CG	-5.17	103.58	113.40

There are no chirality outliers.

All (35) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	1	24	ASN	Peptide
1	2	24	ASN	Peptide
2	A	63	SER	Peptide
2	A	8	TYR	Peptide
2	B	63	SER	Peptide
2	B	8	TYR	Peptide
2	C	63	SER	Peptide
2	C	8	TYR	Peptide
2	D	63	SER	Peptide
2	D	8	TYR	Peptide
2	E	63	SER	Peptide
2	E	8	TYR	Peptide
2	F	63	SER	Peptide
2	F	8	TYR	Peptide
2	G	63	SER	Peptide
2	G	8	TYR	Peptide
1	H	24	ASN	Peptide
1	I	24	ASN	Peptide
1	J	24	ASN	Peptide
1	K	24	ASN	Peptide
1	L	24	ASN	Peptide
1	M	24	ASN	Peptide
1	N	24	ASN	Peptide
1	V	24	ASN	Peptide
1	W	24	ASN	Peptide
1	X	24	ASN	Peptide
1	Y	24	ASN	Peptide
1	Z	24	ASN	Peptide
4	a	207	LEU	Peptide
4	b	207	LEU	Peptide
4	c	207	LEU	Peptide
4	d	207	LEU	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
4	e	207	LEU	Peptide
4	f	207	LEU	Peptide
4	g	207	LEU	Peptide

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	2	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	H	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	I	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	J	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	K	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	L	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	M	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	N	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	V	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	W	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	X	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	Y	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
1	Z	201/203 (99%)	189 (94%)	11 (6%)	1 (0%)	29	61
2	A	225/227 (99%)	215 (96%)	9 (4%)	1 (0%)	34	67
2	B	225/227 (99%)	215 (96%)	9 (4%)	1 (0%)	34	67

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	225/227 (99%)	216 (96%)	8 (4%)	1 (0%)	34	67
2	D	225/227 (99%)	215 (96%)	9 (4%)	1 (0%)	34	67
2	E	225/227 (99%)	215 (96%)	9 (4%)	1 (0%)	34	67
2	F	225/227 (99%)	215 (96%)	9 (4%)	1 (0%)	34	67
2	G	225/227 (99%)	216 (96%)	8 (4%)	1 (0%)	34	67
3	O	214/228 (94%)	203 (95%)	11 (5%)	0	100	100
3	P	214/228 (94%)	203 (95%)	11 (5%)	0	100	100
3	Q	214/228 (94%)	203 (95%)	11 (5%)	0	100	100
3	R	214/228 (94%)	203 (95%)	11 (5%)	0	100	100
3	S	214/228 (94%)	203 (95%)	11 (5%)	0	100	100
3	T	214/228 (94%)	203 (95%)	11 (5%)	0	100	100
3	U	214/228 (94%)	203 (95%)	11 (5%)	0	100	100
4	a	222/224 (99%)	209 (94%)	12 (5%)	1 (0%)	29	61
4	b	222/224 (99%)	209 (94%)	12 (5%)	1 (0%)	29	61
4	c	222/224 (99%)	209 (94%)	12 (5%)	1 (0%)	29	61
4	d	222/224 (99%)	209 (94%)	12 (5%)	1 (0%)	29	61
4	e	222/224 (99%)	209 (94%)	12 (5%)	1 (0%)	29	61
4	f	222/224 (99%)	209 (94%)	12 (5%)	1 (0%)	29	61
4	g	222/224 (99%)	209 (94%)	12 (5%)	1 (0%)	29	61
All	All	7441/7595 (98%)	7037 (95%)	376 (5%)	28 (0%)	38	67

All (28) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	a	208	LYS
4	b	208	LYS
4	c	208	LYS
4	d	208	LYS
4	e	208	LYS
4	f	208	LYS
4	g	208	LYS
1	M	25	PHE
1	Y	25	PHE
1	Z	25	PHE
1	1	25	PHE

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	2	25	PHE
1	H	25	PHE
1	I	25	PHE
1	J	25	PHE
1	K	25	PHE
1	L	25	PHE
1	N	25	PHE
1	V	25	PHE
1	W	25	PHE
1	X	25	PHE
2	A	64	ILE
2	B	64	ILE
2	C	64	ILE
2	E	64	ILE
2	F	64	ILE
2	D	64	ILE
2	G	64	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	2	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	H	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	I	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	J	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	K	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	L	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	M	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	N	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	V	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	W	170/170 (100%)	169 (99%)	1 (1%)	86	94

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	Y	170/170 (100%)	169 (99%)	1 (1%)	86	94
1	Z	170/170 (100%)	169 (99%)	1 (1%)	86	94
2	A	188/188 (100%)	188 (100%)	0	100	100
2	B	188/188 (100%)	188 (100%)	0	100	100
2	C	188/188 (100%)	187 (100%)	1 (0%)	88	94
2	D	188/188 (100%)	188 (100%)	0	100	100
2	E	188/188 (100%)	188 (100%)	0	100	100
2	F	188/188 (100%)	188 (100%)	0	100	100
2	G	188/188 (100%)	187 (100%)	1 (0%)	88	94
3	O	179/187 (96%)	178 (99%)	1 (1%)	86	94
3	P	179/187 (96%)	178 (99%)	1 (1%)	86	94
3	Q	179/187 (96%)	178 (99%)	1 (1%)	86	94
3	R	179/187 (96%)	178 (99%)	1 (1%)	86	94
3	S	179/187 (96%)	178 (99%)	1 (1%)	86	94
3	T	179/187 (96%)	178 (99%)	1 (1%)	86	94
3	U	179/187 (96%)	178 (99%)	1 (1%)	86	94
4	a	185/185 (100%)	183 (99%)	2 (1%)	73	86
4	b	185/185 (100%)	183 (99%)	2 (1%)	73	86
4	c	185/185 (100%)	183 (99%)	2 (1%)	73	86
4	d	185/185 (100%)	183 (99%)	2 (1%)	73	86
4	e	185/185 (100%)	183 (99%)	2 (1%)	73	86
4	f	185/185 (100%)	183 (99%)	2 (1%)	73	86
4	g	185/185 (100%)	183 (99%)	2 (1%)	73	86
All	All	6244/6300 (99%)	6207 (99%)	37 (1%)	86	94

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1	178	ILE
1	2	178	ILE
2	C	222	ARG
2	G	222	ARG
1	H	178	ILE

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	I	178	ILE
1	J	178	ILE
1	K	178	ILE
1	L	178	ILE
1	M	178	ILE
1	N	178	ILE
3	O	184	ARG
3	P	184	ARG
3	Q	184	ARG
3	R	184	ARG
3	S	184	ARG
3	T	184	ARG
3	U	184	ARG
1	V	178	ILE
1	W	178	ILE
1	X	178	ILE
1	Y	178	ILE
1	Z	178	ILE
4	a	111	ASN
4	a	212	ILE
4	b	111	ASN
4	b	212	ILE
4	c	111	ASN
4	c	212	ILE
4	d	111	ASN
4	d	212	ILE
4	e	111	ASN
4	e	212	ILE
4	f	111	ASN
4	f	212	ILE
4	g	111	ASN
4	g	212	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (33) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	219	ASN
2	B	219	ASN
2	C	219	ASN
2	D	158	ASN
2	D	219	ASN
2	E	158	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	E	219	ASN
2	F	219	ASN
2	G	219	ASN
1	J	141	GLN
3	O	47	HIS
3	O	200	HIS
3	P	47	HIS
3	P	200	HIS
3	Q	47	HIS
3	Q	185	GLN
3	Q	200	HIS
3	R	47	HIS
3	R	72	GLN
3	R	200	HIS
3	S	47	HIS
3	S	200	HIS
3	T	47	HIS
3	T	200	HIS
3	U	47	HIS
3	U	200	HIS
4	a	23	GLN
4	b	23	GLN
4	c	23	GLN
4	d	23	GLN
4	e	23	GLN
4	f	23	GLN
4	g	23	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20878. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


### 6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 139




Y Index: 176




Z Index: 156


### 6.3.2 Raw map



X Index: 172



Y Index: 146




Z Index: 187


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.472. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [i](#)




The volume at the recommended contour level is 381 nm<sup>3</sup>; this corresponds to an approximate mass of 344 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.294 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.294 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.26	6.18	4.35

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.26 differs from the reported value 3.4 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-20878 and PDB model 6UTG. Per-residue inclusion information can be found in section [3](#) on page [8](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.472 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.472).









































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.472) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7360	 0.4310
1	 0.7800	 0.4600
2	 0.7810	 0.4600
A	 0.7170	 0.4200
B	 0.7170	 0.4190
C	 0.7140	 0.4180
D	 0.7130	 0.4170
E	 0.7170	 0.4160
F	 0.7180	 0.4170
G	 0.7200	 0.4180
H	 0.7620	 0.4480
I	 0.7720	 0.4500
J	 0.7690	 0.4500
K	 0.7670	 0.4510
L	 0.7690	 0.4510
M	 0.7660	 0.4510
N	 0.7680	 0.4510
O	 0.7690	 0.4320
P	 0.7690	 0.4330
Q	 0.7670	 0.4280
R	 0.7680	 0.4360
S	 0.7700	 0.4330
T	 0.7670	 0.4330
U	 0.7620	 0.4330
V	 0.7810	 0.4580
W	 0.7830	 0.4600
X	 0.7820	 0.4600
Y	 0.7790	 0.4610
Z	 0.7770	 0.4610
a	 0.6590	 0.3980
b	 0.6550	 0.3990
c	 0.6580	 0.3990
d	 0.6540	 0.3980
e	 0.6560	 0.3980
f	 0.6590	 0.3990
g	 0.6580	 0.3980

