



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 05:33 pm BST

PDB ID : 3UUO  
Title : The discovery of potent, selectivity, and orally bioavailable pyroloquinolines as PDE10 inhibitors for the treatment of Schizophrenia  
Authors : Ho, G.D.; Yang, S.; Smotryski, J.; Bercovici, A.; Nechuta, T.; Smith, E.M.; McElroy, W.; Tan, Z.; Tulshian, D.; Mckittrick, B.; Greenlee, W.J.; Hruza, A.; Xiao, L.; Rindgen, D.; Guzzi, M.; Zhang, X.; Bleickardt, C.; Mullins, D.; Hodgson, R.  
Deposited on : 2011-11-28  
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	 4% 88% 9%
1	B	337	 9% 89% 7%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10218 atoms, of which 5019 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	305	4921	1580	2448	423	448	22	0	0	0
1	B	315	5089	1638	2529	436	463	23	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	434	GLY	-	EXPRESSION TAG	UNP Q9Y233
A	435	SER	-	EXPRESSION TAG	UNP Q9Y233
A	436	HIS	-	EXPRESSION TAG	UNP Q9Y233
A	437	MET	-	EXPRESSION TAG	UNP Q9Y233
A	438	GLY	-	EXPRESSION TAG	UNP Q9Y233
A	446	SER	-	EXPRESSION TAG	UNP Q9Y233
A	447	HIS	-	EXPRESSION TAG	UNP Q9Y233
A	448	MET	-	EXPRESSION TAG	UNP Q9Y233
B	434	GLY	-	EXPRESSION TAG	UNP Q9Y233
B	435	SER	-	EXPRESSION TAG	UNP Q9Y233
B	436	HIS	-	EXPRESSION TAG	UNP Q9Y233
B	437	MET	-	EXPRESSION TAG	UNP Q9Y233
B	438	GLY	-	EXPRESSION TAG	UNP Q9Y233
B	446	SER	-	EXPRESSION TAG	UNP Q9Y233
B	447	HIS	-	EXPRESSION TAG	UNP Q9Y233
B	448	MET	-	EXPRESSION TAG	UNP Q9Y233

- Molecule 2 is 6-methoxy-3,8-dimethyl-4-(piperazin-1-yl)-1H-pyrazolo[3,4-b]quinoline (three-letter code: 0CV) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>5</sub>O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	44	17	21	5	1	0	0
2	B	1	44	17	21	5	1	0	0


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	B	1	1	1	0	0
3	A	1	1	1	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
4	B	1	1	1	0	0
4	A	1	1	1	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O P 5 4 1	0	0
5	B	1	Total O P 5 4 1	0	0



- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	55	Total O 55 55	0	0
6	B	51	Total O 51 51	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.57Å 82.23Å 161.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.79 – 2.11 29.67 – 2.11	Depositor EDS
% Data completeness (in resolution range)	94.2 (29.79-2.11) 94.3 (29.67-2.11)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.73 (at 2.12Å)	Xtrriage
Refinement program	BUSTER 2.11.2	Depositor
R, $R_{free}$	0.195 , 0.227 0.198 , 0.227	Depositor DCC
$R_{free}$ test set	1874 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.9	Xtrriage
Anisotropy	0.730	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 49.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10218	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MG, OCV, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/2534	0.57	0/3430
1	B	0.47	0/2624	0.58	0/3553
All	All	0.46	0/5158	0.57	0/6983

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2473	2448	2446	0	0
1	B	2560	2529	2527	1	0
2	A	23	21	21	0	0
2	B	23	21	21	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	55	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	51	0	0	0	0
All	All	5199	5019	5015	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:483:GLU:HA	1:B:486:TRP:CE2	2.51	0.44

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/337 (90%)	299 (99%)	4 (1%)	0	100	100
1	B	313/337 (93%)	308 (98%)	5 (2%)	0	100	100
All	All	616/674 (91%)	607 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/304 (90%)	265 (96%)	10 (4%)	35	35
1	B	284/304 (93%)	272 (96%)	12 (4%)	30	29
All	All	559/608 (92%)	537 (96%)	22 (4%)	32	32

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	467	ARG
1	A	551	LEU
1	A	635	LEU
1	A	642	ARG
1	A	656	LEU
1	A	686	LYS
1	A	711	ILE
1	A	713	MET
1	A	727	LEU
1	A	760	SER
1	B	467	ARG
1	B	484	ASN
1	B	517	LYS
1	B	582	LEU
1	B	585	LEU
1	B	615	SER
1	B	617	SER
1	B	620	GLU
1	B	656	LEU
1	B	686	LYS
1	B	714	MET
1	B	727	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	PO4	A	773	3,4	4,4,4	2.96	3 (75%)	6,6,6	0.78	0
2	0CV	A	1	-	24,26,26	1.49	4 (16%)	27,38,38	3.64	10 (37%)
5	PO4	B	773	3,4	4,4,4	2.96	2 (50%)	6,6,6	0.51	0
2	0CV	B	1	-	24,26,26	1.50	4 (16%)	27,38,38	3.13	8 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0CV	A	1	-	-	3/6/14/14	0/4/4/4
2	0CV	B	1	-	-	3/6/14/14	0/4/4/4

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	773	PO4	P-O1	4.59	1.61	1.50
5	A	773	PO4	P-O1	4.37	1.61	1.50
2	B	1	0CV	C6-N1	-3.75	1.40	1.46
2	A	1	0CV	C9-C12	3.28	1.51	1.43
2	A	1	0CV	C19-C13	-3.14	1.48	1.50
2	B	1	0CV	C7-C8	3.11	1.45	1.40
2	B	1	0CV	C20-C15	2.94	1.42	1.37
5	B	773	PO4	P-O3	2.60	1.62	1.54

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	773	PO4	P-O3	2.52	1.62	1.54
2	A	1	0CV	C20-C15	2.52	1.41	1.37
2	A	1	0CV	C7-C8	2.51	1.44	1.40
5	A	773	PO4	P-O2	2.46	1.62	1.54
2	B	1	0CV	C13-N18	-2.18	1.31	1.34

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	0CV	C7-C8-C10	15.54	123.50	116.31
2	B	1	0CV	C7-C8-C10	12.88	122.27	116.31
2	A	1	0CV	C6-N1-C2	5.38	123.39	111.52
2	B	1	0CV	C8-C10-N14	-4.19	117.53	123.31
2	B	1	0CV	C6-N1-C2	3.79	119.88	111.52
2	A	1	0CV	C8-C7-C9	-3.71	115.40	122.01
2	A	1	0CV	C8-C10-N14	-3.55	118.41	123.31
2	B	1	0CV	C8-C7-C9	-3.44	115.87	122.01
2	A	1	0CV	C23-C15-C10	3.26	123.19	120.03
2	B	1	0CV	C10-N14-C12	3.14	122.98	116.75
2	A	1	0CV	C15-C10-N14	2.98	122.06	118.23
2	A	1	0CV	C11-C8-C7	-2.96	119.09	124.00
2	B	1	0CV	C16-C11-C8	2.96	123.99	120.05
2	B	1	0CV	C23-C15-C10	2.89	122.83	120.03
2	A	1	0CV	C16-C11-C8	2.83	123.82	120.05
2	B	1	0CV	C15-C10-N14	2.24	121.11	118.23
2	A	1	0CV	C23-C15-C20	-2.18	117.56	120.74
2	A	1	0CV	C11-C16-C20	-2.13	118.56	121.12

There are no chirality outliers.


All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1	0CV	C9-C7-N1-C6
2	B	1	0CV	C9-C7-N1-C6
2	B	1	0CV	C9-C7-N1-C2
2	A	1	0CV	C9-C7-N1-C2
2	A	1	0CV	C8-C7-N1-C2
2	B	1	0CV	C8-C7-N1-C6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	305/337 (90%)	0.16	15 (4%) 29 35	28, 46, 84, 107	0
1	B	315/337 (93%)	0.45	30 (9%) 8 10	30, 48, 96, 132	0
All	All	620/674 (91%)	0.31	45 (7%) 15 19	28, 47, 88, 132	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	457	GLN	6.3
1	A	585	LEU	5.1
1	A	467	ARG	4.9
1	B	533	VAL	4.2
1	A	584	ALA	4.1
1	A	502	CYS	3.8
1	B	551	LEU	3.8
1	B	458	GLY	3.5
1	B	560	CYS	3.5
1	A	501	SER	3.4
1	A	533	VAL	3.4
1	B	536	CYS	3.3
1	B	613	THR	3.3
1	A	466	VAL	3.0
1	B	461	GLN	3.0
1	B	585	LEU	3.0
1	B	534	ALA	2.9
1	B	456	TRP	2.9
1	B	606	GLU	2.9
1	B	616	SER	2.9
1	A	647	GLU	2.9
1	B	532	THR	2.7
1	B	673	CYS	2.7
1	B	559	ALA	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	537	MET	2.7
1	B	670	MET	2.6
1	A	768	GLY	2.6
1	A	673	CYS	2.6
1	B	531	VAL	2.5
1	A	505	LEU	2.5
1	B	619	TYR	2.5
1	A	536	CYS	2.5
1	A	766	ILE	2.4
1	B	563	HIS	2.4
1	B	581	PRO	2.4
1	B	557	LEU	2.4
1	A	532	THR	2.4
1	B	501	SER	2.3
1	B	578	PHE	2.2
1	B	506	GLU	2.1
1	B	676	CYS	2.1
1	B	675	LEU	2.1
1	B	671	THR	2.0
1	B	460	MET	2.0
1	A	674	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	0CV	A	1	23/23	0.94	0.13	39,44,48,50	0
2	0CV	B	1	23/23	0.96	0.12	35,40,42,42	0

*Continued on next page...*






*Continued from previous page...*


<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Atoms</b>	<b>RSCC</b>	<b>RSR</b>	<b>B-factors(Å<sup>2</sup>)</b>	<b>Q&lt;0.9</b>
5	PO4	B	773	5/5	0.97	0.15	43,45,45,46	0
3	MG	A	771	1/1	0.98	0.19	29,29,29,29	0
5	PO4	A	773	5/5	0.99	0.16	32,35,38,40	0
3	MG	B	771	1/1	0.99	0.20	37,37,37,37	0
4	ZN	B	772	1/1	1.00	0.12	37,37,37,37	0
4	ZN	A	772	1/1	1.00	0.15	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 0CV A 1:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.