



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 13, 2024 – 05:39 am BST

PDB ID : 1UWV
Title : Crystal Structure of RumA, the iron-sulfur cluster containing E. coli 23S Ribosomal RNA 5-Methyluridine Methyltransferase
Authors : Lee, T.T.; Agarwalla, S.; Stroud, R.M.
Deposited on : 2004-02-11
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	433	75% 19% . .

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 3378 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 23S RRNA (URACIL-5-)-METHYLTRANSFERASE RUMA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	418	3239	2034	586	601	7	11	0	0	1

- Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe S		
2	A	1	8	4 4	0	0

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Ni 2 2	0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	123	Total O 123 123	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 23S rRNA (URACIL-5-)-METHYLTRANSFERASE RUMA

Chain A:  75% 19%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.94Å 99.45Å 58.52Å 90.00° 100.57° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95	Depositor
% Data completeness (in resolution range)	99.6 (50.00-1.95)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.186 , 0.227	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3378	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, NI, CL, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/3289	0.95	13/4439 (0.3%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	431	ARG	O-C-N	-8.22	109.55	122.70
1	A	358	ASP	CB-CG-OD2	6.52	124.17	118.30
1	A	262	ASP	CB-CG-OD2	6.26	123.93	118.30
1	A	128	ARG	NE-CZ-NH1	6.20	123.40	120.30
1	A	122	ASP	CB-CG-OD2	6.04	123.73	118.30
1	A	363	ASP	CB-CG-OD2	5.98	123.68	118.30
1	A	57	ASP	CB-CG-OD2	5.68	123.42	118.30
1	A	235	ASP	CB-CG-OD2	5.61	123.35	118.30
1	A	76	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	115	ASP	CB-CG-OD2	5.36	123.12	118.30
1	A	397	ASP	CB-CG-OD2	5.33	123.10	118.30
1	A	277	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	A	129	ARG	NE-CZ-NH1	5.14	122.87	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3239	0	3260	55	0
2	A	8	0	0	0	0
3	A	2	0	0	0	0
4	A	5	0	0	0	0
5	A	1	0	0	0	0
6	A	123	0	0	4	0
All	All	3378	0	3260	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:HIS:HE1	1:A:160:LYS:H	1.17	0.91
1:A:203:MSE:HE3	1:A:230:LEU:HD13	1.54	0.89
1:A:262:ASP:O	6:A:2082:HOH:O	1.97	0.83
1:A:271:ASN:HD21	1:A:299:ASN:HD21	1.25	0.81
1:A:252:ASN:H	1:A:272:GLN:HE21	1.27	0.80
1:A:16:ILE:HD12	1:A:17:ILE:N	2.05	0.71
1:A:191:HIS:HD1	1:A:206:ARG:HB3	1.57	0.68
1:A:132:ARG:NE	1:A:191:HIS:HD2	1.93	0.67
1:A:351:PRO:O	1:A:354:LYS:HG3	1.97	0.65
1:A:132:ARG:NE	1:A:191:HIS:CD2	2.65	0.65
1:A:39:LEU:HD23	1:A:62:ALA:HB3	1.81	0.63
1:A:191:HIS:ND1	1:A:206:ARG:HB3	2.13	0.62
1:A:132:ARG:CZ	1:A:191:HIS:CD2	2.83	0.61
1:A:132:ARG:CD	1:A:191:HIS:HD2	2.14	0.60
1:A:191:HIS:CE1	1:A:206:ARG:CB	2.85	0.60
1:A:80:ARG:HD2	1:A:125:TRP:CE2	2.38	0.58
1:A:111:LEU:HD23	1:A:391:PRO:HB2	1.88	0.56
1:A:245:GLU:HG3	1:A:246:MSE:N	2.20	0.55
1:A:83:HIS:CE1	1:A:160:LYS:H	2.09	0.55
1:A:246:MSE:HA	1:A:246:MSE:HE2	1.89	0.55
1:A:140:LYS:HB2	6:A:2044:HOH:O	2.08	0.54
1:A:112:MSE:HE1	1:A:398:SER:OG	2.10	0.52
1:A:369:ALA:HB3	1:A:373:MSE:HE3	1.92	0.50
1:A:44:LEU:HD11	1:A:50:ALA:HB3	1.94	0.50
1:A:191:HIS:CE1	1:A:206:ARG:HB2	2.46	0.50
1:A:34:HIS:HD2	1:A:35:ASN:HB2	1.77	0.49
1:A:21:VAL:HG21	1:A:44:LEU:HB3	1.95	0.49
1:A:366:ARG:HD2	1:A:390:ASN:ND2	2.27	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:411:LEU:CD1	1:A:427:VAL:HG22	2.43	0.49
1:A:39:LEU:CD2	1:A:62:ALA:HB3	2.43	0.48
1:A:187:ARG:HG3	1:A:188:HIS:CE1	2.49	0.47
1:A:191:HIS:ND1	1:A:206:ARG:CB	2.77	0.47
1:A:132:ARG:CD	1:A:191:HIS:CD2	2.98	0.46
1:A:354:LYS:HD2	1:A:354:LYS:C	2.36	0.46
1:A:100:GLN:HE22	1:A:419:HIS:CD2	2.33	0.46
1:A:124:PRO:O	1:A:419:HIS:HE1	1.99	0.46
1:A:411:LEU:HD13	1:A:427:VAL:HG22	1.97	0.45
1:A:92:GLN:NE2	1:A:100:GLN:HG2	2.33	0.44
1:A:187:ARG:CG	1:A:188:HIS:CE1	3.01	0.43
1:A:413:MSE:HE3	1:A:423:LEU:HD13	1.98	0.43
1:A:354:LYS:HD2	1:A:355:ASN:N	2.33	0.43
1:A:325:GLN:HG2	6:A:2100:HOH:O	2.19	0.43
1:A:373:MSE:HE2	1:A:373:MSE:N	2.34	0.43
1:A:165:LEU:HG	1:A:169:LEU:HB2	2.02	0.42
1:A:217:GLU:OE1	1:A:221:ARG:HD2	2.19	0.42
1:A:271:ASN:HD21	1:A:299:ASN:ND2	2.05	0.42
1:A:369:ALA:HB1	1:A:372:VAL:HB	2.02	0.42
1:A:354:LYS:HZ3	1:A:355:ASN:HB2	1.84	0.42
1:A:281:TRP:CZ3	1:A:426:MSE:HG2	2.54	0.41
1:A:191:HIS:CE1	1:A:206:ARG:HG2	2.55	0.41
1:A:207:HIS:HD2	1:A:209:ALA:O	2.03	0.41
1:A:302:LEU:HB2	1:A:303:PRO:HD3	2.02	0.41
1:A:16:ILE:HD12	1:A:16:ILE:C	2.42	0.40
1:A:175:LYS:NZ	6:A:2058:HOH:O	2.54	0.40
1:A:422:HIS:HB3	1:A:424:GLU:OE1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	416/433 (96%)	405 (97%)	8 (2%)	3 (1%)	19 10

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	ILE
1	A	59	LYS
1	A	431	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	351/353 (99%)	319 (91%)	32 (9%)	7 2

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ILE
1	A	24	LEU
1	A	27	PHE
1	A	37	LYS
1	A	47	GLN
1	A	53	THR
1	A	54	VAL
1	A	55	THR
1	A	58	LYS
1	A	76	ARG
1	A	102	ARG
1	A	111	LEU
1	A	129	ARG
1	A	142	GLN
1	A	149	ARG
1	A	187	ARG
1	A	217	GLU
1	A	237	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	245	GLU
1	A	261	ARG
1	A	330	LEU
1	A	345	GLU
1	A	349	LYS
1	A	354	LYS
1	A	366	ARG
1	A	378	LYS
1	A	380	GLU
1	A	382	ILE
1	A	386	TYR
1	A	413	MSE
1	A	423	LEU
1	A	431	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	HIS
1	A	49	ASN
1	A	83	HIS
1	A	92	GLN
1	A	136	ASN
1	A	188	HIS
1	A	207	HIS
1	A	252	ASN
1	A	271	ASN
1	A	272	GLN
1	A	419	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PO4	A	1435	-	4,4,4	1.11	0	6,6,6	0.40	0
2	SF4	A	1432	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	A	1432	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.