



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 28, 2023 – 10:31 AM EST

PDB ID : 8UZ7
Title : Crystal structure of a novel triose phosphate isomerase identified on the shrimp transcriptome
Authors : Sotelo-Mundo, R.R.; Gomez-Yanes, A.C.; Lopez-Zavala, A.A.; Lopez-Garcia, J.A.; Ochoa-Leyva, A.
Deposited on : 2023-11-14
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	93% (green), 5% (yellow), 2% (orange), 0% (red)
1	B	249	85% (green), 13% (yellow), 2% (orange), 2% (red)
1	C	249	87% (green), 11% (yellow), 2% (orange), 2% (red)
1	D	249	90% (green), 8% (yellow), 2% (orange), 0% (red)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	D	302	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8125 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Triosephosphate isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	244	1895	1190	341	350	14	0	1	0
1	B	244	1890	1184	341	351	14	0	1	0
1	C	244	1896	1188	341	353	14	0	2	0
1	D	244	1887	1182	341	350	14	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	200	ARG	LYS	conflict	UNP A0A3R7QX74
A	249	GLY	-	expression tag	UNP A0A3R7QX74
B	200	ARG	LYS	conflict	UNP A0A3R7QX74
B	249	GLY	-	expression tag	UNP A0A3R7QX74
C	200	ARG	LYS	conflict	UNP A0A3R7QX74
C	249	GLY	-	expression tag	UNP A0A3R7QX74
D	200	ARG	LYS	conflict	UNP A0A3R7QX74
D	249	GLY	-	expression tag	UNP A0A3R7QX74

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Cl 2	0	0
2	C	2	Total 2	Cl 2	0	0
2	D	2	Total 2	Cl 2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	162	Total	O	0	0
			162	162		
4	B	115	Total	O	0	0
			115	115		
4	C	125	Total	O	0	0
			125	125		
4	D	148	Total	O	0	0
			148	148		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Triosephosphate isomerase

Chain A:  93% 5% .





- Molecule 1: Triosephosphate isomerase

Chain B:  85% 13% .




- Molecule 1: Triosephosphate isomerase

Chain C:  87% 11% .



- Molecule 1: Triosephosphate isomerase

Chain D:  90% 8% .



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.56Å 84.25Å 73.74Å 90.00° 92.69° 90.00°	Depositor
Resolution (Å)	55.45 – 1.70 55.45 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.1 (55.45-1.70) 93.1 (55.45-1.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 1.70Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.189 , 0.191 0.189 , 0.189	Depositor DCC
R_{free} test set	2025 reflections (2.21%)	wwPDB-VP
Wilson B-factor (Å ²)	19.9	Xtrriage
Anisotropy	0.025	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 53.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.004 for l,k,-h 0.037 for h,-k,-l 0.026 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8125	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4720e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1931	0.55	0/2608
1	B	0.35	0/1925	0.55	0/2600
1	C	0.38	0/1934	0.56	0/2612
1	D	0.36	0/1919	0.54	0/2592
All	All	0.37	0/7709	0.55	0/10412

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1895	0	1901	9	0
1	B	1890	0	1897	19	0
1	C	1896	0	1903	16	0
1	D	1887	0	1892	13	0
2	A	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	4	1
3	C	1	0	0	0	0
4	A	162	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	115	0	0	3	1
4	C	125	0	0	3	0
4	D	148	0	0	5	0
All	All	8125	0	7593	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:195:ASN:ND2	2:D:302:CL:CL	2.17	1.11
2:D:302:CL:CL	4:D:424:HOH:O	2.05	1.10
2:D:302:CL:CL	4:D:507:HOH:O	2.18	0.98
1:A:5:ARG:N	4:A:401:HOH:O	1.97	0.97
1:B:5:ARG:N	4:B:302:HOH:O	2.05	0.89
2:D:302:CL:CL	4:D:428:HOH:O	2.30	0.86
1:C:219:ARG:O	4:C:401:HOH:O	2.06	0.74
1:B:139:ARG:NH1	4:B:303:HOH:O	2.21	0.74
1:A:18:ASP:OD2	4:A:402:HOH:O	2.10	0.69
1:B:200:ARG:NH1	4:B:301:HOH:O	2.00	0.67
1:A:146:LYS:HD2	4:A:548:HOH:O	2.01	0.61
1:C:155:ILE:O	1:C:156:ARG:NH1	2.34	0.60
1:C:130:GLU:OE2	1:C:135:ARG:HD3	2.02	0.59
1:B:132:THR:HA	1:B:169:TRP:HB3	1.85	0.59
1:C:179:ASN:HB3	1:C:224:LEU:HD21	1.84	0.58
1:D:200:ARG:NH2	4:D:405:HOH:O	2.36	0.58
1:C:8:PHE:O	1:C:229:GLY:HA3	2.05	0.56
1:B:136:LYS:O	1:B:137:GLU:HB2	2.05	0.56
1:C:130:GLU:OE2	1:C:140:THR:OG1	2.15	0.56
1:D:8:PHE:O	1:D:229:GLY:HA3	2.08	0.53
1:C:215:SER:OG	1:C:243:ASP:OD2	2.25	0.53
1:C:133:GLU:H	1:C:133:GLU:CD	2.12	0.52
1:C:97:PRO:HG3	1:C:129:CYS:HB3	1.92	0.52
1:C:179:ASN:ND2	1:C:220:GLU:OE2	2.38	0.52
1:B:167:ALA:HB1	1:B:169:TRP:CE2	2.47	0.50
1:C:21:ARG:NH2	4:C:403:HOH:O	2.41	0.49
1:D:232:VAL:HG11	1:D:235:ALA:HB3	1.94	0.49
1:D:232:VAL:CG1	1:D:235:ALA:HB3	2.41	0.49
1:B:179:ASN:ND2	1:B:220:GLU:OE2	2.46	0.48
1:B:157:ASP:OD2	1:B:160:ARG:NE	2.44	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:26:VAL:HG22	1:D:56:LEU:HD13	1.96	0.48
1:B:217:ASN:O	1:B:221:MET:HG3	2.14	0.47
1:C:26:VAL:HG22	1:C:56:LEU:HD13	1.96	0.47
1:A:8:PHE:O	1:A:229:GLY:HA3	2.15	0.47
1:D:108:SER:O	1:D:112:GLU:HG3	2.14	0.47
1:B:85:LYS:NZ	1:B:90:ASP:OD1	2.48	0.47
1:A:6:LYS:HD3	1:A:39:GLU:HB2	1.95	0.46
1:C:85:LYS:HE2	4:C:433:HOH:O	2.16	0.46
1:B:8:PHE:O	1:B:229:GLY:HA3	2.16	0.45
1:B:81:PRO:HB3	1:B:122:MET:HG3	1.98	0.45
1:B:130:GLU:HG2	1:B:169:TRP:CD1	2.51	0.45
1:C:132:THR:HA	1:C:169:TRP:HB3	1.97	0.45
1:D:81:PRO:HB3	1:D:122:MET:HG3	1.98	0.45
1:C:68:TYR:HB2	1:C:78:GLU:HB3	1.99	0.45
1:C:34:LEU:HD22	1:C:245:ILE:HG21	1.98	0.45
1:D:238:LYS:HB2	1:D:238:LYS:HE3	1.85	0.44
1:B:238:LYS:HB2	1:B:238:LYS:HE2	1.66	0.43
1:D:167:ALA:HB1	1:D:169:TRP:CE2	2.54	0.43
1:D:188:LEU:HD22	4:D:468:HOH:O	2.18	0.43
1:B:26:VAL:HG22	1:B:56:LEU:HG	2.01	0.43
1:B:135:ARG:NH2	1:B:181:GLN:HG2	2.33	0.42
1:B:140:THR:O	1:B:144:LEU:HB2	2.19	0.42
1:A:72:ARG:HD2	1:A:72:ARG:C	2.41	0.41
1:B:130:GLU:HG2	1:B:169:TRP:HD1	1.86	0.41
1:B:97:PRO:HG3	1:B:129:CYS:HB3	2.02	0.41
1:D:130:GLU:OE2	1:D:135:ARG:NH2	2.43	0.41
1:A:72:ARG:NH2	1:D:70:VAL:HA	2.35	0.41
1:A:213:VAL:CG1	1:A:232:VAL:HG12	2.51	0.40
1:A:214:ASN:HB2	1:A:240:ASP:OD2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:302:CL:CL	4:B:322:HOH:O[1_655]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/249 (98%)	236 (97%)	7 (3%)	0	100	100
1	B	243/249 (98%)	234 (96%)	9 (4%)	0	100	100
1	C	244/249 (98%)	238 (98%)	6 (2%)	0	100	100
1	D	242/249 (97%)	235 (97%)	7 (3%)	0	100	100
All	All	972/996 (98%)	943 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/205 (98%)	200 (99%)	2 (1%)	76	67
1	B	202/205 (98%)	200 (99%)	2 (1%)	76	67
1	C	203/205 (99%)	202 (100%)	1 (0%)	88	83
1	D	201/205 (98%)	200 (100%)	1 (0%)	88	83
All	All	808/820 (98%)	802 (99%)	6 (1%)	84	77

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	ARG
1	A	206	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	188	LEU
1	B	214	ASN
1	C	206	ARG
1	D	206	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	244/249 (97%)	-0.33	0 100 100	12, 19, 29, 44	0
1	B	244/249 (97%)	-0.14	3 (1%) 79 82	13, 23, 39, 57	0
1	C	244/249 (97%)	-0.17	6 (2%) 57 61	13, 21, 38, 48	0
1	D	244/249 (97%)	-0.35	0 100 100	13, 21, 31, 35	0
All	All	976/996 (97%)	-0.25	9 (0%) 84 87	12, 21, 36, 57	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	176	PHE	5.0
1	C	172	ASN	4.4
1	C	174	GLY	3.1
1	B	176	PHE	3.0
1	B	133	GLU	2.9
1	C	170	ALA	2.7
1	C	140	THR	2.4
1	C	175	VAL	2.3
1	B	138	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	D	302	1/1	0.96	0.04	17,17,17,17	1
2	CL	C	302	1/1	0.97	0.07	21,21,21,21	1
2	CL	D	301	1/1	0.98	0.06	20,20,20,20	0
3	MG	C	303	1/1	0.98	0.05	16,16,16,16	0
2	CL	A	302	1/1	0.99	0.05	18,18,18,18	1
2	CL	C	301	1/1	0.99	0.06	18,18,18,18	0
2	CL	A	301	1/1	1.00	0.06	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.