



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 22, 2024 – 06:04 PM EST

PDB ID : 8V04
Title : High resolution TMPRSS2 structure following acylation by nafamostat
Authors : Fraser, B.J.; Dong, A.; Kutera, M.; Seitova, A.; Li, Y.; Hutchinson, A.; Edwards, A.; Benard, F.; Levon, H.; Arrowsmith, C.
Deposited on : 2023-11-16
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
Sidechain outliers	138945	5703 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	 14% 75% 17% • 7%
2	B	249	 % 85% 11% •
3	C	4	 50% 50%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 3228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transmembrane protease serine 2 non-catalytic chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	102	830	522	147	149	12	0	9	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	146	ALA	-	expression tag	UNP O15393
A	147	ALA	-	expression tag	UNP O15393
A	251	ASP	SER	engineered mutation	UNP O15393
A	252	ASP	ARG	engineered mutation	UNP O15393
A	253	ASP	GLN	engineered mutation	UNP O15393
A	254	ASP	SER	engineered mutation	UNP O15393
A	255	LYS	ARG	engineered mutation	UNP O15393

- Molecule 2 is a protein called Transmembrane protease serine 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	238	1949	1262	320	346	21	0	25	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	493	GLU	-	expression tag	UNP O15393
B	494	PHE	-	expression tag	UNP O15393
B	495	VAL	-	expression tag	UNP O15393
B	496	GLU	-	expression tag	UNP O15393
B	497	HIS	-	expression tag	UNP O15393
B	498	HIS	-	expression tag	UNP O15393
B	499	HIS	-	expression tag	UNP O15393
B	500	HIS	-	expression tag	UNP O15393
B	501	HIS	-	expression tag	UNP O15393

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	502	HIS	-	expression tag	UNP O15393
B	503	HIS	-	expression tag	UNP O15393
B	504	HIS	-	expression tag	UNP O15393

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	4	52	30	3	19	0	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	4	2	2	0	0
4	A	1	4	2	2	0	0
4	A	1	4	2	2	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	X	0	0
			1	1		

- Molecule 6 is 4-carbamimidamidobenzoic acid (three-letter code: GBS) (formula: C₈H₉N₃O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			12	8	3	1		

- Molecule 7 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 13 6 7	0	0

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total Ca 1 1	0	0

- Molecule 9 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	75	Total	O	0	0
			75	75		
10	B	245	Total	O	0	1
			246	246		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transmembrane protease serine 2 non-catalytic chain

Chain A: 



- Molecule 2: Transmembrane protease serine 2

Chain B: 



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.76Å 50.50Å 64.58Å 90.00° 91.60° 90.00°	Depositor
Resolution (Å)	39.81 – 1.58 39.78 – 1.58	Depositor EDS
% Data completeness (in resolution range)	99.0 (39.81-1.58) 99.0 (39.78-1.58)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 1.58Å)	Xtrriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.156 , 0.181 0.168 , 0.188	Depositor DCC
R_{free} test set	2463 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtrriage
Anisotropy	1.111	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.028 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3228	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GBS, EDO, FUC, NAG, CIT, UNX, CA, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/870	0.82	0/1172
2	B	0.54	0/2076	0.86	1/2829 (0.0%)
All	All	0.52	0/2946	0.85	1/4001 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	316	ARG	NE-CZ-NH2	-5.34	117.63	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	150[A]	ARG	Sidechain
1	A	150[B]	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	830	0	803	21	0
2	B	1949	0	1938	20	0
3	C	52	0	46	2	0
4	A	12	0	18	4	0
4	B	24	0	36	6	0
5	A	1	0	0	0	0
6	B	12	0	0	0	0
7	B	13	0	5	1	0
8	B	1	0	0	0	0
9	B	13	0	18	3	0
10	A	75	0	0	1	0
10	B	246	0	0	4	0
All	All	3228	0	2864	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:360:LEU:HD13	4:B:603:EDO:H21	1.56	0.85
2:B:407:THR:HA	2:B:410[B]:CYS:SG	2.24	0.78
2:B:438[B]:GLN:NE2	10:B:702:HOH:O	2.21	0.73
1:A:150[B]:ARG:HG2	1:A:168:TRP:CZ3	2.28	0.68
4:B:605:EDO:H11	10:B:742:HOH:O	1.94	0.66
2:B:409:ARG:HD3	2:B:430:LEU:HD11	1.77	0.66
1:A:172[A]:CYS:SG	1:A:174:ASP:OD1	2.55	0.65
1:A:213:ASN:HD22	4:A:304:EDO:C2	2.11	0.64
1:A:172[A]:CYS:SG	1:A:174:ASP:CG	2.76	0.64
2:B:409:ARG:HD3	2:B:430:LEU:CD1	2.29	0.62
1:A:209:PHE:CG	1:A:231[B]:CYS:SG	2.93	0.61
1:A:150[B]:ARG:HH21	1:A:150[B]:ARG:HB3	1.65	0.61
4:A:304:EDO:H22	3:C:1:NAG:H2	1.84	0.59
1:A:209:PHE:CD1	1:A:231[B]:CYS:SG	2.97	0.58
2:B:341:THR:O	9:B:606:PG4:H51	2.03	0.58
2:B:481[A]:THR:HG22	2:B:485:TYR:CE2	2.40	0.57
1:A:191:LYS:NZ	2:B:491:ASP:O	2.37	0.57
2:B:419:LEU:O	9:B:606:PG4:H42	2.06	0.55
1:A:150[B]:ARG:HH21	1:A:150[B]:ARG:CB	2.19	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:ASN:HD22	4:A:304:EDO:H22	1.74	0.53
2:B:368:ASN:O	2:B:371:MET:HG2	2.08	0.52
2:B:281[B]:CYS:HB2	2:B:441:SER:O	2.10	0.52
1:A:165:ARG:HH11	1:A:165:ARG:HG3	1.75	0.51
1:A:176:TRP:CH2	1:A:221:ILE:HD11	2.46	0.51
2:B:352[B]:GLN:CD	10:B:703:HOH:O	2.50	0.50
1:A:169:HIS:CE1	1:A:208:SER:HB3	2.46	0.50
2:B:264:PRO:HB3	4:B:603:EDO:H12	1.96	0.48
2:B:367:PRO:O	2:B:481[B]:THR:HG21	2.13	0.48
1:A:191:LYS:HE3	10:B:840:HOH:O	2.14	0.47
1:A:166[B]:LYS:HE2	10:A:429:HOH:O	2.15	0.47
1:A:180:TYR:CE2	1:A:212:LEU:HD21	2.50	0.47
2:B:410[B]:CYS:CB	2:B:426:CYS:SG	3.04	0.46
1:A:188[B]:MET:HA	4:B:604:EDO:H22	1.96	0.46
1:A:172[B]:CYS:SG	1:A:209:PHE:HB3	2.56	0.45
1:A:165:ARG:HG3	1:A:165:ARG:NH1	2.31	0.45
7:B:601:CIT:O2	7:B:601:CIT:O5	2.35	0.44
2:B:329:GLU:HB2	2:B:352[B]:GLN:HG3	1.99	0.43
1:A:188[A]:MET:HA	4:B:604:EDO:H22	1.99	0.43
2:B:486:ARG:NE	2:B:489:ARG:HH11	2.15	0.43
2:B:457:GLY:HA2	2:B:475:GLY:O	2.19	0.43
1:A:148[B]:CYS:HB2	1:A:161:TYR:HB3	2.00	0.41
4:A:301:EDO:H22	3:C:1:NAG:H83	2.02	0.41
2:B:341:THR:O	9:B:606:PG4:C5	2.67	0.41
2:B:331:VAL:O	4:B:608:EDO:H12	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	107/110 (97%)	104 (97%)	3 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	261/249 (105%)	256 (98%)	4 (2%)	1 (0%)	34	15
All	All	368/359 (102%)	360 (98%)	7 (2%)	1 (0%)	41	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	415	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	93/95 (98%)	92 (99%)	1 (1%)	73	55
2	B	216/210 (103%)	216 (100%)	0	100	100
All	All	309/305 (101%)	308 (100%)	1 (0%)	91	86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	0.50	0	17,19,21	0.57	0
3	NAG	C	2	3	14,14,15	0.44	0	17,19,21	0.57	0
3	NAG	C	3	3	14,14,15	0.44	0	17,19,21	1.42	1 (5%)
3	FUC	C	4	3	10,10,11	0.37	0	14,14,16	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	NAG	C	3	3	-	4/6/23/26	0/1/1/1
3	FUC	C	4	3	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3	NAG	O5-C1-C2	4.69	118.69	111.29

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	3	NAG	C4-C5-C6-O6
3	C	3	NAG	C8-C7-N2-C2
3	C	3	NAG	O7-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	C	3	NAG	O5-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 1 is unknown and 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	B	603	-	3,3,3	0.42	0	2,2,2	0.45	0
4	EDO	A	301	-	3,3,3	0.27	0	2,2,2	0.89	0
4	EDO	A	303	-	3,3,3	0.06	0	2,2,2	0.34	0
4	EDO	B	607	-	3,3,3	0.11	0	2,2,2	0.06	0
6	GBS	B	600	2	12,12,13	0.77	0	15,15,17	0.90	0
7	CIT	B	601	-	12,12,12	1.25	1 (8%)	17,17,17	1.11	2 (11%)
4	EDO	B	608	-	3,3,3	0.07	0	2,2,2	0.20	0
9	PG4	B	606	-	12,12,12	0.34	0	11,11,11	0.37	0
4	EDO	B	604	-	3,3,3	0.28	0	2,2,2	0.38	0
4	EDO	B	605	-	3,3,3	0.42	0	2,2,2	0.69	0
4	EDO	A	304	-	3,3,3	0.33	0	2,2,2	0.25	0
4	EDO	B	609	-	3,3,3	0.08	0	2,2,2	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	603	-	-	1/1/1/1	-
4	EDO	A	301	-	-	1/1/1/1	-
4	EDO	A	303	-	-	1/1/1/1	-
4	EDO	B	607	-	-	1/1/1/1	-
6	GBS	B	600	2	-	2/6/6/8	0/1/1/1
7	CIT	B	601	-	-	5/16/16/16	-
4	EDO	B	608	-	-	0/1/1/1	-
9	PG4	B	606	-	-	5/10/10/10	-
4	EDO	B	604	-	-	1/1/1/1	-
4	EDO	B	605	-	-	1/1/1/1	-
4	EDO	A	304	-	-	1/1/1/1	-
4	EDO	B	609	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	601	CIT	C3-C6	2.43	1.55	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	601	CIT	O3-C5-C4	-2.17	116.61	122.94
7	B	601	CIT	O5-C6-C3	-2.11	119.27	122.25

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	601	CIT	C2-C3-C6-O5
7	B	601	CIT	C2-C3-C6-O6
7	B	601	CIT	O7-C3-C6-O5
7	B	601	CIT	O7-C3-C6-O6
9	B	606	PG4	O3-C5-C6-O4
9	B	606	PG4	C5-C6-O4-C7
9	B	606	PG4	O4-C7-C8-O5
4	A	303	EDO	O1-C1-C2-O2
4	B	604	EDO	O1-C1-C2-O2
9	B	606	PG4	O2-C3-C4-O3
6	B	600	GBS	C4-C5-C6-O
4	B	607	EDO	O1-C1-C2-O2
6	B	600	GBS	C-C5-C6-O
4	B	603	EDO	O1-C1-C2-O2
4	B	605	EDO	O1-C1-C2-O2
9	B	606	PG4	C8-C7-O4-C6
7	B	601	CIT	O7-C3-C4-C5
4	A	304	EDO	O1-C1-C2-O2
4	A	301	EDO	O1-C1-C2-O2

There are no ring outliers.

8 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	603	EDO	2	0
4	A	301	EDO	1	0
7	B	601	CIT	1	0
4	B	608	EDO	1	0
9	B	606	PG4	3	0
4	B	604	EDO	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	605	EDO	1	0
4	A	304	EDO	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	102/110 (92%)	0.54	15 (14%) 2 2	13, 26, 54, 75	0
2	B	238/249 (95%)	-0.15	3 (1%) 77 78	10, 16, 29, 59	0
All	All	340/359 (94%)	0.06	18 (5%) 26 26	10, 18, 43, 75	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	218	ASN	5.1
2	B	414	TYR	5.0
1	A	201	VAL	4.4
1	A	207	THR	4.2
1	A	206	SER	3.8
2	B	492	GLY	3.5
1	A	249	ASN	3.4
1	A	148[A]	CYS	3.3
1	A	222	TYR	3.3
1	A	203	ASP	3.2
2	B	322	TYR	3.2
1	A	147	ALA	3.0
1	A	209	PHE	2.9
1	A	231[A]	CYS	2.4
1	A	149	VAL	2.3
1	A	202	ASP	2.3
1	A	219	VAL	2.2
1	A	250	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	3	14/15	0.71	0.31	55,62,81,81	0
3	FUC	C	4	10/11	0.78	0.37	30,30,30,30	0
3	NAG	C	1	14/15	0.89	0.09	22,26,31,32	0
3	NAG	C	2	14/15	0.92	0.13	35,43,53,53	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	A	304	4/4	0.48	0.38	42,47,48,52	0
4	EDO	B	607	4/4	0.73	0.23	63,63,65,67	0
8	CA	B	602	1/1	0.76	0.07	83,83,83,83	0
9	PG4	B	606	13/13	0.76	0.32	33,53,59,59	0
5	UNX	A	302	1/1	0.78	0.17	32,32,32,32	0
4	EDO	B	608	4/4	0.79	0.29	52,54,56,60	0
4	EDO	B	605	4/4	0.80	0.25	41,44,44,45	0
4	EDO	B	603	4/4	0.82	0.17	34,37,39,39	0
4	EDO	A	303	4/4	0.84	0.16	42,48,48,49	0
4	EDO	B	604	4/4	0.90	0.20	46,49,51,53	0
4	EDO	A	301	4/4	0.90	0.14	34,39,39,41	0
4	EDO	B	609	4/4	0.90	0.14	58,63,63,67	0
7	CIT	B	601	13/13	0.94	0.09	21,24,35,40	0
6	GBS	B	600	12/13	0.97	0.06	15,16,17,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.