



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 24, 2024 – 03:07 PM EDT

PDB ID : 6V1V  
Title : VIP3B (VIP3B\_2160) adapted for crystallization  
Authors : Evdokimov, A.G.; Zheng, M.; Moshiri, F.; Haas, J.; Lowder, C.  
Deposited on : 2019-11-21  
Resolution : 3.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1467 (3.20-3.16)
Clashscore	141614	1599 (3.20-3.16)
Ramachandran outliers	138981	1574 (3.20-3.16)
Sidechain outliers	138945	1573 (3.20-3.16)
RSRZ outliers	127900	1423 (3.20-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	796	 2% 74% 20% 6%
1	B	796	 5% 73% 20% 6%
1	C	796	 3% 70% 22% 7%
1	D	796	 4% 69% 22% 8%

## 2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 23683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vegetative insecticidal protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	752	6002	3826	976	1185	15	0	0	0
1	B	750	5985	3816	972	1182	15	0	0	0
1	C	739	5874	3747	955	1157	15	0	0	0
1	D	731	5822	3718	946	1143	15	0	0	0

There are 112 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	initiating methionine	UNP A0A290WPI2
A	-8	GLY	-	expression tag	UNP A0A290WPI2
A	-7	SER	-	expression tag	UNP A0A290WPI2
A	-6	SER	-	expression tag	UNP A0A290WPI2
A	-5	HIS	-	expression tag	UNP A0A290WPI2
A	-4	HIS	-	expression tag	UNP A0A290WPI2
A	-3	HIS	-	expression tag	UNP A0A290WPI2
A	-2	HIS	-	expression tag	UNP A0A290WPI2
A	-1	HIS	-	expression tag	UNP A0A290WPI2
A	0	HIS	-	expression tag	UNP A0A290WPI2
A	1	HIS	-	expression tag	UNP A0A290WPI2
A	?	-	TRP	deletion	UNP A0A290WPI2
A	?	-	LYS	deletion	UNP A0A290WPI2
A	?	-	GLU	deletion	UNP A0A290WPI2
A	?	-	LYS	deletion	UNP A0A290WPI2
A	?	-	SER	deletion	UNP A0A290WPI2
A	?	-	CYS	deletion	UNP A0A290WPI2
A	?	-	GLU	deletion	UNP A0A290WPI2
A	?	-	GLU	deletion	UNP A0A290WPI2
A	466	SER	ASP	engineered mutation	UNP A0A290WPI2
A	514	ALA	GLU	engineered mutation	UNP A0A290WPI2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	515	ALA	LYS	engineered mutation	UNP A0A290WPI2
A	517	ALA	GLN	engineered mutation	UNP A0A290WPI2
A	518	ALA	LYS	engineered mutation	UNP A0A290WPI2
A	?	-	ASP	deletion	UNP A0A290WPI2
A	?	-	THR	deletion	UNP A0A290WPI2
A	?	-	ILE	deletion	UNP A0A290WPI2
A	?	-	LYS	deletion	UNP A0A290WPI2
B	-9	MET	-	initiating methionine	UNP A0A290WPI2
B	-8	GLY	-	expression tag	UNP A0A290WPI2
B	-7	SER	-	expression tag	UNP A0A290WPI2
B	-6	SER	-	expression tag	UNP A0A290WPI2
B	-5	HIS	-	expression tag	UNP A0A290WPI2
B	-4	HIS	-	expression tag	UNP A0A290WPI2
B	-3	HIS	-	expression tag	UNP A0A290WPI2
B	-2	HIS	-	expression tag	UNP A0A290WPI2
B	-1	HIS	-	expression tag	UNP A0A290WPI2
B	0	HIS	-	expression tag	UNP A0A290WPI2
B	1	HIS	-	expression tag	UNP A0A290WPI2
B	?	-	TRP	deletion	UNP A0A290WPI2
B	?	-	LYS	deletion	UNP A0A290WPI2
B	?	-	GLU	deletion	UNP A0A290WPI2
B	?	-	LYS	deletion	UNP A0A290WPI2
B	?	-	SER	deletion	UNP A0A290WPI2
B	?	-	CYS	deletion	UNP A0A290WPI2
B	?	-	GLU	deletion	UNP A0A290WPI2
B	?	-	GLU	deletion	UNP A0A290WPI2
B	466	SER	ASP	engineered mutation	UNP A0A290WPI2
B	514	ALA	GLU	engineered mutation	UNP A0A290WPI2
B	515	ALA	LYS	engineered mutation	UNP A0A290WPI2
B	517	ALA	GLN	engineered mutation	UNP A0A290WPI2
B	518	ALA	LYS	engineered mutation	UNP A0A290WPI2
B	?	-	ASP	deletion	UNP A0A290WPI2
B	?	-	THR	deletion	UNP A0A290WPI2
B	?	-	ILE	deletion	UNP A0A290WPI2
B	?	-	LYS	deletion	UNP A0A290WPI2
C	-9	MET	-	initiating methionine	UNP A0A290WPI2
C	-8	GLY	-	expression tag	UNP A0A290WPI2
C	-7	SER	-	expression tag	UNP A0A290WPI2
C	-6	SER	-	expression tag	UNP A0A290WPI2
C	-5	HIS	-	expression tag	UNP A0A290WPI2
C	-4	HIS	-	expression tag	UNP A0A290WPI2
C	-3	HIS	-	expression tag	UNP A0A290WPI2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	HIS	-	expression tag	UNP A0A290WPI2
C	-1	HIS	-	expression tag	UNP A0A290WPI2
C	0	HIS	-	expression tag	UNP A0A290WPI2
C	1	HIS	-	expression tag	UNP A0A290WPI2
C	?	-	TRP	deletion	UNP A0A290WPI2
C	?	-	LYS	deletion	UNP A0A290WPI2
C	?	-	GLU	deletion	UNP A0A290WPI2
C	?	-	LYS	deletion	UNP A0A290WPI2
C	?	-	SER	deletion	UNP A0A290WPI2
C	?	-	CYS	deletion	UNP A0A290WPI2
C	?	-	GLU	deletion	UNP A0A290WPI2
C	?	-	GLU	deletion	UNP A0A290WPI2
C	466	SER	ASP	engineered mutation	UNP A0A290WPI2
C	514	ALA	GLU	engineered mutation	UNP A0A290WPI2
C	515	ALA	LYS	engineered mutation	UNP A0A290WPI2
C	517	ALA	GLN	engineered mutation	UNP A0A290WPI2
C	518	ALA	LYS	engineered mutation	UNP A0A290WPI2
C	?	-	ASP	deletion	UNP A0A290WPI2
C	?	-	THR	deletion	UNP A0A290WPI2
C	?	-	ILE	deletion	UNP A0A290WPI2
C	?	-	LYS	deletion	UNP A0A290WPI2
D	-9	MET	-	initiating methionine	UNP A0A290WPI2
D	-8	GLY	-	expression tag	UNP A0A290WPI2
D	-7	SER	-	expression tag	UNP A0A290WPI2
D	-6	SER	-	expression tag	UNP A0A290WPI2
D	-5	HIS	-	expression tag	UNP A0A290WPI2
D	-4	HIS	-	expression tag	UNP A0A290WPI2
D	-3	HIS	-	expression tag	UNP A0A290WPI2
D	-2	HIS	-	expression tag	UNP A0A290WPI2
D	-1	HIS	-	expression tag	UNP A0A290WPI2
D	0	HIS	-	expression tag	UNP A0A290WPI2
D	1	HIS	-	expression tag	UNP A0A290WPI2
D	?	-	TRP	deletion	UNP A0A290WPI2
D	?	-	LYS	deletion	UNP A0A290WPI2
D	?	-	GLU	deletion	UNP A0A290WPI2
D	?	-	LYS	deletion	UNP A0A290WPI2
D	?	-	SER	deletion	UNP A0A290WPI2
D	?	-	CYS	deletion	UNP A0A290WPI2
D	?	-	GLU	deletion	UNP A0A290WPI2
D	?	-	GLU	deletion	UNP A0A290WPI2
D	466	SER	ASP	engineered mutation	UNP A0A290WPI2
D	514	ALA	GLU	engineered mutation	UNP A0A290WPI2

*Continued on next page...*


*Continued from previous page...*

<b>Chain</b>	<b>Residue</b>	<b>Modelled</b>	<b>Actual</b>	<b>Comment</b>	<b>Reference</b>
D	515	ALA	LYS	engineered mutation	UNP A0A290WPI2
D	517	ALA	GLN	engineered mutation	UNP A0A290WPI2
D	518	ALA	LYS	engineered mutation	UNP A0A290WPI2
D	?	-	ASP	deletion	UNP A0A290WPI2
D	?	-	THR	deletion	UNP A0A290WPI2
D	?	-	ILE	deletion	UNP A0A290WPI2
D	?	-	LYS	deletion	UNP A0A290WPI2









## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.49Å 106.54Å 117.73Å 96.13° 70.93° 69.52°	Depositor
Resolution (Å)	49.89 – 3.19 49.84 – 3.19	Depositor EDS
% Data completeness (in resolution range)	98.3 (49.89-3.19) 98.3 (49.84-3.19)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 3.19Å)	Xtrriage
Refinement program	REFMAC 5.8.0257	Depositor
R, $R_{free}$	0.218 , 0.259 0.215 , 0.254	Depositor DCC
$R_{free}$ test set	3538 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	102.1	Xtrriage
Anisotropy	0.053	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 58.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	23683	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	117.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.59	0/6105	0.79	0/8254
1	B	0.59	0/6087	0.80	0/8229
1	C	0.59	0/5976	0.80	0/8082
1	D	0.59	0/5919	0.81	1/7999 (0.0%)
All	All	0.59	0/24087	0.80	1/32564 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	228	ASP	CB-CA-C	5.03	120.45	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6002	0	5970	119	0
1	B	5985	0	5950	142	0
1	C	5874	0	5847	171	0
1	D	5822	0	5811	170	0
All	All	23683	0	23578	567	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 567 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:554:ILE:HG12	1:D:674:ILE:HG23	1.14	1.08
1:C:371:ASP:CB	1:C:372:PRO:HD3	1.83	1.07
1:C:371:ASP:HB3	1:C:372:PRO:HD3	1.09	1.04
1:B:652:PRO:HA	1:B:655:ILE:HD11	1.45	0.99
1:C:371:ASP:HB3	1:C:372:PRO:CD	1.91	0.99

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	740/796 (93%)	691 (93%)	48 (6%)	1 (0%)	51 83
1	B	736/796 (92%)	693 (94%)	43 (6%)	0	100 100
1	C	729/796 (92%)	682 (94%)	46 (6%)	1 (0%)	51 83
1	D	715/796 (90%)	673 (94%)	41 (6%)	1 (0%)	51 83
All	All	2920/3184 (92%)	2739 (94%)	178 (6%)	3 (0%)	51 83

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	371	ASP
1	D	229	MET
1	A	204	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	680/718 (95%)	670 (98%)	10 (2%)	65	85
1	B	678/718 (94%)	667 (98%)	11 (2%)	62	83
1	C	663/718 (92%)	650 (98%)	13 (2%)	55	79
1	D	658/718 (92%)	650 (99%)	8 (1%)	71	87
All	All	2679/2872 (93%)	2637 (98%)	42 (2%)	62	83

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	373	LEU
1	D	339	SER
1	C	391	ARG
1	C	613	ILE
1	D	498	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	670	ASN
1	D	647	ASN
1	C	102	ASN
1	D	74	ASN
1	C	101	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	752/796 (94%)	-0.02	16 (2%) 63 49	58, 106, 163, 202	0
1	B	750/796 (94%)	0.07	38 (5%) 28 15	59, 109, 183, 251	0
1	C	739/796 (92%)	-0.03	22 (2%) 50 34	63, 118, 172, 221	0
1	D	731/796 (91%)	0.13	32 (4%) 34 20	64, 118, 176, 230	0
All	All	2972/3184 (93%)	0.04	108 (3%) 42 27	58, 112, 174, 251	0

The worst 5 of 108 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	577	VAL	6.2
1	D	585	LEU	6.1
1	B	786	PHE	5.0
1	D	659	PHE	4.7
1	D	657	PHE	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.