



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 04:24 pm BST

PDB ID : 2V2X  
Title : T cell cross-reactivity and conformational changes during TCR engagement.  
Authors : Lee, J.K.; Stewart-Jones, G.; Dong, T.; harlos, K.; Di Gleria, K.; Dorrell, L.;  
Douek, D.C.; van der Merwe, P.A.; Jones, E.Y.; McMichael, A.J.  
Deposited on : 2007-06-07  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 5% 88% 11%
1	D	276	 3% 86% 13%
2	B	100	 % 93% 7%
2	E	100	 % 96% .
3	C	9	 11% 100%
3	F	9	 100%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	Total 2220	C 1387	N 404	O 420	S 9	0	0	1
1	D	276	Total 2220	C 1387	N 404	O 420	S 9	0	0	1

- Molecule 2 is a protein called BETA-2 MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total 817	C 520	N 140	O 153	S 4	0	0	0
2	E	100	Total 817	C 520	N 140	O 153	S 4	0	0	0

- Molecule 3 is a protein called HIV P17.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	9	Total 68	C 44	N 10	O 14	0	0	0
3	F	9	Total 68	C 44	N 10	O 14	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	428	Total 428	O 428	0	0
4	B	196	Total 196	O 196	0	0
4	C	18	Total 18	O 18	0	0
4	D	444	Total 444	O 444	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	E	215	Total 215	O 215	0	0
4	F	22	Total 22	O 22	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN




- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN



- Molecule 2: BETA-2 MICROGLOBULIN




- Molecule 2: BETA-2 MICROGLOBULIN



- Molecule 3: HIV P17

Chain C:  100%



- Molecule 3: HIV P17

Chain F:  100%

There are no outlier residues recorded for this chain.

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.31Å 63.61Å 74.74Å 81.83° 76.21° 77.73°	Depositor
Resolution (Å)	72.17 – 1.60 22.33 – 1.60	Depositor EDS
% Data completeness (in resolution range)	87.2 (72.17-1.60) 87.2 (22.33-1.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.27 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.192 , 0.238 0.190 , 0.237	Depositor DCC
$R_{free}$ test set	5039 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.3	Xtrriage
Anisotropy	0.365	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 56.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7533	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.59	1/2284 (0.0%)	0.69	0/3100
1	D	0.62	1/2284 (0.0%)	0.75	2/3100 (0.1%)
2	B	0.60	0/840	0.67	0/1135
2	E	0.65	0/840	0.69	0/1135
3	C	0.59	0/68	0.66	0/91
3	F	0.61	0/68	0.61	0/91
All	All	0.61	2/6384 (0.0%)	0.71	2/8652 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	275	GLU	C-N	-5.14	1.24	1.34
1	A	275	GLU	C-N	-5.01	1.24	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	6	ARG	NE-CZ-NH1	8.16	124.38	120.30
1	D	6	ARG	NE-CZ-NH2	-7.16	116.72	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2220	0	2063	25	0
1	D	2220	0	2063	37	0
2	B	817	0	776	6	0
2	E	817	0	776	4	0
3	C	68	0	72	0	0
3	F	68	0	72	0	0
4	A	428	0	0	7	0
4	B	196	0	0	3	0
4	C	18	0	0	0	0
4	D	444	0	0	10	0
4	E	215	0	0	3	0
4	F	22	0	0	0	0
All	All	7533	0	5822	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:GLU:HG3	4:A:2136:HOH:O	1.28	1.25
2:E:47:GLU:CA	4:E:2132:HOH:O	2.00	1.08
2:E:75:LYS:HG2	4:E:2167:HOH:O	1.77	0.85
1:D:6:ARG:HH11	1:D:6:ARG:HG2	1.46	0.80
1:D:263:HIS:CD2	1:D:265:GLY:H	2.00	0.78
1:A:263:HIS:CD2	1:A:265:GLY:H	2.02	0.78
1:A:255:GLN:HG2	4:A:2401:HOH:O	1.86	0.76
1:A:274:TRP:O	1:A:276:PRO:N	2.21	0.73
1:D:93:HIS:HD2	1:D:119:ASP:OD2	1.71	0.72
1:D:263:HIS:HD2	1:D:265:GLY:H	1.36	0.72
1:A:170:ARG:O	1:A:173:GLU:HG2	1.92	0.69
1:D:93:HIS:HE1	4:D:2196:HOH:O	1.76	0.69
1:A:72:GLN:HE22	1:A:75:ARG:HH11	1.40	0.69
1:D:97:ARG:HH21	1:D:114:HIS:HE1	1.41	0.69
1:A:263:HIS:HD2	1:A:265:GLY:H	1.39	0.68
1:A:93:HIS:HD2	1:A:119:ASP:OD2	1.77	0.68
1:D:81:LEU:HD13	1:D:118:TYR:CD1	2.28	0.68
1:A:97:ARG:HE	1:A:114:HIS:CE1	2.14	0.65
1:A:229:GLU:HG3	4:A:2377:HOH:O	1.99	0.63
1:A:97:ARG:HE	1:A:114:HIS:HE1	1.46	0.62
1:A:93:HIS:HE1	4:A:2186:HOH:O	1.83	0.60
1:D:111:ARG:HD2	1:D:113:TYR:OH	2.02	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:ARG:HG3	4:A:2029:HOH:O	2.03	0.58
1:D:192:HIS:CA	4:D:2338:HOH:O	2.51	0.58
1:D:6:ARG:HH11	1:D:6:ARG:CG	2.15	0.58
1:D:97:ARG:HE	1:D:114:HIS:CE1	2.21	0.57
1:D:186:LYS:HE3	4:D:2324:HOH:O	2.05	0.57
1:D:30:ASP:HB2	4:D:2351:HOH:O	2.05	0.56
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.40	0.56
1:A:253:GLN:NE2	1:A:256:ARG:HH11	2.04	0.55
1:A:19:GLU:HG3	1:A:20:PRO:HD2	1.88	0.55
2:E:75:LYS:HB3	4:E:2168:HOH:O	2.06	0.54
1:D:189:MET:HE2	1:D:274:TRP:HE3	1.71	0.53
2:B:3:ARG:HD3	4:B:2142:HOH:O	2.07	0.53
1:D:155:GLN:NE2	4:D:2291:HOH:O	2.43	0.52
1:A:255:GLN:HG3	4:A:2403:HOH:O	2.09	0.52
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.45	0.52
1:D:201:LEU:HD12	1:D:249:VAL:HG21	1.93	0.51
1:D:186:LYS:HD3	4:D:2326:HOH:O	2.11	0.51
1:D:97:ARG:HH21	1:D:114:HIS:CE1	2.23	0.51
1:A:58:GLU:CG	4:A:2136:HOH:O	2.12	0.51
1:D:121:LYS:HE3	4:D:2236:HOH:O	2.11	0.51
1:A:72:GLN:HE22	1:A:75:ARG:NH1	2.06	0.50
1:D:255:GLN:H	1:D:255:GLN:CD	2.15	0.49
2:B:4:THR:HG23	2:B:86:THR:OG1	2.13	0.48
1:D:131:ARG:HD3	4:D:2262:HOH:O	2.13	0.48
2:B:4:THR:HG22	2:B:5:PRO:HD2	1.96	0.47
1:D:6:ARG:HG2	1:D:6:ARG:NH1	2.16	0.47
2:B:77:GLU:HG2	4:B:2086:HOH:O	2.14	0.47
1:D:72:GLN:HG2	4:D:2049:HOH:O	2.14	0.47
1:A:210:PRO:O	1:A:263:HIS:HE1	1.98	0.47
1:A:275:GLU:OE2	1:A:275:GLU:N	2.48	0.47
1:D:97:ARG:HE	1:D:114:HIS:HE1	1.61	0.46
1:A:19:GLU:HG3	1:A:20:PRO:CD	2.46	0.46
1:D:93:HIS:CD2	1:D:119:ASP:OD2	2.61	0.46
1:A:111:ARG:HD3	1:A:113:TYR:OH	2.15	0.45
1:D:253:GLN:NE2	1:D:256:ARG:HH11	2.13	0.45
1:D:186:LYS:HD2	1:D:207:SER:HB3	1.98	0.45
1:D:189:MET:HE1	1:D:274:TRP:HB2	1.98	0.45
1:D:20:PRO:HG2	1:D:75:ARG:HG2	1.99	0.44
1:D:81:LEU:HD13	1:D:118:TYR:CG	2.52	0.44
1:D:82:ARG:HD2	1:D:88:SER:O	2.18	0.44
1:D:65:ARG:NH1	4:D:2151:HOH:O	2.50	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:3:ARG:NH2	4:B:2012:HOH:O	2.47	0.43
1:D:97:ARG:NH2	1:D:114:HIS:HE1	2.11	0.43
1:A:51:TRP:CZ2	1:A:179:LEU:HD11	2.54	0.43
1:D:253:GLN:HA	1:D:255:GLN:HE22	1.83	0.43
1:D:210:PRO:O	1:D:263:HIS:HE1	2.02	0.42
1:A:97:ARG:HH21	1:A:114:HIS:HE1	1.68	0.41
1:D:6:ARG:CG	1:D:6:ARG:NH1	2.80	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/276 (99%)	270 (98%)	3 (1%)	1 (0%)	34	15
1	D	274/276 (99%)	271 (99%)	3 (1%)	0	100	100
2	B	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	E	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	F	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	758/770 (98%)	747 (98%)	10 (1%)	1 (0%)	51	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	275	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/232 (97%)	221 (98%)	5 (2%)	52	27
1	D	226/232 (97%)	221 (98%)	5 (2%)	52	27
2	B	91/95 (96%)	90 (99%)	1 (1%)	73	57
2	E	91/95 (96%)	90 (99%)	1 (1%)	73	57
3	C	8/8 (100%)	8 (100%)	0	100	100
3	F	8/8 (100%)	8 (100%)	0	100	100
All	All	650/670 (97%)	638 (98%)	12 (2%)	59	36

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ARG
1	A	74	HIS
1	A	163	THR
1	A	194	VAL
1	A	268	LYS
2	B	70	PHE
1	D	6	ARG
1	D	35	ARG
1	D	74	HIS
1	D	110	LEU
1	D	268	LYS
2	E	70	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	74	HIS
1	A	93	HIS
1	A	114	HIS
1	A	174	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	253	GLN
1	A	263	HIS
1	D	32	GLN
1	D	74	HIS
1	D	93	HIS
1	D	114	HIS
1	D	155	GLN
1	D	174	ASN
1	D	180	GLN
1	D	253	GLN
1	D	263	HIS
2	E	2	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	276/276 (100%)	-0.01	13 (4%) 31 28	11, 20, 30, 39	0
1	D	276/276 (100%)	-0.08	8 (2%) 51 49	9, 18, 30, 37	0
2	B	100/100 (100%)	-0.13	1 (1%) 82 82	12, 20, 29, 32	0
2	E	100/100 (100%)	-0.32	1 (1%) 82 82	11, 16, 24, 29	0
3	C	9/9 (100%)	0.13	1 (11%) 5 4	14, 22, 24, 26	0
3	F	9/9 (100%)	-0.13	0 100 100	13, 18, 22, 23	0
All	All	770/770 (100%)	-0.09	24 (3%) 49 46	9, 19, 29, 39	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	276	PRO	11.2
1	D	90	ALA	4.0
1	D	16	GLY	3.7
2	B	48	LYS	3.6
1	D	86	ASN	3.0
1	A	194	VAL	2.8
1	A	225	THR	2.8
1	A	197	HIS	2.7
1	A	196	ASP	2.6
1	A	50	PRO	2.6
1	D	268	LYS	2.6
1	A	275	GLU	2.5
2	E	48	LYS	2.5
1	D	196	ASP	2.4
1	D	15	PRO	2.4
1	A	41	ALA	2.4
1	A	226	GLN	2.4
1	A	273	ARG	2.3
1	A	136	ALA	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	86	ASN	2.2
1	D	113	TYR	2.2
1	A	113	TYR	2.1
1	D	194	VAL	2.1
3	C	6	VAL	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.