



Full wwPDB X-ray Structure Validation Report i

Dec 3, 2024 – 04:05 PM EST

PDB ID : 8V5Q
Title : Varicella Zoster Virus (VZV) glycoprotein E (gE) gI binding domain in complex with human Fab 1E3
Authors : Holzapfel, G.; Seraj, N.; Harshbarger, W.
Deposited on : 2023-11-30
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	H	301	-	-	X	-

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 4451 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 1E3 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	220	Total	C 1666	N 1056	O 276	S 330	4	0	0

- Molecule 2 is a protein called Fab 1E3 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	211	Total	C 1622	N 1013	O 276	S 327	6	0	0


- Molecule 3 is a protein called Envelope glycoprotein E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	116	Total	C 948	N 612	O 154	S 181	1	0	0

There are 13 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	306	GLU	-	expression tag	UNP A6XEF7
G	307	ASN	-	expression tag	UNP A6XEF7
G	308	LEU	-	expression tag	UNP A6XEF7
G	309	TYR	-	expression tag	UNP A6XEF7
G	310	PHE	-	expression tag	UNP A6XEF7
G	311	GLN	-	expression tag	UNP A6XEF7
G	312	GLY	-	expression tag	UNP A6XEF7
G	313	HIS	-	expression tag	UNP A6XEF7
G	314	HIS	-	expression tag	UNP A6XEF7
G	315	HIS	-	expression tag	UNP A6XEF7
G	316	HIS	-	expression tag	UNP A6XEF7
G	317	HIS	-	expression tag	UNP A6XEF7
G	318	HIS	-	expression tag	UNP A6XEF7

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total C O 7 4 3	0	0
4	H	1	Total C O 7 4 3	0	0
4	G	1	Total C O 7 4 3	0	0
4	G	1	Total C O 7 4 3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	G	1	Total O S 5 4 1	0	0
5	G	1	Total O S 5 4 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	56	Total O 56 56	0	0
6	L	89	Total O 89 89	0	0
6	G	32	Total O 32 32	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	44.62 Å 193.75 Å 179.85 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.72 – 1.90 33.72 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (33.72-1.90) 89.2 (33.72-1.90)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	0.63 (at 1.91 Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R , R_{free}	0.226 , 0.233 0.227 , 0.234	Depositor DCC
R_{free} test set	59768 reflections (3.28%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtriage
Anisotropy	0.737	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4451	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.45	0/1708	0.70	0/2334
2	L	0.44	0/1656	0.66	0/2247
3	G	0.47	0/972	0.70	0/1325
All	All	0.45	0/4336	0.68	0/5906

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	210	ARG	Sidechain
2	L	93	ARG	Sidechain

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1666	0	1633	13	0
2	L	1622	0	1583	7	0
3	G	948	0	907	13	0
4	G	14	0	20	4	0
4	H	14	0	20	9	0
5	G	10	0	0	0	0
6	G	32	0	0	2	0
6	H	56	0	0	0	0
6	L	89	0	0	0	0
All	All	4451	0	4163	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:103:SER:HA	4:H:301:PEG:H31	1.55	0.85
1:H:104:SER:H	4:H:301:PEG:C3	1.99	0.76
1:H:103:SER:CA	4:H:301:PEG:H31	2.19	0.73
1:H:221:ARG:NH1	1:H:223:GLU:OE2	2.23	0.70
3:G:217:GLU:HG2	3:G:235:VAL:HB	1.74	0.69
1:H:204:THR:HG23	1:H:221:ARG:HE	1.59	0.67
1:H:104:SER:H	4:H:301:PEG:H31	1.63	0.61
3:G:165:ARG:HD3	3:G:221:ARG:HH21	1.65	0.60
1:H:104:SER:N	4:H:301:PEG:C3	2.69	0.56
1:H:104:SER:N	4:H:301:PEG:H31	2.21	0.55
2:L:37:GLN:HB2	2:L:47:LEU:HD11	1.91	0.53
3:G:165:ARG:CD	3:G:221:ARG:HH21	2.20	0.53
3:G:257:LYS:HE2	6:G:505:HOH:O	2.10	0.52
2:L:145:VAL:HG22	2:L:195:VAL:HG22	1.92	0.52
2:L:192:ALA:HB2	2:L:207:SER:HB3	1.90	0.51
2:L:28:GLY:H	4:G:404:PEG:C3	2.25	0.49
3:G:252:GLU:HB2	3:G:255:VAL:CG1	2.43	0.48
1:H:50:ILE:HG23	1:H:65:LEU:HD13	1.95	0.48
4:H:301:PEG:H32	6:G:504:HOH:O	2.15	0.47
3:G:245:GLU:OE1	4:G:404:PEG:H22	2.15	0.47
3:G:167:THR:HG22	3:G:219:SER:HB3	1.98	0.45
3:G:239:THR:O	3:G:243:GLU:HG2	2.16	0.45
2:L:55:GLN:HG3	2:L:56:ASN:N	2.33	0.43
4:H:301:PEG:H41	4:H:301:PEG:H21	1.69	0.43
3:G:164:VAL:HB	3:G:166:TYR:CE2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:18:LEU:HD12	1:H:119:VAL:HG11	2.01	0.42
1:H:60:TYR:CZ	3:G:151:PRO:HB2	2.55	0.42
3:G:272:ASN:HA	3:G:281:THR:O	2.20	0.42
4:H:302:PEG:H41	4:H:302:PEG:H22	1.79	0.41
2:L:93:ARG:O	2:L:94:THR:C	2.57	0.41
2:L:28:GLY:HA3	4:G:404:PEG:H11	2.03	0.41
3:G:214:GLN:O	3:G:238:SER:HB2	2.21	0.41
3:G:263:LYS:HE3	3:G:263:LYS:HB2	1.90	0.41
1:H:9:PRO:HD2	1:H:19:SER:O	2.21	0.41
1:H:59:THR:OG1	4:G:403:PEG:H42	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	216/245 (88%)	205 (95%)	11 (5%)	0	100 100
2	L	209/213 (98%)	200 (96%)	8 (4%)	1 (0%)	25 17
3	G	110/203 (54%)	108 (98%)	2 (2%)	0	100 100
All	All	535/661 (81%)	513 (96%)	21 (4%)	1 (0%)	44 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	210	ARG

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	H	191/213 (90%)	185 (97%)	6 (3%)	35 29
2	L	185/187 (99%)	184 (100%)	1 (0%)	86 88
3	G	103/182 (57%)	102 (99%)	1 (1%)	73 74
All	All	479/582 (82%)	471 (98%)	8 (2%)	56 54

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	5	GLN
1	H	13	LYS
1	H	18	LEU
1	H	64	SER
1	H	78	ILE
1	H	202	THR
2	L	175	SER
3	G	175	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	G	404	-	6,6,6	0.10	0	5,5,5	0.12	0
5	SO4	G	401	-	4,4,4	0.47	0	6,6,6	0.09	0
4	PEG	H	302	-	6,6,6	0.13	0	5,5,5	0.10	0
4	PEG	H	301	-	6,6,6	0.10	0	5,5,5	0.10	0
5	SO4	G	402	-	4,4,4	0.40	0	6,6,6	0.07	0
4	PEG	G	403	-	6,6,6	0.14	0	5,5,5	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	G	403	-	-	3/4/4/4	-
4	PEG	H	302	-	-	2/4/4/4	-
4	PEG	G	404	-	-	4/4/4/4	-
4	PEG	H	301	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	403	PEG	C1-C2-O2-C3
4	H	301	PEG	C4-C3-O2-C2
4	H	302	PEG	C4-C3-O2-C2
4	G	404	PEG	O1-C1-C2-O2
4	G	404	PEG	O2-C3-C4-O4
4	G	403	PEG	O1-C1-C2-O2
4	G	403	PEG	C4-C3-O2-C2
4	G	404	PEG	C1-C2-O2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	H	301	PEG	C1-C2-O2-C3
4	G	404	PEG	C4-C3-O2-C2
4	H	302	PEG	O2-C3-C4-O4

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	404	PEG	3	0
4	H	302	PEG	1	0
4	H	301	PEG	8	0
4	G	403	PEG	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	220/245 (89%)	1.01	29 (13%) 8 8	23, 43, 67, 84	0
2	L	211/213 (99%)	0.80	25 (11%) 10 11	22, 41, 70, 75	0
3	G	116/203 (57%)	0.93	18 (15%) 6 6	23, 39, 68, 95	0
All	All	547/661 (82%)	0.91	72 (13%) 8 8	22, 42, 68, 95	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	176	LEU	6.2
1	H	125	ALA	4.7
1	H	28	SER	4.3
1	H	126	SER	4.3
1	H	1	GLN	3.9
3	G	223	GLN	3.7
1	H	27	GLY	3.7
3	G	228	ALA	3.6
2	L	144	LYS	3.6
1	H	78	ILE	3.5
1	H	43	PRO	3.5
3	G	294	LYS	3.4
1	H	9	PRO	3.4
3	G	162	TYR	3.4
1	H	225	LYS	3.3
1	H	20	LEU	3.2
2	L	191	TYR	3.2
2	L	208	PHE	3.2
2	L	210	ARG	3.2
3	G	236	ASN	3.2
1	H	75	THR	3.1
1	H	138	SER	3.1
1	H	2	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	226	SER	3.0
3	G	175	SER	2.9
2	L	198	GLN	2.9
2	L	200	LEU	2.9
2	L	146	GLN	2.9
1	H	124	ALA	2.8
3	G	212	GLU	2.8
1	H	30	SER	2.7
3	G	230	GLN	2.7
2	L	149	VAL	2.7
1	H	202	THR	2.7
3	G	231	PRO	2.6
2	L	196	THR	2.6
2	L	193	CYS	2.6
1	H	44	GLY	2.6
3	G	234	VAL	2.5
1	H	29	ILE	2.5
2	L	195	VAL	2.5
3	G	213	ASP	2.5
3	G	233	ILE	2.5
1	H	32	ASN	2.5
3	G	235	VAL	2.4
2	L	192	ALA	2.4
1	H	16	GLN	2.4
2	L	178	LEU	2.3
2	L	187	LYS	2.3
2	L	152	ALA	2.3
2	L	153	LEU	2.3
2	L	190	VAL	2.3
1	H	148	ALA	2.2
3	G	232	TRP	2.2
1	H	169	ALA	2.2
1	H	10	GLY	2.2
2	L	157	ASN	2.2
2	L	188	HIS	2.2
2	L	131	VAL	2.2
2	L	183	ALA	2.2
2	L	204	VAL	2.1
1	H	145	GLY	2.1
2	L	113	SER	2.1
3	G	140	ARG	2.1
1	H	31	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	189	LYS	2.1
3	G	218	ILE	2.1
1	H	73	VAL	2.0
1	H	76	SER	2.0
3	G	276	SER	2.0
2	L	211	GLY	2.0
1	H	123	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	G	402	5/5	0.69	0.16	48,50,61,67	0
4	PEG	G	404	7/7	0.74	0.18	39,44,49,52	0
4	PEG	H	301	7/7	0.76	0.19	35,40,54,55	0
4	PEG	H	302	7/7	0.77	0.18	41,53,57,60	0
4	PEG	G	403	7/7	0.80	0.16	47,49,59,59	0
5	SO4	G	401	5/5	0.98	0.06	29,33,35,35	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.