



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 4, 2023 – 04:16 PM EDT

PDB ID : 3V8H  
Title : Crystal structure of Thymidylate Synthase from Burkholderia thailandensis  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2011-12-22  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	 4% 90% 5% 5%
1	B	327	 6% 92% 5% 5%
1	C	327	 4% 84% 12% 5%
1	D	327	 4% 84% 11% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	324	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	312	Total 2510	C 1610	N 445	O 448	S 7	0	7	0
1	B	311	Total 2463	C 1579	N 434	O 443	S 7	0	5	0
1	C	288	Total 2306	C 1473	N 412	O 415	S 6	0	5	0
1	D	290	Total 2305	C 1478	N 408	O 414	S 5	0	4	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q2SXY2
A	-2	PRO	-	expression tag	UNP Q2SXY2
A	-1	GLY	-	expression tag	UNP Q2SXY2
A	0	SER	-	expression tag	UNP Q2SXY2
B	-3	GLY	-	expression tag	UNP Q2SXY2
B	-2	PRO	-	expression tag	UNP Q2SXY2
B	-1	GLY	-	expression tag	UNP Q2SXY2
B	0	SER	-	expression tag	UNP Q2SXY2
C	-3	GLY	-	expression tag	UNP Q2SXY2
C	-2	PRO	-	expression tag	UNP Q2SXY2
C	-1	GLY	-	expression tag	UNP Q2SXY2
C	0	SER	-	expression tag	UNP Q2SXY2
D	-3	GLY	-	expression tag	UNP Q2SXY2
D	-2	PRO	-	expression tag	UNP Q2SXY2
D	-1	GLY	-	expression tag	UNP Q2SXY2
D	0	SER	-	expression tag	UNP Q2SXY2

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		
2	C	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	306	Total O 306 306	0	0
4	B	284	Total O 284 284	0	0
4	C	284	Total O 284 284	0	0
4	D	288	Total O 288 288	0	0







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.06Å 107.69Å 117.24Å 90.00° 98.06° 90.00°	Depositor
Resolution (Å)	49.51 – 1.65 19.84 – 1.65	Depositor EDS
% Data completeness (in resolution range)	95.4 (49.51-1.65) 95.4 (19.84-1.65)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.45 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.175 , 0.203 0.183 , 0.209	Depositor DCC
$R_{free}$ test set	8295 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.1	Xtrriage
Anisotropy	0.051	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10809	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	2/2593 (0.1%)	0.82	3/3525 (0.1%)
1	B	0.69	0/2540	0.77	2/3458 (0.1%)
1	C	0.69	0/2374	0.79	0/3226
1	D	0.70	1/2372 (0.0%)	0.84	2/3223 (0.1%)
All	All	0.70	3/9879 (0.0%)	0.80	7/13432 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	179	TRP	CD2-CE2	5.99	1.48	1.41
1	A	113	TRP	CD2-CE2	5.58	1.48	1.41
1	D	297	TRP	CD2-CE2	5.25	1.47	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	163	ASP	CB-CG-OD1	5.81	123.53	118.30
1	D	111	ARG	NE-CZ-NH1	5.80	123.20	120.30
1	B	163	ASP	CB-CG-OD1	5.62	123.36	118.30
1	B	73	ARG	NE-CZ-NH1	5.61	123.10	120.30
1	A	184	LEU	CB-CG-CD2	5.45	120.27	111.00
1	A	96	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	D	204	ARG	NE-CZ-NH1	5.18	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2510	0	2476	10	0
1	B	2463	0	2390	7	0
1	C	2306	0	2253	8	0
1	D	2305	0	2250	11	0
2	A	13	0	5	5	0
2	B	13	0	5	0	0
2	C	13	0	5	0	0
3	A	12	0	18	2	0
3	B	4	0	6	4	0
3	C	4	0	6	1	0
3	D	4	0	6	1	0
4	A	306	0	0	0	0
4	B	284	0	0	1	0
4	C	284	0	0	1	0
4	D	288	0	0	2	0
All	All	10809	0	9420	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192[B]:CYS:SG	1:B:215:SER:O	2.25	0.95
1:B:125:ALA:HB3	3:B:324:EDO:H22	1.64	0.78
1:C:20:ASN:HB2	1:C:22:THR:HG22	1.66	0.77
1:C:22:THR:HG23	1:C:24:ILE:H	1.52	0.75
1:B:125:ALA:HB3	3:B:324:EDO:C2	2.17	0.74
1:D:193:HIS:HB3	1:D:213:ILE:HD12	1.78	0.64
1:C:140[B]:ARG:NH1	4:C:893:HOH:O	2.27	0.58
1:D:193:HIS:HA	1:D:213:ILE:HD13	1.86	0.58
1:A:214:ARG:HH22	2:A:400:CIT:C5	2.18	0.56
2:A:400:CIT:O5	1:D:172:ARG:NH1	2.39	0.56
1:A:75:LEU:HD23	3:A:325:EDO:H21	1.89	0.55
2:A:400:CIT:C5	2:A:400:CIT:O6	2.51	0.54

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:ALA:HB3	3:B:324:EDO:H21	1.93	0.50
1:C:204[A]:ARG:NH2	3:C:324:EDO:O2	2.45	0.49
1:C:192[A]:CYS:SG	1:C:215:SER:O	2.71	0.48
1:D:193:HIS:HB3	1:D:213:ILE:CD1	2.42	0.48
3:D:324:EDO:H22	4:D:385:HOH:O	2.12	0.48
1:D:193:HIS:CB	1:D:213:ILE:CD1	2.93	0.46
1:A:250:PHE:CD1	1:D:250:PHE:CD1	3.04	0.46
1:C:279:GLU:OE2	1:C:309:VAL:HG21	2.15	0.45
1:D:193:HIS:HA	1:D:213:ILE:CD1	2.47	0.45
1:D:292:LYS:NZ	4:D:969:HOH:O	2.49	0.45
1:A:215:SER:OG	2:A:400:CIT:C2	2.67	0.43
1:A:84:ALA:HB2	1:A:102:LEU:HD21	1.99	0.43
1:A:213:ILE:HG13	1:A:249:TYR:CZ	2.55	0.42
1:B:213:ILE:HG13	1:B:249:TYR:CZ	2.54	0.42
1:D:196:TYR:OH	1:D:228[A]:GLU:OE2	2.38	0.42
3:B:324:EDO:H22	4:B:349:HOH:O	2.19	0.41
1:A:75:LEU:HA	3:A:325:EDO:H21	2.02	0.41
1:B:42:PHE:CD2	1:B:308:LEU:HD21	2.55	0.41
1:D:59:LEU:HB2	1:D:228[A]:GLU:HG3	2.03	0.41
1:D:38:LEU:HD12	1:D:233:LEU:HD23	2.02	0.41
1:A:38:LEU:HD12	1:A:233:LEU:HD23	2.03	0.41
1:A:215:SER:HG	2:A:400:CIT:C2	2.35	0.40
1:B:250:PHE:CD1	1:C:250:PHE:CD1	3.09	0.40
1:C:26:THR:HG22	1:C:257:TYR:CD1	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	315/327 (96%)	308 (98%)	7 (2%)	0	<b>100</b> <b>100</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	312/327 (95%)	305 (98%)	7 (2%)	0	100	100
1	C	287/327 (88%)	276 (96%)	11 (4%)	0	100	100
1	D	286/327 (88%)	275 (96%)	11 (4%)	0	100	100
All	All	1200/1308 (92%)	1164 (97%)	36 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/269 (96%)	256 (100%)	1 (0%)	91	85
1	B	248/269 (92%)	247 (100%)	1 (0%)	91	85
1	C	234/269 (87%)	233 (100%)	1 (0%)	91	85
1	D	232/269 (86%)	231 (100%)	1 (0%)	91	85
All	All	971/1076 (90%)	967 (100%)	4 (0%)	91	85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	216	ASN
1	B	216	ASN
1	C	216	ASN
1	D	216	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	89	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	CIT	B	400	-	12,12,12	1.10	0	17,17,17	1.46	3 (17%)
3	EDO	A	324	-	3,3,3	0.39	0	2,2,2	0.33	0
2	CIT	A	400	-	12,12,12	1.25	1 (8%)	17,17,17	1.68	5 (29%)
3	EDO	A	326	-	3,3,3	0.42	0	2,2,2	0.17	0
3	EDO	D	324	-	3,3,3	0.38	0	2,2,2	0.46	0
3	EDO	C	324	-	3,3,3	0.42	0	2,2,2	0.50	0
2	CIT	C	400	-	12,12,12	1.09	0	17,17,17	1.28	2 (11%)
3	EDO	A	325	-	3,3,3	0.45	0	2,2,2	0.36	0
3	EDO	B	324	-	3,3,3	0.74	0	2,2,2	1.81	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	400	-	-	6/16/16/16	-
3	EDO	A	324	-	-	1/1/1/1	-
2	CIT	A	400	-	-	6/16/16/16	-
3	EDO	A	326	-	-	0/1/1/1	-
3	EDO	D	324	-	-	0/1/1/1	-
3	EDO	C	324	-	-	1/1/1/1	-
2	CIT	C	400	-	-	8/16/16/16	-
3	EDO	A	325	-	-	1/1/1/1	-
3	EDO	B	324	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	CIT	C3-C6	-2.65	1.50	1.53

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	CIT	O6-C6-C3	3.36	118.88	113.05
2	A	400	CIT	O5-C6-C3	-3.32	117.55	122.25
2	B	400	CIT	O6-C6-C3	3.26	118.72	113.05
2	A	400	CIT	O7-C3-C4	-2.83	102.77	109.40
2	C	400	CIT	O6-C6-C3	2.68	117.70	113.05
2	A	400	CIT	C3-C4-C5	-2.51	107.74	113.81
3	B	324	EDO	O1-C1-C2	-2.47	94.16	111.91
2	A	400	CIT	C4-C3-C2	2.36	115.33	109.16
2	C	400	CIT	O7-C3-C6	-2.28	105.66	108.86
2	B	400	CIT	O5-C6-C3	-2.23	119.09	122.25
2	B	400	CIT	C4-C3-C2	2.08	114.57	109.16

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	400	CIT	O7-C3-C6-O5
2	A	400	CIT	O7-C3-C6-O6
2	A	400	CIT	C4-C3-C6-O5
2	A	400	CIT	C4-C3-C6-O6
2	B	400	CIT	C2-C3-C6-O5
2	B	400	CIT	C2-C3-C6-O6
2	B	400	CIT	O7-C3-C6-O5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	B	400	CIT	O7-C3-C6-O6
2	C	400	CIT	C2-C3-C6-O5
2	C	400	CIT	C2-C3-C6-O6
2	C	400	CIT	O7-C3-C6-O5
2	C	400	CIT	O7-C3-C6-O6
3	A	324	EDO	O1-C1-C2-O2
3	C	324	EDO	O1-C1-C2-O2
2	A	400	CIT	C1-C2-C3-O7
2	C	400	CIT	C2-C3-C4-C5
2	C	400	CIT	O7-C3-C4-C5
3	A	325	EDO	O1-C1-C2-O2
2	C	400	CIT	C6-C3-C4-C5
2	A	400	CIT	C1-C2-C3-C4
2	B	400	CIT	C1-C2-C3-O7
2	C	400	CIT	C1-C2-C3-O7
2	B	400	CIT	C1-C2-C3-C4
3	B	324	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	400	CIT	5	0
3	D	324	EDO	1	0
3	C	324	EDO	1	0
3	A	325	EDO	2	0
3	B	324	EDO	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	312/327 (95%)	0.02	12 (3%) 40 40	10, 18, 40, 52	0
1	B	311/327 (95%)	0.23	20 (6%) 19 18	10, 21, 50, 65	0
1	C	288/327 (88%)	0.08	12 (4%) 36 35	10, 18, 43, 50	0
1	D	290/327 (88%)	0.09	13 (4%) 33 32	10, 17, 41, 61	0
All	All	1201/1308 (91%)	0.10	57 (4%) 31 30	10, 18, 44, 65	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	50	LEU	6.5
1	C	50	LEU	5.8
1	B	273	PHE	5.1
1	B	268	LEU	5.1
1	B	272	PRO	5.0
1	D	310	GLY	5.0
1	B	24	ILE	4.9
1	D	309	VAL	4.9
1	B	314	HIS	4.3
1	C	310	GLY	4.2
1	C	5	LEU	3.9
1	C	309	VAL	3.9
1	C	22	THR	3.9
1	D	266	GLN	3.8
1	A	269	GLU	3.7
1	A	23	GLY	3.7
1	A	145	GLY	3.6
1	B	262	ASP	3.5
1	D	220	LEU	3.4
1	B	313	HIS	3.4
1	B	266	GLN	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	263	MET	3.2
1	C	13	ASP	3.2
1	D	5	LEU	3.1
1	B	312	ARG	3.1
1	B	19	SER	3.1
1	A	144	GLY	3.0
1	A	24	ILE	2.9
1	C	220	LEU	2.9
1	B	20	ASN	2.7
1	C	20	ASN	2.7
1	D	263	MET	2.7
1	A	289	LYS	2.6
1	D	275	SER	2.6
1	B	261	LEU	2.6
1	A	277	ARG	2.6
1	B	13	ASP	2.6
1	D	18	GLN	2.5
1	D	6	ASP	2.4
1	C	12	LEU	2.4
1	D	264	LEU	2.4
1	A	262	ASP	2.4
1	D	257	TYR	2.4
1	B	263	MET	2.3
1	C	21	ARG	2.3
1	B	259	ASN	2.3
1	B	40	GLN	2.2
1	C	260	GLN	2.2
1	A	312	ARG	2.2
1	D	267	GLN	2.2
1	A	20	ASN	2.2
1	B	11	ILE	2.2
1	B	2	LYS	2.1
1	A	273	PHE	2.1
1	B	146	ALA	2.0
1	C	257	TYR	2.0
1	B	289	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CIT	B	400	13/13	0.52	0.34	44,47,53,54	0
2	CIT	C	400	13/13	0.64	0.29	62,65,71,73	0
2	CIT	A	400	13/13	0.65	0.27	29,38,44,44	13
3	EDO	A	325	4/4	0.74	0.26	38,40,40,43	0
3	EDO	C	324	4/4	0.80	0.21	37,38,39,41	0
3	EDO	D	324	4/4	0.86	0.15	35,36,38,39	0
3	EDO	A	324	4/4	0.93	0.22	26,32,32,33	0
3	EDO	A	326	4/4	0.94	0.18	34,38,39,40	0
3	EDO	B	324	4/4	0.94	0.23	19,23,23,23	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.