



wwPDB EM Validation Summary Report

Feb 27, 2024 – 07:06 AM EST


PDB ID : 6V92
EMDB ID : EMD-21114
Title : RSC-NCP
Authors : Patel, A.B.; Moore, C.M.; Greber, B.J.; Nogales, E.
Deposited on : 2019-12-13
Resolution : 20.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 20.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	477	
2	R	1359	
3	B	467	
4	P	157	
5	C	78	
6	D	180	
7	E	435	
8	F	889	



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	G	885	5% . 94%
10	H	625	16% 5% 79%
11	I	557	44% 8% . 47%
11	J	557	18% . 79%
11	K	557	17% . 80%
11	L	557	44% 10% 46%
12	M	483	52% 12% 36%
13	N	581	58% 13% . 29%
14	O	502	64% 13% 24%
15	Q	426	48% 13% . 38%
16	S	883	96%
17	2	28	96%
18	3	19	47% 100%
18	4	19	95% 5%
19	5	14	43% 100%
20	6	15	100%
21	7	49	96%
22	i	146	6% 100%
22	j	146	10% 100%
23	a	136	5% 69% . 29%
23	e	136	71% . 27%
24	b	103	76% 24%
24	f	103	79% . 17%
25	c	130	78% 5% 17%
25	g	130	8% 74% 6% 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
26	d	126	 10% 71% 5% 24%
26	h	126	 10% 69% 6% 25%

2 Entry composition [i](#)

There are 27 unique types of molecules in this entry. The entry contains 44583 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin-related protein 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	399	3227	2081	528	603	15	3	0

- Molecule 2 is a protein called Nuclear protein STH1/NPS1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	R	325	2663	1668	487	504	4	0	0

- Molecule 3 is a protein called Actin-like protein ARP9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	B	396	3198	2053	523	615	7	1	0

- Molecule 4 is a protein called Regulator of Ty1 transposition protein 102.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	P	54	490	313	84	92	1	0	0

- Molecule 5 is a protein called High temperature lethal protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	C	60	493	301	92	96	4	0	0

- Molecule 6 is a protein called Chromatin structure-remodeling complex protein RSC14.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	D	100	772	490	132	148	2	0	0

- Molecule 7 is a protein called Chromatin structure-remodeling complex subunit RSC7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	E	120	978	610	166	200	2	0	0

- Molecule 8 is a protein called Chromatin structure-remodeling complex subunit RSC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	F	67	536	346	94	95	1	0	0

- Molecule 9 is a protein called Chromatin structure-remodeling complex protein RSC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	G	53	422	270	71	79	2	0	0

- Molecule 10 is a protein called Chromatin structure-remodeling complex subunit RSC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	H	131	1083	696	175	205	7	0	0

- Molecule 11 is a protein called Chromatin structure-remodeling complex protein RSC8.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	I	293	2416	1537	423	448	8	0	0
11	J	115	924	579	149	190	6	0	0
11	K	109	878	554	139	179	6	0	0
11	L	298	2445	1557	428	452	8	0	0

- Molecule 12 is a protein called Chromatin structure-remodeling complex protein RSC6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	M	310	2474	1558	414	496	6	0	0

- Molecule 13 is a protein called Chromatin structure-remodeling complex subunit RSC9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	N	412	3275	2105	540	612	18	0	0

- Molecule 14 is a protein called Chromatin structure-remodeling complex protein RSC58.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
14	O	384	3145	2025	529	581	10	0	0

- Molecule 15 is a protein called Chromatin structure-remodeling complex subunit SFH1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
15	Q	264	2137	1349	362	418	8	0	0

- Molecule 16 is a protein called Chromatin structure-remodeling complex protein RSC30.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
16	S	34	278	182	41	54	1	0	0

- Molecule 17 is a protein called Unknown Protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
17	2	28	140	84	28	28	0	0

- Molecule 18 is a protein called Unknown Protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
18	3	19	95	57	19	19	0	0
18	4	19	95	57	19	19	0	0

- Molecule 19 is a protein called Unknown Protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
19	5	14	70	42	14	14	0	0

- Molecule 20 is a protein called Unknown Protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
20	6	15	Total	C	N	O	0	0
			75	45	15	15		

- Molecule 21 is a protein called Unknown Protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
21	7	49	Total	C	N	O	0	0
			245	147	49	49		

- Molecule 22 is a DNA chain called DNA (146-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
22	i	146	Total	C	N	O	P	0	0
			2990	1431	540	874	145		
22	j	146	Total	C	N	O	P	0	0
			2990	1431	540	874	145		

- Molecule 23 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	a	97	Total	C	N	O	S	0	0
			801	505	155	137	4		
23	e	99	Total	C	N	O	S	0	0
			816	514	158	140	4		

- Molecule 24 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	b	78	Total	C	N	O	S	0	0
			619	391	120	107	1		
24	f	85	Total	C	N	O	S	0	0
			683	430	136	116	1		

- Molecule 25 is a protein called Histone H2A type 1-B/E.


Mol	Chain	Residues	Atoms				AltConf	Trace
25	c	108	Total	C	N	O	0	0
			835	526	165	144		
25	g	104	Total	C	N	O	0	0
			805	508	157	140		

- Molecule 26 is a protein called Histone H2B type 1-K.


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
26	d	96	754	473	138	141	2	0	0
26	h	94	735	461	134	138	2	0	0

- Molecule 27 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
27	I	1	Total	Zn	0
			1	1	




● Molecule 11: Chromatin structure-remodeling complex protein RSC8




● Molecule 11: Chromatin structure-remodeling complex protein RSC8







• Molecule 11: Chromatin structure-remodeling complex protein RSC8




• Molecule 12: Chromatin structure-remodeling complex protein RSC6







• Molecule 13: Chromatin structure-remodeling complex subunit RSC9




• Molecule 14: Chromatin structure-remodeling complex protein RSC58







• Molecule 15: Chromatin structure-remodeling complex subunit SFH1




• Molecule 16: Chromatin structure-remodeling complex protein RSC30




• Molecule 21: Unknown Protein




• Molecule 22: DNA (146-MER)




• Molecule 22: DNA (146-MER)





• Molecule 23: Histone H3.1





• Molecule 23: Histone H3.1





• Molecule 24: Histone H4





• Molecule 24: Histone H4





• Molecule 25: Histone H2A type 1-B/E




• Molecule 25: Histone H2A type 1-B/E



• Molecule 26: Histone H2B type 1-K



• Molecule 26: Histone H2B type 1-K



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	13337	Depositor
Resolution determination method	FSC 0.5 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TALOS ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.110	Depositor
Minimum map value	-0.064	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.03	Depositor
Map size (Å)	437.76, 437.76, 437.76	wwPDB
Map dimensions	128, 128, 128	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	3.42, 3.42, 3.42	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/3303	0.60	1/4465 (0.0%)
2	R	0.36	2/2697 (0.1%)	0.49	3/3616 (0.1%)
3	B	0.45	0/3269	0.59	0/4432
4	P	0.41	0/501	0.57	0/669
5	C	0.24	0/495	0.36	0/662
6	D	0.27	0/786	0.38	0/1062
7	E	0.28	0/997	0.44	0/1356
8	F	0.27	0/551	0.41	0/748
9	G	0.26	0/431	0.43	0/584
10	H	0.29	0/1108	0.48	0/1497
11	I	0.32	0/2474	0.41	0/3343
11	J	0.30	0/926	0.40	0/1233
11	K	0.27	0/879	0.39	0/1172
11	L	0.29	0/2502	0.40	0/3376
12	M	0.28	0/2515	0.42	0/3423
13	N	0.30	0/3334	0.42	0/4515
14	O	0.30	0/3216	0.42	1/4358 (0.0%)
15	Q	0.28	0/2181	0.45	0/2964
16	S	0.26	0/281	0.35	0/378
22	i	0.38	0/3354	0.79	0/5175
22	j	0.36	0/3354	0.78	0/5175
23	a	0.40	0/813	0.59	0/1090
23	e	0.46	0/828	0.61	0/1109
24	b	0.38	0/626	0.62	0/837
24	f	0.44	0/691	0.63	0/923
25	c	0.39	0/845	0.60	0/1139
25	g	0.39	0/815	0.65	0/1100
26	d	0.41	0/765	0.63	0/1025
26	h	0.40	0/746	0.58	0/1003
All	All	0.35	2/45283 (0.0%)	0.55	5/62429 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	328	LYS	C-O	5.96	1.34	1.23
2	R	333	ILE	C-O	5.93	1.34	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	380	SER	C-N-CD	5.93	140.86	128.40
14	O	231	VAL	C-N-CA	5.36	135.09	121.70
2	R	333	ILE	CA-C-O	5.28	131.19	120.10
2	R	367	ASP	CB-CG-OD2	5.17	122.95	118.30
2	R	328	LYS	CA-C-O	5.10	130.81	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3227	0	3251	150	0
2	R	2663	0	2746	159	0
3	B	3198	0	3187	136	0
4	P	490	0	467	19	0
5	C	493	0	507	4	0
6	D	772	0	754	11	0
7	E	978	0	935	25	0
8	F	536	0	533	13	0
9	G	422	0	423	9	0
10	H	1083	0	1062	21	0
11	I	2416	0	2358	36	0
11	J	924	0	976	12	0
11	K	878	0	930	13	0
11	L	2445	0	2402	48	0
12	M	2474	0	2465	48	0
13	N	3275	0	3373	47	0
14	O	3145	0	3168	50	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
15	Q	2137	0	2069	42	0
16	S	278	0	287	6	0
17	2	140	0	34	1	0
18	3	95	0	21	0	0
18	4	95	0	21	1	0
19	5	70	0	16	0	0
20	6	75	0	17	0	0
21	7	245	0	63	2	0
22	i	2990	0	1652	0	0
22	j	2990	0	1652	0	0
23	a	801	0	839	0	0
23	e	816	0	856	0	0
24	b	619	0	659	0	0
24	f	683	0	729	0	0
25	c	835	0	897	0	0
25	g	805	0	861	0	0
26	d	754	0	782	0	0
26	h	735	0	756	0	0
27	I	1	0	0	0	0
All	All	44583	0	41748	652	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 652 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:R:353:PHE:CZ	3:B:447:PHE:CE2	2.05	1.45
2:R:353:PHE:CZ	3:B:447:PHE:HE2	1.34	1.41
2:R:353:PHE:CE1	3:B:447:PHE:HE2	1.47	1.33
2:R:365:GLU:HG2	3:B:415:TYR:CD2	1.65	1.32
2:R:353:PHE:CE2	3:B:447:PHE:CE2	2.21	1.29

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/477 (82%)	369 (94%)	18 (5%)	5 (1%)	12	48
2	R	319/1359 (24%)	296 (93%)	23 (7%)	0	100	100
3	B	391/467 (84%)	368 (94%)	21 (5%)	2 (0%)	29	69
4	P	46/157 (29%)	43 (94%)	2 (4%)	1 (2%)	6	35
5	C	58/78 (74%)	56 (97%)	2 (3%)	0	100	100
6	D	96/180 (53%)	90 (94%)	6 (6%)	0	100	100
7	E	118/435 (27%)	108 (92%)	10 (8%)	0	100	100
8	F	63/889 (7%)	56 (89%)	7 (11%)	0	100	100
9	G	51/885 (6%)	49 (96%)	2 (4%)	0	100	100
10	H	127/625 (20%)	114 (90%)	10 (8%)	3 (2%)	6	33
11	I	289/557 (52%)	274 (95%)	15 (5%)	0	100	100
11	J	113/557 (20%)	109 (96%)	4 (4%)	0	100	100
11	K	105/557 (19%)	101 (96%)	4 (4%)	0	100	100
11	L	292/557 (52%)	277 (95%)	15 (5%)	0	100	100
12	M	302/483 (62%)	281 (93%)	21 (7%)	0	100	100
13	N	408/581 (70%)	380 (93%)	28 (7%)	0	100	100
14	O	376/502 (75%)	352 (94%)	24 (6%)	0	100	100
15	Q	258/426 (61%)	224 (87%)	34 (13%)	0	100	100
16	S	32/883 (4%)	32 (100%)	0	0	100	100
23	a	95/136 (70%)	92 (97%)	3 (3%)	0	100	100
23	e	97/136 (71%)	95 (98%)	2 (2%)	0	100	100
24	b	76/103 (74%)	75 (99%)	1 (1%)	0	100	100
24	f	83/103 (81%)	80 (96%)	3 (4%)	0	100	100
25	c	106/130 (82%)	101 (95%)	5 (5%)	0	100	100
25	g	102/130 (78%)	98 (96%)	3 (3%)	1 (1%)	15	55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
26	d	94/126 (75%)	89 (95%)	4 (4%)	1 (1%)	14	52
26	h	92/126 (73%)	88 (96%)	0	4 (4%)	2	22
All	All	4581/11645 (39%)	4297 (94%)	267 (6%)	17 (0%)	38	72

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	48	ASP
10	H	528	ILE
26	h	120	SER
1	A	343	SER
26	d	101	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/420 (85%)	338 (95%)	19 (5%)	22	47
2	R	297/1228 (24%)	294 (99%)	3 (1%)	76	86
3	B	363/423 (86%)	343 (94%)	20 (6%)	21	47
4	P	53/140 (38%)	52 (98%)	1 (2%)	57	75
5	C	58/75 (77%)	58 (100%)	0	100	100
6	D	82/151 (54%)	82 (100%)	0	100	100
7	E	113/388 (29%)	105 (93%)	8 (7%)	14	39
8	F	60/810 (7%)	60 (100%)	0	100	100
9	G	48/832 (6%)	47 (98%)	1 (2%)	53	72
10	H	126/578 (22%)	124 (98%)	2 (2%)	62	79
11	I	268/500 (54%)	261 (97%)	7 (3%)	46	66
11	J	111/500 (22%)	110 (99%)	1 (1%)	78	87
11	K	106/500 (21%)	106 (100%)	0	100	100
11	L	270/500 (54%)	270 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
12	M	286/435 (66%)	281 (98%)	5 (2%)	60	78
13	N	374/521 (72%)	366 (98%)	8 (2%)	53	72
14	O	358/462 (78%)	355 (99%)	3 (1%)	81	89
15	Q	243/384 (63%)	233 (96%)	10 (4%)	30	55
16	S	33/824 (4%)	32 (97%)	1 (3%)	41	63
23	a	85/111 (77%)	82 (96%)	3 (4%)	36	59
23	e	86/111 (78%)	83 (96%)	3 (4%)	36	59
24	b	63/79 (80%)	63 (100%)	0	100	100
24	f	70/79 (89%)	66 (94%)	4 (6%)	20	45
25	c	85/100 (85%)	79 (93%)	6 (7%)	14	39
25	g	83/100 (83%)	76 (92%)	7 (8%)	11	33
26	d	82/105 (78%)	77 (94%)	5 (6%)	18	44
26	h	80/105 (76%)	77 (96%)	3 (4%)	33	57
All	All	4240/10461 (40%)	4120 (97%)	120 (3%)	46	65

5 of 120 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	I	270	VAL
25	g	35	ARG
13	N	166	ARG
24	f	93	GLN
26	h	96	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 62 such sidechains are listed below:

Mol	Chain	Res	Type
12	M	61	ASN
26	d	92	GLN
13	N	559	HIS
26	d	60	ASN
25	g	73	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
21	7	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	7	57:UNK	C	60:UNK	N	4.98


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21114. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)

6.1.1 Primary map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 64



Y Index: 64



Z Index: 64

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 68



Y Index: 71




Z Index: 59


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.03. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 1633 nm³; this corresponds to an approximate mass of 1475 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.050 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-21114 and PDB model 6V92. Per-residue inclusion information can be found in section 3 on page 10.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.03 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.03).









































































9.4 Atom inclusion [i](#)



At the recommended contour level, 91% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.03) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9000	 0.0530
2	 1.0000	 0.0480
3	 0.5790	 0.0250
4	 0.9900	 0.0580
5	 0.5710	 0.0050
6	 1.0000	 0.0500
7	 0.9880	 0.0500
A	 0.6960	 0.0330
B	 0.7390	 0.0370
C	 0.9920	 0.0650
D	 0.8760	 0.0380
E	 0.9690	 0.0740
F	 1.0000	 0.0290
G	 0.8230	 0.0470
H	 0.9570	 0.0530
I	 0.9710	 0.0510
J	 0.9720	 0.0800
K	 0.9690	 0.0810
L	 0.9720	 0.0510
M	 0.9430	 0.0480
N	 0.9370	 0.0490
O	 0.9940	 0.0650
P	 0.6980	 0.0590
Q	 0.9560	 0.0660
R	 0.8230	 0.0410
S	 0.7630	 0.0680
a	 0.9360	 0.0280
b	 0.9900	 0.0440
c	 0.9580	 0.0380
d	 0.8560	 0.0200
e	 0.9550	 0.0430
f	 0.9950	 0.0050
g	 0.8830	 0.0110
h	 0.9370	 0.0120
i	 0.8990	 0.0860
j	 0.8970	 0.0930

