

wwPDB X-ray Structure Validation Summary Report (i)

Dec 3, 2023 - 07.58 pm GMT

PDB ID : 2VB2

Title : Crystal structure of Cu(I)CusF

Authors: Xue, Y.; Davis, A.V.; Balakrishnan, G.; Stasser, J.P.; Staehlin, B.M.; Focia,

P.; Spiro, T.G.; Penner-Hahn, J.E.; O'Halloran, T.V.

Deposited on : 2007-09-06

Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

There are no overall percentile quality scores available for this entry.

MolProbity failed to run properly; EDS was not executed - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 634 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called CATION EFFLUX SYSTEM PROTEIN CUSF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	76	Total	С	N	0	S	0	0	1
_			582	371	99	109	3			

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	X	1	Total 5	O 4	S 1	0	0

• Molecule 3 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	X	1	Total Cu 1 1	0	0	

• Molecule 4 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
4	X	46	Total O 46 46	0	0	

MolProbity failed to run properly; EDS was not executed - this section is therefore empty.



3 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source		
Space group	P 21 21 21	Depositor		
Cell constants	38.47Å 41.26Å 44.57Å	Donogitor		
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor		
Resolution (Å)	30.28 - 1.70	Depositor		
% Data completeness	96.3 (30.28-1.70)	Depositor		
(in resolution range)	30.3 (30.25-1.70)			
R_{merge}	0.05	Depositor		
R_{sym}	(Not available)	Depositor		
Refinement program	REFMAC 5.2.0019	Depositor		
R, R_{free}	0.238 , 0.263	Depositor		
Estimated twinning fraction	No twinning to report.	Xtriage		
Total number of atoms	634	wwPDB-VP		
Average B, all atoms (Å ²)	37.0	wwPDB-VP		



4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	$_{ m gths}$	Bond angles		
WIOI	or Type	Chain	nes Li	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	X	1088	-	4,4,4	0.15	0	6,6,6	0.10	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

5.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

5.4 Ligands (i)

EDS was not executed - this section is therefore empty.

5.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

