

wwPDB X-ray Structure Validation Summary Report (i)

Oct 5, 2023 – 01:53 AM EDT

PDB ID : 6VBP

Title: Crystal structure of anti-HIV-1 antibody DH815 bound to gp120 V2 peptide

Authors: Janus, B.M.; Ofek, G.

Deposited on : 2019-12-19

Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 27397 atoms, of which 13232 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DH815 light chain.

| Mol | Chain | Residues | | Atoms | | | | ZeroOcc | AltConf | Trace | |
|-----|-------|----------|-------|-------|------|-----|-----|---------|---------|-------|---|
| 1 | Т | 218 | Total | С | Н | N | О | S | 0 | 0 | 0 |
| 1 | ь | 210 | 3298 | 1047 | 1623 | 284 | 339 | 5 | 0 | 0 | U |
| 1 | В | 219 | Total | С | Н | N | О | S | 0 | 0 | 0 |
| 1 | Ъ | 219 | 3313 | 1051 | 1631 | 285 | 341 | 5 | 0 | 0 | 0 |
| 1 | D | 219 | Total | С | Н | N | О | S | 0 | 0 | 0 |
| 1 | ע | 219 | 3320 | 1053 | 1634 | 285 | 343 | 5 | 0 | U | U |
| 1 | Т | 219 | Total | С | Н | N | О | S | 0 | 0 | 0 |
| 1 | 1 | 219 | 3320 | 1053 | 1634 | 285 | 343 | 5 | | | |

• Molecule 2 is a protein called DH815 heavy chain.

| Mol | Chain | Residues | | | Atoms | S | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-------|-----|-----|---|---------|---------|-------|
| 2 | Н | 214 | Total | С | Н | N | О | S | 0 | 0 | 0 |
| 2 | 11 | 214 | 3177 | 1024 | 1564 | 271 | 313 | 5 | | U | 0 |
| 2 | Λ | 213 | Total | С | Н | N | О | S | 0 | 0 | 0 |
| 2 | A | 210 | 3176 | 1023 | 1567 | 270 | 311 | 5 | 0 | | |
| 2 | С | 221 | Total | С | Н | N | О | S | 0 | 0 | 0 |
| 2 | | 221 | 3296 | 1055 | 1630 | 280 | 325 | 6 | | | 0 |
| 2 | ī | 214 | Total | С | Н | N | О | S | 0 | 0 | 0 |
| | J | 214 | 3149 | 1021 | 1541 | 270 | 312 | 5 | U | U | U |

• Molecule 3 is a protein called Envelope glycoprotein gp160.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------------|----------|-------|----|-----|----|----|---------|---------|-------|
| 3 | F | 1.4 | Total | С | Н | N | О | 0 | 0 | 0 |
| 3 | I' | 14 | 231 | 79 | 118 | 17 | 17 | 0 | U | U |
| 3 | Е | 13 | Total | С | Н | N | О | 0 | 0 | 0 |
| 3 | <u> 1</u> 2 | 10 | 190 | 68 | 90 | 16 | 16 | 0 0 | 0 | U |
| 3 | G | 14 | Total | С | Н | N | О | 0 | 0 | 0 |
| 3 | G | 14 | 230 | 79 | 117 | 17 | 17 | 0 | U | U |
| 3 | K | 12 | Total | С | Н | N | О | 0 | 0 | 0 |
|) | 17 | 12 | 173 | 62 | 83 | 13 | 15 | | U | |



• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 4 | С | 1 | Total Cl 1 1 | 0 | 0 |

• Molecule 5 is water.

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|------------------|---------|---------|
| 5 | L | 75 | Total O 75 75 | 0 | 0 |
| 5 | Н | 56 | Total O 56 56 | 0 | 0 |
| 5 | В | 63 | Total O 63 63 | 0 | 0 |
| 5 | A | 54 | Total O 54 54 | 0 | 0 |
| 5 | D | 88 | Total O 88 88 | 0 | 0 |
| 5 | С | 70 | Total O 70 70 | 0 | 0 |
| 5 | F | 3 | Total O 3 3 | 0 | 0 |
| 5 | Е | 4 | Total O 4 4 | 0 | 0 |
| 5 | I | 82 | Total O 82 82 | 0 | 0 |
| 5 | J | 28 | Total O 28 28 | 0 | 0 |

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

| Property | Value | Source |
|---|--|-----------|
| Space group | P 1 21 1 | Depositor |
| Cell constants | 69.90Å 158.53Å 92.65Å | Depositor |
| a, b, c, α , β , γ | 90.00° 110.27° 90.00° | Depositor |
| Resolution (Å) | 45.15 - 2.30 | Depositor |
| % Data completeness | 96.0 (45.15-2.30) | Depositor |
| (in resolution range) | , | _ |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | (Not available) | Depositor |
| $< I/\sigma(I) > 1$ | 0.37 (at 2.29Å) | Xtriage |
| Refinement program | PHENIX 1.14_3228 | Depositor |
| R, R_{free} | 0.180 , 0.229 | Depositor |
| Wilson B-factor (\mathring{A}^2) | 37.5 | Xtriage |
| Anisotropy | 0.118 | Xtriage |
| L-test for twinning ² | $< L > = 0.46, < L^2> = 0.29$ | Xtriage |
| Estimated twinning fraction | 0.033 for h,-k,-h-l | Xtriage |
| Total number of atoms | 27397 | wwPDB-VP |
| Average B, all atoms (\mathring{A}^2) | 54.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

