



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 12:05 am BST

PDB ID : 1VCA
Title : CRYSTAL STRUCTURE OF AN INTEGRIN-BINDING FRAGMENT OF VASCULAR CELL ADHESION MOLECULE-1 AT 1.8 ANGSTROMS RESOLUTION
Authors : Jones, E.Y.; Harlos, K.; Bottomley, M.J.; Robinson, R.C.; Driscoll, P.C.; Edwards, R.M.; Clements, J.M.; Dudgeon, T.J.; Stuart, D.I.
Deposited on : 1995-03-21
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	202	
1	B	202	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN VASCULAR CELL ADHESION MOLECULE-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	199	1553	980	253	311	9	0	0	0
1	B	199	1553	980	253	311	9	0	0	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	148	148	148	0	0
2	B	106	106	106	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: HUMAN VASCULAR CELL ADHESION MOLECULE-1

Chain A:  76% 20% ..



- Molecule 1: HUMAN VASCULAR CELL ADHESION MOLECULE-1

Chain B:  80% 15% ..



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.70Å 66.50Å 113.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (15.00-1.80)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.202 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3360	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/1583	0.90	1/2147 (0.0%)
1	B	0.67	0/1583	0.90	2/2147 (0.1%)
All	All	0.66	0/3166	0.90	3/4294 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	171	CYS	CA-CB-SG	-5.76	103.63	114.00
1	B	154	LEU	CA-CB-CG	5.32	127.54	115.30
1	A	171	CYS	CA-CB-SG	-5.13	104.77	114.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	119	TYR	Sidechain
1	B	119	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1553	0	1544	22	0
1	B	1553	0	1544	12	0
2	A	148	0	0	1	0
2	B	106	0	0	1	0
All	All	3360	0	3088	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:ARG:HG2	1:A:147:LYS:H	1.39	0.87
1:B:197:ILE:O	1:B:199:PRO:HD3	1.93	0.68
1:B:85:GLN:HG3	2:B:472:HOH:O	1.96	0.64
1:A:168:VAL:HG12	1:A:192:GLU:HG2	1.82	0.59
1:A:162:ILE:HD11	1:A:197:ILE:HG22	1.87	0.57
1:A:14:GLN:HE21	1:A:91:PHE:H	1.54	0.56
1:B:185:THR:HG23	1:B:186:VAL:HG23	1.89	0.55
1:A:185:THR:HG22	1:A:186:VAL:HG23	1.91	0.53
1:A:129:LEU:HB2	1:A:170:VAL:HG22	1.91	0.52
1:B:10:ARG:HD2	1:B:87:GLU:OE1	2.09	0.52
1:A:12:LEU:HD22	1:A:89:TYR:CD1	2.45	0.51
1:B:53:THR:HG22	1:B:54:SER:N	2.26	0.51
1:A:121:PHE:C	1:A:121:PHE:CD1	2.83	0.50
1:A:140:PHE:O	1:A:142:GLU:N	2.45	0.50
1:A:5:THR:HB	1:A:84:ILE:HD11	1.93	0.49
1:B:112:LYS:HG3	1:B:155:GLU:HG2	1.94	0.49
1:A:146:ARG:HG2	1:A:147:LYS:N	2.20	0.49
1:A:172:ARG:HD3	2:A:449:HOH:O	2.13	0.48
1:B:5:THR:HB	1:B:84:ILE:HD11	1.95	0.47
1:A:99:LEU:HD23	1:A:111:VAL:HG22	1.97	0.47
1:A:135:MET:HE3	1:A:156:VAL:HG11	1.98	0.45
1:A:59:ASN:HA	1:A:60:PRO:HA	1.76	0.45
1:B:49:ASN:ND2	1:B:54:SER:OG	2.50	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:LYS:HG2	1:A:186:VAL:HG22	1.99	0.45
1:A:49:ASN:ND2	1:A:54:SER:OG	2.50	0.44
1:B:59:ASN:HA	1:B:60:PRO:HA	1.77	0.44
1:B:107:LYS:O	1:B:160:PRO:HD2	2.18	0.43
1:B:121:PHE:C	1:B:121:PHE:CD1	2.92	0.42
1:A:127:ASP:OD2	1:A:172:ARG:NH2	2.51	0.42
1:A:120:PRO:HG2	1:A:123:ARG:CG	2.49	0.42
1:A:128:LEU:HD23	1:A:135:MET:CE	2.50	0.41
1:B:146:ARG:HB3	1:B:147:LYS:H	1.62	0.41
1:A:185:THR:HG22	1:A:186:VAL:CG2	2.51	0.40
1:A:120:PRO:HG2	1:A:123:ARG:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/202 (98%)	188 (95%)	6 (3%)	3 (2%)	10	2
1	B	197/202 (98%)	181 (92%)	10 (5%)	6 (3%)	4	0
All	All	394/404 (98%)	369 (94%)	16 (4%)	9 (2%)	6	1

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	LEU
1	A	182	SER
1	B	142	GLU
1	B	144	ALA
1	B	145	ASP
1	B	179	GLU
1	A	181	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	143	ASP
1	B	181	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/185 (98%)	170 (93%)	12 (7%)	16	5
1	B	182/185 (98%)	169 (93%)	13 (7%)	14	5
All	All	364/370 (98%)	339 (93%)	25 (7%)	15	5

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	21	LEU
1	A	70	LEU
1	A	121	PHE
1	A	141	LEU
1	A	149	LEU
1	A	162	ILE
1	A	169	LEU
1	A	178	ASP
1	A	183	VAL
1	A	185	THR
1	A	198	SER
1	B	21	LEU
1	B	59	ASN
1	B	68	SER
1	B	92	PRO
1	B	109	ILE
1	B	121	PHE
1	B	143	ASP
1	B	146	ARG
1	B	154	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	169	LEU
1	B	170	VAL
1	B	179	GLU
1	B	191	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	A	38	GLN
1	A	49	ASN
1	A	59	ASN
1	B	38	GLN
1	B	49	ASN
1	B	59	ASN
1	B	85	GLN
1	B	138	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.