



wwPDB X-ray Structure Validation Summary Report

Dec 9, 2024 – 10:04 AM EST


PDB ID : 8VC7
Title : Crystal Structure of Human BTN2A1 ectodomain in complex with Antagonist 2A1.9 Fab
Authors : Ramesh, A.; Roy, S.; Adams, E.
Deposited on : 2023-12-13
Resolution : 2.76 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1606 (2.78-2.74)
Clashscore	180529	1689 (2.78-2.74)
Ramachandran outliers	177936	1665 (2.78-2.74)
Sidechain outliers	177891	1665 (2.78-2.74)
RSRZ outliers	164620	1606 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	
1	D	215	
2	B	233	
2	J	233	
3	C	231	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	231	 <p>5% 76% 15% • 7%</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9797 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Human IgG1 Fragment Antibody Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	214	Total 1614	C 1003	N 271	O 334	S 6	0	0	0
1	A	214	Total 1596	C 992	N 268	O 330	S 6	0	0	0

- Molecule 2 is a protein called Human IgG1 Fragment Antibody Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	J	215	Total 1603	C 1017	N 263	O 317	S 6	0	0	0
2	B	215	Total 1606	C 1020	N 263	O 317	S 6	0	0	0

- Molecule 3 is a protein called Butyrophilin subfamily 2 member A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	215	Total 1650	C 1039	N 288	O 312	S 11	11	0	0
3	E	215	Total 1658	C 1047	N 288	O 312	S 11	11	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	ALA	-	expression tag	UNP Q7KYR7
C	-1	ASP	-	expression tag	UNP Q7KYR7
C	0	LEU	-	expression tag	UNP Q7KYR7
C	219	SER	CYS	engineered mutation	UNP Q7KYR7
C	220	GLY	-	expression tag	UNP Q7KYR7
C	221	SER	-	expression tag	UNP Q7KYR7
C	222	GLY	-	expression tag	UNP Q7KYR7
C	223	LEU	-	expression tag	UNP Q7KYR7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	224	GLU	-	expression tag	UNP Q7KYR7
C	225	VAL	-	expression tag	UNP Q7KYR7
C	226	LEU	-	expression tag	UNP Q7KYR7
C	227	PHE	-	expression tag	UNP Q7KYR7
C	228	GLN	-	expression tag	UNP Q7KYR7
E	-2	ALA	-	expression tag	UNP Q7KYR7
E	-1	ASP	-	expression tag	UNP Q7KYR7
E	0	LEU	-	expression tag	UNP Q7KYR7
E	219	SER	CYS	engineered mutation	UNP Q7KYR7
E	220	GLY	-	expression tag	UNP Q7KYR7
E	221	SER	-	expression tag	UNP Q7KYR7
E	222	GLY	-	expression tag	UNP Q7KYR7
E	223	LEU	-	expression tag	UNP Q7KYR7
E	224	GLU	-	expression tag	UNP Q7KYR7
E	225	VAL	-	expression tag	UNP Q7KYR7
E	226	LEU	-	expression tag	UNP Q7KYR7
E	227	PHE	-	expression tag	UNP Q7KYR7
E	228	GLN	-	expression tag	UNP Q7KYR7

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	E	1	Total	C	N	O	0	0
			14	8	1	5		
4	E	1	Total	C	N	O	0	0
			14	8	1	5		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Human IgG1 Fragment Antibody Light Chain

Chain D:  90% 10%





- Molecule 1: Human IgG1 Fragment Antibody Light Chain

Chain A:  82% 16% 2%




- Molecule 2: Human IgG1 Fragment Antibody Heavy Chain

Chain J:  80% 12% 8%




- Molecule 2: Human IgG1 Fragment Antibody Heavy Chain

Chain B:  76% 15% 8%





- Molecule 3: Butyrophilin subfamily 2 member A1

Chain C: 



- Molecule 3: Butyrophilin subfamily 2 member A1

Chain E: 



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.91Å 200.98Å 206.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.46 – 2.76 39.46 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.46-2.76) 91.1 (39.46-2.76)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.77Å)	Xtrriage
Refinement program	PHENIX 1.21rc1_5015	Depositor
R, R_{free}	0.218 , 0.261 0.218 , 0.260	Depositor DCC
R_{free} test set	47432 reflections (4.04%)	wwPDB-VP
Wilson B-factor (Å ²)	37.0	Xtrriage
Anisotropy	0.199	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 30.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.000 for -h,l,k	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9797	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.75 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.9301e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1628	0.68	1/2218 (0.0%)
1	D	0.47	0/1646	0.68	0/2237
2	B	0.48	1/1649 (0.1%)	0.66	0/2255
2	J	0.46	0/1646	0.69	0/2252
3	C	0.44	0/1686	0.69	0/2290
3	E	0.44	0/1695	0.68	0/2300
All	All	0.46	1/9950 (0.0%)	0.68	1/13552 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	99	CYS	CB-SG	-6.11	1.71	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	12	LEU	CA-CB-CG	5.16	127.18	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	54	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1596	0	1512	24	0
1	D	1614	0	1555	8	0
2	B	1606	0	1534	20	0
2	J	1603	0	1528	14	0
3	C	1650	0	1591	24	0
3	E	1658	0	1602	25	0
4	C	42	0	39	0	0
4	E	28	0	26	0	0
All	All	9797	0	9387	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 111 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:3:ILE:HG12	3:E:26:SER:HB2	1.23	1.08
3:C:3:ILE:HG13	3:C:26:SER:HB3	1.52	0.91
3:E:3:ILE:H	3:E:26:SER:HB3	1.40	0.87
1:A:91:GLN:HE21	1:A:98:THR:HG22	1.44	0.83
1:D:21:THR:HG21	1:A:205:PRO:HD3	1.65	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/215 (99%)	206 (97%)	6 (3%)	0	100	100
1	D	212/215 (99%)	205 (97%)	7 (3%)	0	100	100
2	B	211/233 (91%)	205 (97%)	5 (2%)	1 (0%)	25	41
2	J	211/233 (91%)	205 (97%)	6 (3%)	0	100	100
3	C	213/231 (92%)	207 (97%)	5 (2%)	1 (0%)	25	41
3	E	213/231 (92%)	204 (96%)	8 (4%)	1 (0%)	25	41
All	All	1272/1358 (94%)	1232 (97%)	37 (3%)	3 (0%)	44	63

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	5	VAL
3	C	214	PRO
3	E	28	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/190 (95%)	173 (96%)	7 (4%)	27	48
1	D	186/190 (98%)	179 (96%)	7 (4%)	28	49
2	B	175/195 (90%)	168 (96%)	7 (4%)	27	47
2	J	175/195 (90%)	170 (97%)	5 (3%)	37	59
3	C	177/199 (89%)	169 (96%)	8 (4%)	23	42
3	E	177/199 (89%)	170 (96%)	7 (4%)	27	47
All	All	1070/1168 (92%)	1029 (96%)	41 (4%)	28	49

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	53	TYR
3	E	24	HIS
2	B	68	LYS
2	B	162	VAL
3	E	117	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	J	6	GLN
3	C	42	GLN
1	A	4	GLN
2	B	176	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	303	3	14,14,15	0.73	0	17,19,21	1.87	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	302	3	14,14,15	1.27	2 (14%)	17,19,21	3.21	7 (41%)
4	NAG	E	301	3	14,14,15	1.19	1 (7%)	17,19,21	3.88	8 (47%)
4	NAG	E	302	3	14,14,15	0.64	0	17,19,21	1.97	2 (11%)
4	NAG	C	301	3	14,14,15	0.98	1 (7%)	17,19,21	1.39	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	303	3	-	2/6/23/26	0/1/1/1
4	NAG	C	302	3	-	2/6/23/26	0/1/1/1
4	NAG	E	301	3	-	2/6/23/26	0/1/1/1
4	NAG	E	302	3	-	0/6/23/26	0/1/1/1
4	NAG	C	301	3	-	3/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	302	NAG	C1-C2	3.15	1.56	1.52
4	E	301	NAG	C1-C2	3.06	1.56	1.52
4	C	301	NAG	C1-C2	2.04	1.55	1.52
4	C	302	NAG	O5-C5	-2.01	1.39	1.43

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	301	NAG	C2-N2-C7	11.00	137.63	122.90
4	C	302	NAG	C2-N2-C7	10.12	136.47	122.90
4	E	301	NAG	C1-O5-C5	8.54	123.63	112.19
4	E	302	NAG	C1-O5-C5	6.97	121.53	112.19
4	C	303	NAG	C1-O5-C5	6.75	121.23	112.19

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	302	NAG	O5-C5-C6-O6
4	C	302	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	C	301	NAG	C8-C7-N2-C2
4	C	301	NAG	O7-C7-N2-C2
4	C	303	NAG	C8-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/215 (99%)	-0.09	4 (1%) 66 67	24, 34, 57, 87	0
1	D	214/215 (99%)	-0.35	1 (0%) 87 89	23, 33, 45, 83	0
2	B	215/233 (92%)	-0.36	1 (0%) 87 89	22, 33, 49, 68	0
2	J	215/233 (92%)	-0.40	2 (0%) 81 83	23, 32, 48, 69	0
3	C	215/231 (93%)	0.05	5 (2%) 61 61	22, 41, 59, 72	3 (1%)
3	E	215/231 (93%)	0.35	11 (5%) 34 36	28, 44, 71, 90	4 (1%)
All	All	1288/1358 (94%)	-0.13	24 (1%) 66 67	22, 36, 58, 90	7 (0%)

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	215	SER	6.6
3	E	1	GLN	3.5
1	A	215	CYS	3.5
3	E	186	VAL	3.2
3	E	26	SER	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	302	14/15	0.50	0.22	60,70,78,81	0
4	NAG	E	301	14/15	0.51	0.19	47,70,78,80	0
4	NAG	C	301	14/15	0.56	0.17	54,64,71,79	0
4	NAG	E	302	14/15	0.88	0.12	39,53,57,66	0
4	NAG	C	303	14/15	0.89	0.10	46,49,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.