



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 1, 2024 – 04:08 PM EDT

PDB ID : 8VDL  
Title : HB3VAR03 CIDRa1.4 domain with C7 Fab  
Authors : Hurlburt, N.K.; Pancera, M.  
Deposited on : 2023-12-15  
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4708 (2.70-2.66)
Clashscore	180529	5138 (2.70-2.66)
Ramachandran outliers	177936	5071 (2.70-2.66)
Sidechain outliers	177891	5071 (2.70-2.66)
RSRZ outliers	164620	4708 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	164	
2	H	241	
3	L	212	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4067 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HB3VAR03 CIDRa1.4 domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	76	676	452	106	117	1	0	0	0

- Molecule 2 is a protein called C7 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	225	1666	1047	279	331	9	0	0	0

- Molecule 3 is a protein called C7 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	208	1579	989	264	319	7	0	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Zn	0	0
			2	2		
4	H	3	Total	Zn	0	0
			3	3		
4	L	1	Total	Zn	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	7	Total	O	0	0
			7	7		
5	H	63	Total	O	0	0
			63	63		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	L	70	Total	O	0	0
			70	70		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HB3VAR03 CIDRa1.4 domain



- Molecule 2: C7 Heavy Chain



- Molecule 3: C7 Light Chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.65Å 65.35Å 250.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.30 – 2.68 46.30 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.30-2.68) 99.6 (46.30-2.68)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, $R_{free}$	0.217 , 0.266 0.218 , 0.267	Depositor DCC
$R_{free}$ test set	1357 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtrriage
Anisotropy	1.382	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4067	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	C	0.45	0/693	0.54	0/927
2	H	0.46	0/1705	0.64	0/2318
3	L	0.46	0/1618	0.62	1/2209 (0.0%)
All	All	0.46	0/4016	0.61	1/5454 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	1
3	L	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	107	GLY	C-N-CA	-6.49	105.49	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	65	ARG	Sidechain
3	L	36	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	676	0	661	21	0
2	H	1666	0	1614	29	0
3	L	1579	0	1533	20	0
4	C	2	0	0	0	1
4	H	3	0	0	0	0
4	L	1	0	0	0	0
5	C	7	0	0	0	0
5	H	63	0	0	4	0
5	L	70	0	0	3	0
All	All	4067	0	3808	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:108:GLN:HE22	3:L:170:ASN:HD22	1.19	0.90
1:C:582:LEU:O	1:C:586:THR:OG1	2.07	0.73
2:H:215:ASN:HD22	2:H:222:LYS:HG3	1.52	0.73
2:H:209:THR:HG23	2:H:226:LYS:HE3	1.72	0.70
3:L:108:GLN:HE22	3:L:170:ASN:ND2	1.90	0.68
3:L:120:PRO:HD3	3:L:132:LEU:HD23	1.80	0.63
3:L:108:GLN:NE2	3:L:170:ASN:HD22	1.95	0.61
2:H:48:LEU:HD22	2:H:64:VAL:HG11	1.83	0.59
3:L:58:ARG:NH2	3:L:79:ASP:OD1	2.38	0.57
2:H:67:ARG:HG2	2:H:84:ASN:O	2.07	0.55
2:H:68:PHE:CE1	2:H:83:MET:HB3	2.42	0.54
2:H:59:TYR:OH	5:H:401:HOH:O	2.15	0.54
2:H:226:LYS:NZ	5:H:406:HOH:O	2.42	0.53
2:H:61:SER:OG	2:H:64:VAL:HG22	2.10	0.52
1:C:702:HIS:O	1:C:706:ILE:N	2.42	0.51
2:H:143:SER:O	5:H:402:HOH:O	2.19	0.50
2:H:56:LYS:HA	2:H:56:LYS:HE2	1.93	0.50
1:C:582:LEU:HG	1:C:699:LEU:HD21	1.94	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:697:LYS:HG2	1:C:698:VAL:HG23	1.94	0.49
1:C:619:VAL:O	1:C:623:GLU:HB2	2.13	0.49
1:C:660:TYR:O	1:C:662:PHE:N	2.46	0.49
3:L:58:ARG:HB3	3:L:73:THR:O	2.13	0.49
2:H:12:ALA:HB2	2:H:18:LEU:HD13	1.94	0.48
2:H:51:ILE:HD12	2:H:58:SER:OG	2.13	0.48
2:H:49:SER:HB3	2:H:60:TYR:CD1	2.49	0.48
2:H:105:SER:OG	5:H:403:HOH:O	2.20	0.47
1:C:660:TYR:C	1:C:662:PHE:H	2.16	0.47
1:C:589:TRP:O	1:C:590:GLU:HB3	2.13	0.47
3:L:120:PRO:HD3	3:L:132:LEU:CD2	2.45	0.47
1:C:650:LEU:H	1:C:650:LEU:HD22	1.80	0.47
3:L:198:GLU:OE1	5:L:401:HOH:O	2.21	0.47
3:L:51:ARG:HD2	5:L:439:HOH:O	2.15	0.47
1:C:620:LYS:HB3	1:C:620:LYS:HE3	1.44	0.46
2:H:155:GLY:HA2	2:H:170:TRP:CH2	2.50	0.46
1:C:657:GLN:HG3	2:H:104:PRO:HG2	1.97	0.46
1:C:580:ARG:O	1:C:584:ILE:HG13	2.14	0.46
1:C:663:ASN:O	1:C:668:LYS:NZ	2.47	0.46
2:H:30:ARG:O	2:H:53:ALA:HB1	2.16	0.46
2:H:57:MET:HE3	2:H:58:SER:H	1.80	0.46
1:C:660:TYR:CD2	2:H:107:THR:HG22	2.51	0.45
2:H:155:GLY:HA3	2:H:196:SER:O	2.16	0.45
3:L:58:ARG:HH21	3:L:79:ASP:CG	2.20	0.45
1:C:617:LYS:HE3	1:C:618:TRP:HB2	1.99	0.45
1:C:701:ASN:OD1	1:C:701:ASN:N	2.50	0.45
3:L:115:VAL:O	3:L:204:LYS:HE2	2.16	0.45
3:L:16:GLN:HG2	3:L:17:THR:N	2.32	0.44
3:L:156:LYS:HD2	3:L:156:LYS:HA	1.82	0.44
3:L:80:GLU:HG3	3:L:103:LEU:O	2.18	0.44
1:C:679:GLN:HG3	1:C:698:VAL:CG1	2.47	0.44
1:C:580:ARG:HG3	1:C:654:PHE:CD1	2.53	0.44
1:C:580:ARG:HG3	1:C:654:PHE:HD1	1.83	0.44
1:C:655:PHE:O	1:C:659:ILE:HG12	2.18	0.43
2:H:2:VAL:HA	2:H:26:GLY:HA3	2.00	0.43
2:H:98:LYS:NZ	2:H:117:ASP:OD2	2.52	0.43
2:H:161:TYR:CE2	2:H:166:VAL:HG13	2.54	0.42
3:L:130:ALA:HB3	3:L:180:LEU:HD12	2.01	0.42
2:H:53:ALA:HA	2:H:72:ARG:NH1	2.35	0.42
2:H:186:LEU:HD13	2:H:192:TYR:CE1	2.55	0.42
3:L:183:GLU:HB2	5:L:442:HOH:O	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:83:MET:HB2	2:H:86:LEU:HD21	2.02	0.41
3:L:159:VAL:HG22	3:L:178:LEU:HD13	2.02	0.41
2:H:87:THR:O	2:H:90:ASP:HB2	2.20	0.41
1:C:617:LYS:O	1:C:621:GLN:HG3	2.21	0.41
3:L:65:GLU:O	3:L:66:ASN:HB2	2.21	0.41
2:H:104:PRO:O	2:H:107:THR:OG1	2.24	0.41
3:L:132:LEU:HD12	3:L:178:LEU:HD23	2.03	0.41
2:H:35:THR:HB	2:H:50:TYR:CD2	2.56	0.40
3:L:167:GLN:HE21	3:L:173:ALA:HB2	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:802:ZN:ZN	4:C:802:ZN:ZN[2_576]	0.72	1.48

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	68/164 (42%)	65 (96%)	2 (3%)	1 (2%)	8	20
2	H	221/241 (92%)	211 (96%)	9 (4%)	1 (0%)	25	46
3	L	206/212 (97%)	196 (95%)	10 (5%)	0	100	100
All	All	495/617 (80%)	472 (95%)	21 (4%)	2 (0%)	30	52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	661	LYS
2	H	160	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	73/154 (47%)	64 (88%)	9 (12%)	4	8
2	H	186/202 (92%)	174 (94%)	12 (6%)	14	31
3	L	179/183 (98%)	172 (96%)	7 (4%)	27	52
All	All	438/539 (81%)	410 (94%)	28 (6%)	14	32

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	580	ARG
1	C	617	LYS
1	C	620	LYS
1	C	627	THR
1	C	662	PHE
1	C	697	LYS
1	C	700	PHE
1	C	701	ASN
1	C	709	ILE
2	H	51	ILE
2	H	52	THR
2	H	56	LYS
2	H	65	ARG
2	H	67	ARG
2	H	76	LYS
2	H	131	SER
2	H	136	SER
2	H	199	THR
2	H	203	SER
2	H	212	CYS
2	H	230	LYS
3	L	17	THR
3	L	44	VAL
3	L	90	SER
3	L	121	SER
3	L	165	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	L	179	SER
3	L	180	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	187	GLN
2	H	215	ASN
3	L	108	GLN
3	L	167	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	76/164 (46%)	2.04	34 (44%) <b>1</b> <b>1</b>	29, 62, 90, 95	0
2	H	225/241 (93%)	0.21	8 (3%) 46 45	11, 29, 49, 64	0
3	L	208/212 (98%)	-0.05	4 (1%) 66 65	9, 24, 41, 56	0
All	All	509/617 (82%)	0.37	46 (9%) <b>17</b> <b>15</b>	9, 28, 75, 95	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	709	ILE	8.0
1	C	589	TRP	5.7
1	C	680	ILE	5.0
1	C	627	THR	4.4
1	C	650	LEU	4.2
1	C	697	LYS	4.2
1	C	699	LEU	4.1
1	C	707	ALA	3.9
1	C	703	ILE	3.9
1	C	626	TRP	3.9
1	C	649	ASP	3.8
1	C	614	CYS	3.8
1	C	618	TRP	3.7
3	L	3	GLU	3.6
1	C	651	PHE	3.4
1	C	615	PHE	3.1
1	C	700	PHE	3.1
3	L	209	THR	3.1
3	L	78	LEU	3.1
1	C	676	LEU	3.0
1	C	575	PHE	3.0
2	H	149	GLY	2.9
1	C	578	TRP	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	620	LYS	2.7
1	C	679	GLN	2.7
3	L	2	SER	2.7
1	C	698	VAL	2.6
1	C	708	THR	2.5
1	C	617	LYS	2.5
2	H	105	SER	2.5
1	C	619	VAL	2.4
1	C	616	ASP	2.4
2	H	129	SER	2.4
2	H	62	ASP	2.4
1	C	577	PHE	2.4
1	C	706	ILE	2.4
1	C	585	ASP	2.4
2	H	103	CYS	2.4
2	H	224	ASP	2.4
2	H	33	GLY	2.3
1	C	662	PHE	2.3
1	C	581	LYS	2.2
1	C	705	GLU	2.1
2	H	30	ARG	2.1
1	C	590	GLU	2.0
1	C	648	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

4	ZN	C	802	1/1	0.87	0.19	140,140,140,140	0
4	ZN	H	303	1/1	0.93	0.07	82,82,82,82	0
4	ZN	H	301	1/1	0.96	0.10	76,76,76,76	0
4	ZN	C	801	1/1	0.96	0.06	86,86,86,86	0
4	ZN	L	301	1/1	0.97	0.11	54,54,54,54	0
4	ZN	H	302	1/1	0.99	0.02	40,40,40,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.