

Full wwPDB X-ray Structure Validation Report (i)

Sep 25, 2023 – 04:16 AM EDT

PDB ID	:	5VEW
Title	:	Structure of the human GLP-1 receptor complex with PF-06372222
Authors	:	Song, G.; Yang, D.; Wang, Y.; Graaf, C.D.; Zhou, Q.; Jiang, S.; Liu, K.; Cai,
		X.; Dai, A.; Lin, G.; Liu, D.; Wu, F.; Wu, Y.; Zhao, S.; Ye, L.; Han, G.W.;
		Lau, J.; Wu, B.; Hanson, M.A.; Liu, ZJ.; Wang, MW.; Stevens, R.C.
Deposited on	:	2017-04-05
Resolution	:	2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	455	8%	10%	7%
1	В	455	9%	10%	7%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6849 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	493	Total	С	Ν	0	\mathbf{S}	0	0	0
	420	3302	2176	544	564	18	0	0	0	
1	Р	494	Total	С	Ν	0	S	0	0	0
	424	3305	2176	548	563	18	0	0	0	

• Molecule 1 is a protein called Glucagon-like peptide 1 receptor, Endolysin chimera.

Chain	Residue	Modelled	Actual	Comment	Reference
А	127	SER	-	expression tag	UNP P43220
А	193	CYS	SER	engineered mutation	UNP P43220
A	196	PHE	ILE	engineered mutation	UNP P43220
А	212	GLY	TYR	linker	UNP P43220
А	?	-	THR	deletion	UNP P43220
А	?	-	ALA	deletion	UNP P43220
А	?	-	ALA	deletion	UNP P43220
А	?	-	GLN	deletion	UNP P43220
А	?	-	GLN	deletion	UNP P43220
А	?	-	HIS	deletion	UNP P43220
А	?	-	GLN	deletion	UNP P43220
А	214	GLY	TRP	linker	UNP P43220
А	225	ALA	SER	engineered mutation	UNP P43220
A	233	CSD	MET	engineered mutation	UNP P43220
А	1012	GLY	ARG	engineered mutation	UNP P00720
А	1054	THR	CYS	engineered mutation	UNP P00720
А	1097	ALA	CYS	engineered mutation	UNP P00720
А	1137	ARG	ILE	engineered mutation	UNP P00720
A	271	ALA	SER	engineered mutation	UNP P43220
А	317	CYS	ILE	engineered mutation	UNP P43220
А	318	ILE	GLY	engineered mutation	UNP P43220
А	346	ALA	LYS	engineered mutation	UNP P43220
А	347	PHE	CYS	engineered mutation	UNP P43220
А	361	CYS	GLY	engineered mutation	UNP P43220
В	127	SER	-	expression tag	UNP P43220

There are 48 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
В	193	CYS	SER	engineered mutation	UNP P43220
В	196	PHE	ILE	engineered mutation	UNP P43220
В	212	GLY	TYR	linker	UNP P43220
В	?	-	THR	deletion	UNP P43220
В	?	-	ALA	deletion	UNP P43220
В	?	-	ALA	deletion	UNP P43220
В	?	-	GLN	deletion	UNP P43220
В	?	-	GLN	deletion	UNP P43220
В	?	-	HIS	deletion	UNP P43220
В	?	-	GLN	deletion	UNP P43220
В	214	GLY	TRP	linker	UNP P43220
В	225	ALA	SER	engineered mutation	UNP P43220
В	233	CSD	MET	engineered mutation	UNP P43220
В	1012	GLY	ARG	engineered mutation	UNP P00720
В	1054	THR	CYS	engineered mutation	UNP P00720
В	1097	ALA	CYS	engineered mutation	UNP P00720
В	1137	ARG	ILE	engineered mutation	UNP P00720
В	271	ALA	SER	engineered mutation	UNP P43220
В	317	CYS	ILE	engineered mutation	UNP P43220
В	318	ILE	GLY	engineered mutation	UNP P43220
В	346	ALA	LYS	engineered mutation	UNP P43220
В	347	PHE	CYS	engineered mutation	UNP P43220
В	361	CYS	GLY	engineered mutation	UNP P43220

• Molecule 2 is N-{4-[(R)-(3,3-dimethylcyclobutyl)({6-[4-(trifluoromethyl)-1H-imidazol-1 -yl]pyridin-3-yl}amino)methyl]benzene-1-carbonyl}-beta-alanine (three-letter code: 97Y) (formula: $C_{26}H_{28}F_3N_5O_3$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
0	٨	1	Total	С	F	Ν	Ο	0	0	
	1	37	26	3	5	3	0	0		
0	D	В	1	Total	С	F	Ν	Ο	0	0
2 B	1	37	26	3	5	3	0	0		

• Molecule 3 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C O 25 21 4	0	0
3	А	1	Total C O 19 15 4	0	0
3	В	1	Total C O 25 21 4	0	0
3	В	1	Total C O 25 21 4	0	0
3	В	1	Total C O 19 15 4	0	0
3	В	1	Total C O 15 11 4	0	0

• Molecule 4 is OLEIC ACID (three-letter code: OLA) (formula: $C_{18}H_{34}O_2$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	А	1	Total 20	C 18	O 2	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	13	Total O 13 13	0	0
5	В	7	Total O 7 7	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Glucagon-like peptide 1 receptor, Endolysin chimera

 3%
 1330
 3%

 1345
 1345
 1003
 4%

 1345
 1003
 1003
 1003

 1446
 1003
 1003
 1003

 1357
 10040
 1003
 100

 1358
 10040
 1008
 100

 1357
 1005
 1005
 1005

 1358
 1006
 1006
 1006

 1000
 1006
 1006
 1006

 1000
 1006
 1006
 1047

 1000
 1006
 10147
 10147

 1000
 1006
 1147
 1147

 1000
 1147
 1147
 1147

 1000
 1147
 1147
 1147

 1000
 1147
 1147
 1147

 1000
 1147
 1147
 1147

 1001
 1147
 1147
 1147

 1001
 1147
 1147
 1147

 1001
 1147
 1147
 1147

 1001
 1147
 1147
 1147

 100101
 1147
 1147
 <

• Molecule 1: Glucagon-like peptide 1 receptor, Endolysin chimera



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	64.77Å 66.43 Å 83.44 Å	Deperitor
a, b, c, α , β , γ	90.54° 90.18° 107.73°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	30.00 - 2.70	Depositor
Resolution (A)	29.73 - 2.70	EDS
% Data completeness	95.2 (30.00-2.70)	Depositor
(in resolution range)	95.2 (29.73-2.70)	EDS
R _{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.59 (at 2.68 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
B B.	0.228 , 0.246	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.261 , 0.276	DCC
R_{free} test set	1743 reflections $(5.04%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	75.9	Xtriage
Anisotropy	0.352	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.28, 76.1	EDS
L-test for twinning ²	$< L > = 0.48, < L^2 > = 0.31$	Xtriage
	0.029 for -h,-k,l	
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
	0.000 for -k,-h,-l	
F_o, F_c correlation	0.93	EDS
Total number of atoms	6849	wwPDB-VP
Average B, all atoms $(Å^2)$	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 13.40% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 97Y, OLA, CSD, OLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.51	0/3371	0.55	0/4590
1	В	0.51	0/3374	0.57	0/4593
All	All	0.51	0/6745	0.56	0/9183

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3302	0	3250	27	0
1	В	3305	0	3251	29	0
2	А	37	0	0	1	0
2	В	37	0	0	1	0
3	А	44	0	65	1	0
3	В	84	0	124	2	0
4	А	20	0	33	1	0
5	А	13	0	0	0	0
5	В	7	0	0	0	0
All	All	6849	0	6723	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:303:MET:HG2	1:B:304:ASN:H	1.45	0.79
1:A:1050:ILE:HD11	1:A:1062:GLU:HB3	1.73	0.69
1:B:1050:ILE:HD11	1:B:1062:GLU:HB3	1.74	0.69
1:B:387:GLU:HG3	3:B:1203:OLC:H17A	1.79	0.64
1:B:360:LEU:HG	1:B:394:GLN:HE21	1.67	0.60
1:B:288:LYS:HE3	1:B:297:TRP:HB3	1.83	0.59
1:A:403:CYS:O	1:A:409:VAL:HG21	2.06	0.55
1:A:390:PHE:O	1:A:394:GLN:HG3	2.06	0.54
1:B:403:CYS:O	1:B:409:VAL:HG21	2.08	0.54
1:A:292:GLU:OE2	1:A:302:ASN:HB3	2.09	0.52
1:B:227:ARG:HG2	1:B:296:CYS:HA	1.92	0.52
1:B:415:LYS:O	1:B:419:ARG:HG2	2.11	0.51
1:A:415:LYS:O	1:A:419:ARG:HG2	2.11	0.50
1:A:1095:ARG:HG2	1:A:1153:PHE:HA	1.94	0.50
1:B:1095:ARG:HG2	1:B:1153:PHE:HA	1.94	0.50
1:A:396:LEU:O	1:A:400:ILE:HG12	2.12	0.49
1:B:1034:THR:HG22	1:B:1036:SER:H	1.77	0.49
1:B:292:GLU:OE2	1:B:302:ASN:HB3	2.13	0.48
1:A:167:LEU:HD22	4:A:1204:OLA:H42	1.96	0.47
1:B:396:LEU:O	1:B:400:ILE:HG12	2.14	0.47
1:A:288:LYS:CE	1:A:297:TRP:HB3	2.44	0.47
1:B:1095:ARG:HD3	1:B:1153:PHE:O	2.16	0.46
1:A:1095:ARG:HD3	1:A:1153:PHE:O	2.16	0.46
1:B:362:THR:CG2	3:B:1203:OLC:H12	2.45	0.46
1:A:348:ARG:HB3	2:A:1201:97Y:C36	2.45	0.46
1:B:348:ARG:HB3	2:B:1202:97Y:C36	2.46	0.46
1:A:181:LEU:HB3	3:A:1202:OLC:H4	1.98	0.46
1:B:1103:VAL:HG22	1:B:1111:VAL:HG21	1.98	0.46
1:A:1103:VAL:HG22	1:A:1111:VAL:HG21	1.99	0.45
1:A:227:ARG:HG2	1:A:296:CYS:HA	1.99	0.44
1:B:288:LYS:O	1:B:297:TRP:CE2	2.71	0.44
1:A:326:ARG:O	1:A:330:ILE:HG12	2.18	0.43
1:A:169:PHE:HB2	1:A:172:LEU:HD12	2.01	0.42
1:A:280:PHE:C	1:A:283:PRO:HD2	2.40	0.42
1:B:326:ARG:O	1:B:330:ILE:HG12	2.19	0.42
1:A:345:ILE:HG13	1:A:346:ALA:N	2.35	0.42
1:B:288:LYS:HD2	1:B:292:GLU:HB2	2.01	0.42
1:A:288:LYS:HE3	1:A:297:TRP:HB3	2.02	0.42



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:227:ARG:HB3	1:B:297:TRP:CZ2	2.55	0.42
1:B:169:PHE:HB2	1:B:172:LEU:HD12	2.02	0.42
1:B:1087:VAL:O	1:B:1091:LEU:HG	2.20	0.41
1:A:1087:VAL:O	1:A:1091:LEU:HG	2.20	0.41
1:A:245:LEU:HD13	1:A:319:VAL:HG23	2.02	0.41
1:A:1137:ARG:H	1:A:1137:ARG:HD3	1.84	0.41
1:B:345:ILE:HG13	1:B:346:ALA:N	2.36	0.41
1:A:320:ASN:HB3	1:A:357:ILE:HG23	2.02	0.41
1:A:319:VAL:O	1:A:323:ILE:HG12	2.21	0.41
1:B:245:LEU:HD13	1:B:319:VAL:HG23	2.03	0.41
1:B:189:LEU:HB3	1:B:236:CYS:SG	2.60	0.41
1:A:288:LYS:HE2	1:A:297:TRP:HB3	2.03	0.41
1:B:319:VAL:O	1:B:323:ILE:HG12	2.21	0.41
1:B:230:PHE:CE2	1:B:284:TRP:HH2	2.39	0.41
1:B:1137:ARG:H	1:B:1137:ARG:HD3	1.85	0.41
1:A:189:LEU:HA	1:A:192:LEU:HD12	2.02	0.40
1:B:320:ASN:HB3	1:B:357:ILE:HG23	2.03	0.40
1:A:1022:GLU:HB2	1:A:1024:TYR:HD1	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	416/455 (91%)	392 (94%)	24~(6%)	0	100 100
1	В	417/455~(92%)	391 (94%)	26~(6%)	0	100 100
All	All	833/910~(92%)	783~(94%)	50~(6%)	0	100 100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	330/394~(84%)	326~(99%)	4 (1%)	71 88		
1	В	329/394~(84%)	326~(99%)	3 (1%)	78 92		
All	All	659/788~(84%)	652~(99%)	7(1%)	73 90		

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	180	HIS
1	А	1068	ASN
1	А	1137	ARG
1	А	397	MET
1	В	180	HIS
1	В	1068	ASN
1	В	1137	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	240	ASN
1	А	1068	ASN
1	В	240	ASN
1	В	1068	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul



statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tink	B	ond leng	gths	I	Bond an	gles
INIOI	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
1	CSD	А	233	1	3,7,8	0.79	0	1,8,10	4.07	1 (100%)
1	CSD	В	233	1	3,7,8	0.80	0	1,8,10	3.72	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	А	233	1	-	1/2/6/8	-
1	CSD	В	233	1	-	2/2/6/8	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	233	CSD	OD1-SG-CB	4.07	113.29	105.54
1	В	233	CSD	OD1-SG-CB	3.72	112.61	105.54

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	233	CSD	CA-CB-SG-OD1
1	В	233	CSD	N-CA-CB-SG
1	В	233	CSD	CA-CB-SG-OD1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trune	Chain	Dag	Tinle	Bond lengths		Bond angles			
IVIOI	туре	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	OLC	А	1203	-	18,18,24	1.15	1 (5%)	18,19,25	0.93	1 (5%)
3	OLC	В	1205	-	14,14,24	1.27	1 (7%)	15,15,25	1.12	2 (13%)
3	OLC	В	1203	-	24,24,24	1.00	1 (4%)	25,25,25	0.82	1 (4%)
2	97Y	А	1201	-	39,40,40	2.17	10 (25%)	50,59,59	1.79	10 (20%)
3	OLC	А	1202	-	$24,\!24,\!24$	1.02	1 (4%)	25,25,25	0.94	2 (8%)
2	97Y	В	1202	-	39,40,40	2.17	10 (25%)	50,59,59	1.79	10 (20%)
3	OLC	В	1201	-	24,24,24	0.99	1 (4%)	25,25,25	0.91	2 (8%)
3	OLC	В	1204	-	18,18,24	1.10	1 (5%)	18,19,25	0.88	1 (5%)
4	OLA	А	1204	-	$19,\!19,\!19$	0.49	0	19,19,19	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLC	А	1203	-	-	10/18/18/24	-
3	OLC	В	1205	-	-	10/14/14/24	-
3	OLC	В	1203	-	-	12/24/24/24	-
2	97Y	А	1201	-	-	3/30/42/42	0/4/4/4
3	OLC	А	1202	-	-	12/24/24/24	-
2	97Y	В	1202	-	-	3/30/42/42	0/4/4/4
3	OLC	В	1201	-	-	14/24/24/24	-
3	OLC	В	1204	-	-	4/18/18/24	-
4	OLA	А	1204	-	-	15/17/17/17	-

All (26) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(\text{\AA})$	Ideal(Å)
2	А	1201	97Y	C22-N07	-6.15	1.32	1.39
2	В	1202	97Y	C22-N07	-6.03	1.32	1.39
2	В	1202	97Y	C34-N11	5.81	1.46	1.33
2	А	1201	97Y	C34-N11	5.67	1.46	1.33
3	А	1202	OLC	O20-C1	4.62	1.46	1.33
3	А	1203	OLC	O20-C1	4.60	1.46	1.33
3	В	1203	OLC	O20-C1	4.59	1.46	1.33
3	В	1205	OLC	O20-C1	4.51	1.46	1.33
3	В	1201	OLC	O20-C1	4.48	1.46	1.33
3	В	1204	OLC	O20-C1	4.32	1.46	1.33
2	А	1201	97Y	C20-N08	3.72	1.46	1.39
2	В	1202	97Y	C20-N08	3.68	1.46	1.39
2	А	1201	97Y	C15-C12	-3.48	1.50	1.55
2	В	1202	97Y	C15-C12	-3.47	1.51	1.55
2	А	1201	97Y	C23-N07	-3.31	1.33	1.36
2	В	1202	97Y	C23-N07	-3.30	1.33	1.36
2	В	1202	97Y	C12-C13	-3.15	1.48	1.53
2	А	1201	97Y	C12-C13	-3.15	1.48	1.53
2	А	1201	97Y	C16-C14	-3.05	1.51	1.55
2	В	1202	97Y	C16-C14	-2.93	1.51	1.55
2	В	1202	97Y	C17-C13	2.91	1.55	1.52
2	В	1202	97Y	C16-C12	-2.87	1.51	1.55
2	А	1201	97Y	C16-C12	-2.84	1.51	1.55
2	А	1201	97Y	C17-C13	2.81	1.55	1.52
2	В	1202	97Y	C15-C14	-2.46	1.51	1.55
2	А	1201	97Y	C15-C14	-2.37	1.52	1.55

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	1202	97Y	C21-C18-N09	6.21	127.07	119.76
2	А	1201	97Y	C21-C18-N09	6.12	126.96	119.76
2	В	1202	97Y	C17-C13-C12	-4.96	107.26	112.15
2	А	1201	97Y	C17-C13-C12	-4.92	107.30	112.15
2	А	1201	97Y	C24-C19-N10	-4.61	119.53	125.48
2	В	1202	97Y	C24-C19-N10	-4.48	119.70	125.48
2	А	1201	97Y	C28-C24-C19	3.20	120.94	116.66
3	А	1202	OLC	O20-C1-C2	2.94	121.13	111.91
2	В	1202	97Y	C28-C24-C19	2.93	120.58	116.66
3	В	1205	OLC	O20-C1-C2	2.88	120.95	111.91
3	В	1201	OLC	O20-C1-C2	2.80	120.71	111.91
2	В	1202	97Y	C20-C27-N10	-2.76	121.65	124.13
3	А	1203	OLC	O20-C1-C2	2.71	120.43	111.91



Mol	Chain	Res	Type	Atoms		$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	А	1201	97Y	C24-C28-C20	-2.63	117.25	120.30
3	В	1203	OLC	O20-C1-C2	2.58	120.00	111.91
2	А	1201	97Y	C20-C27-N10	-2.50	121.89	124.13
2	В	1202	97Y	C24-C28-C20	-2.50	117.41	120.30
3	В	1204	OLC	O20-C1-C2	2.43	119.52	111.91
2	В	1202	97Y	C27-N10-C19	2.39	121.61	117.29
2	В	1202	97Y	F02-C21-C18	-2.39	108.39	112.47
2	В	1202	97Y	O05-C37-C36	2.32	121.47	114.03
2	А	1201	97Y	C27-N10-C19	2.29	121.42	117.29
3	В	1205	OLC	O20-C1-O19	-2.28	117.84	123.59
2	А	1201	97Y	O05-C37-C36	2.28	121.35	114.03
3	В	1201	OLC	O20-C1-O19	-2.26	117.89	123.59
3	А	1202	OLC	O20-C1-O19	-2.24	117.95	123.59
2	В	1202	97Y	F03-C21-C18	-2.10	108.88	112.47
2	A	1201	97Y	F03-C21-C18	-2.10	108.89	112.47
2	A	1201	97Y	F02-C21-C18	-2.04	108.98	112.47

There are no chirality outliers.

All	(83)) torsion	outliers	are	listed	below:
-----	------	-----------	----------	-----	--------	--------

Mol	Chain	\mathbf{Res}	Type	Atoms
3	А	1202	OLC	C21-C22-C24-O25
3	А	1202	OLC	C2-C1-O20-C21
3	А	1202	OLC	O19-C1-O20-C21
3	А	1203	OLC	C9-C10-C11-C12
3	А	1203	OLC	O20-C21-C22-C24
3	В	1205	OLC	C21-C22-C24-O25
3	В	1201	OLC	O19-C1-O20-C21
3	В	1205	OLC	O19-C1-O20-C21
3	А	1203	OLC	C2-C1-O20-C21
3	В	1205	OLC	C2-C1-O20-C21
3	В	1201	OLC	C2-C1-O20-C21
3	В	1204	OLC	C2-C1-O20-C21
4	А	1204	OLA	С11-С10-С9-С8
3	А	1203	OLC	O19-C1-O20-C21
3	А	1203	OLC	O20-C21-C22-O23
3	В	1204	OLC	O19-C1-O20-C21
4	А	1204	OLA	C1-C2-C3-C4
3	В	1203	OLC	C2-C1-O20-C21
4	А	1204	OLA	C10-C11-C12-C13
3	В	1203	OLC	C12-C13-C14-C15
3	А	1203	OLC	C2-C3-C4-C5



00.000	itaea jien	t procee	, ao page.	
Mol	Chain	Res	Type	Atoms
3	A	1202	OLC	C13-C14-C15-C16
3	А	1202	OLC	C12-C13-C14-C15
4	А	1204	OLA	C2-C3-C4-C5
4	А	1204	OLA	C11-C12-C13-C14
3	А	1202	OLC	C14-C15-C16-C17
3	В	1203	OLC	C4-C5-C6-C7
3	В	1203	OLC	C2-C3-C4-C5
4	А	1204	OLA	C4-C5-C6-C7
3	В	1203	OLC	C5-C6-C7-C8
3	А	1203	OLC	C21-C22-C24-O25
3	В	1201	OLC	C6-C7-C8-C9
4	А	1204	OLA	C6-C7-C8-C9
3	В	1205	OLC	C3-C4-C5-C6
3	В	1201	OLC	C2-C3-C4-C5
4	А	1204	OLA	C5-C6-C7-C8
3	В	1201	OLC	C3-C4-C5-C6
4	А	1204	OLA	C12-C13-C14-C15
3	В	1201	OLC	C14-C15-C16-C17
3	В	1203	OLC	O19-C1-O20-C21
3	А	1202	OLC	O23-C22-C24-O25
3	А	1203	OLC	O23-C22-C24-O25
3	А	1202	OLC	C2-C3-C4-C5
3	В	1201	OLC	C11-C12-C13-C14
3	А	1203	OLC	C4-C5-C6-C7
4	А	1204	OLA	C13-C14-C15-C16
3	А	1202	OLC	C10-C11-C12-C13
3	В	1201	OLC	C10-C11-C12-C13
3	В	1201	OLC	C1-C2-C3-C4
3	А	1202	OLC	C3-C4-C5-C6
4	А	1204	OLA	C3-C4-C5-C6
3	В	1205	OLC	C5-C6-C7-C8
3	В	1205	OLC	O23-C22-C24-O25
3	В	1201	OLC	C15-C16-C17-C18
3	В	1203	OLC	C15-C16-C17-C18
4	А	1204	OLA	C15-C16-C17-C18
3	В	1201	OLC	C12-C13-C14-C15
3	В	1201	OLC	C4-C5-C6-C7
3	В	1201	OLC	C13-C14-C15-C16
2	А	1201	97Y	N11-C35-C36-C37
2	В	1202	97Y	N11-C35-C36-C37
3	А	1202	OLC	C11-C12-C13-C14
3	А	1202	OLC	C4-C5-C6-C7

Continued from previous page...



Conti	nued fron	n previo	ous page.	
Mol	Chain	Res	Type	Atoms
3	В	1203	OLC	O20-C21-C22-O23
3	В	1203	OLC	C13-C14-C15-C16
3	В	1205	OLC	C4-C5-C6-C7
3	В	1203	OLC	O20-C21-C22-C24
3	В	1203	OLC	C11-C12-C13-C14
3	В	1204	OLC	C2-C3-C4-C5
4	А	1204	OLA	O2-C1-C2-C3
3	В	1204	OLC	C4-C5-C6-C7
4	А	1204	OLA	O1-C1-C2-C3
4	А	1204	OLA	C7-C8-C9-C10
2	В	1202	97Y	C35-C36-C37-O06
3	В	1203	OLC	C14-C15-C16-C17
3	В	1205	OLC	C2-C3-C4-C5
3	В	1201	OLC	C9-C10-C11-C12
2	А	1201	97Y	C35-C36-C37-O05
2	А	1201	97Y	C35-C36-C37-O06
2	В	1202	97Y	C35-C36-C37-O05
3	А	1203	OLC	C3-C4-C5-C6
3	В	1205	OLC	O20-C1-C2-C3
3	В	1205	OLC	O19-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	1203	OLC	2	0
2	А	1201	97Y	1	0
3	А	1202	OLC	1	0
2	В	1202	97Y	1	0
4	А	1204	OLA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient





equivalents in the CSD to analyse the geometry.













5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	422/455~(92%)	0.54	38 (9%) 9 7	63, 99, 159, 185	0
1	В	423/455~(92%)	0.51	39 (9%) 9 7	63, 96, 158, 185	0
All	All	845/910~(92%)	0.52	77 (9%) 9 7	63, 97, 159, 185	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	1037	PRO	12.3
1	А	218	LEU	8.6
1	А	302	ASN	7.9
1	А	1037	PRO	6.2
1	В	422	LEU	6.0
1	А	1040	ASN	5.9
1	В	216	GLY	5.9
1	В	1038	SER	5.8
1	В	302	ASN	5.6
1	В	217	LEU	5.3
1	А	1038	SER	4.6
1	В	1040	ASN	4.4
1	А	419	ARG	4.3
1	А	418	GLU	4.2
1	В	414	ARG	3.8
1	В	140	GLN	3.7
1	А	301	SER	3.6
1	А	416	SER	3.6
1	В	380	ARG	3.4
1	A	219	SER	3.4
1	A	220	TYR	3.4
1	А	1061	ASP	3.3
1	A	1057	VAL	3.3
1	А	1025	TYR	3.3



r
7

Mol	Chain	Res	Type	RSRZ
1	В	196	PHE	3.3
1	В	144	LEU	3.3
1	В	305	TYR	3.2
1	А	397	MET	3.2
1	В	419	ARG	3.1
1	В	1057	VAL	3.0
1	А	1024	TYR	3.0
1	А	306	TRP	2.9
1	А	382	ILE	2.9
1	А	414	ARG	2.9
1	В	1025	TYR	2.9
1	А	1015	LEU	2.8
1	В	397	MET	2.7
1	А	417	TRP	2.7
1	В	1061	ASP	2.7
1	А	390	PHE	2.7
1	А	1034	THR	2.7
1	А	415	LYS	2.7
1	В	147	ILE	2.6
1	А	1041	ALA	2.5
1	В	224	LEU	2.5
1	В	415	LYS	2.5
1	В	381	PHE	2.5
1	А	1039	LEU	2.4
1	В	139	GLU	2.4
1	А	1060	LYS	2.4
1	А	1058	ILE	2.4
1	А	380	ARG	2.3
1	В	1058	ILE	2.3
1	А	420	TRP	2.3
1	В	1015	LEU	2.3
1	В	137	PRO	2.3
1	В	300	ASN	2.3
1	A	1013	LEU	2.3
1	В	145	TYR	2.3
1	В	301	SER	2.3
1	А	140	GLN	2.2
1	A	200	ALA	2.2
1	А	237	VAL	2.2
1	В	1050	ILE	2.2
1	В	1119	ARG	2.2
1	В	420	TRP	2.2



		1	1 0	
Mol	Chain	\mathbf{Res}	Type	RSRZ
1	А	196	PHE	2.1
1	В	136	SER	2.1
1	А	222	ASP	2.1
1	В	218	LEU	2.1
1	В	1144	ASN	2.1
1	В	390	PHE	2.1
1	А	383	LYS	2.1
1	В	1060	LYS	2.1
1	В	220	TYR	2.0
1	А	1042	ALA	2.0
1	В	199	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
1	CSD	В	233	8/9	0.84	0.23	86,91,97,97	0
1	CSD	А	233	8/9	0.87	0.26	93,95,101,101	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
3	OLC	В	1203	25/25	0.65	0.31	95,99,107,107	0
3	OLC	В	1205	15/25	0.65	0.20	137,141,143,143	0
4	OLA	А	1204	20/20	0.69	0.37	131,135,140,141	0
3	OLC	А	1203	19/25	0.74	0.16	142,145,152,153	0
3	OLC	В	1201	25/25	0.75	0.29	104,110,113,114	0



e ontontaca front procesa ao pagoni									
Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9	
3	OLC	А	1202	25/25	0.81	0.23	109,112,115,116	0	
3	OLC	В	1204	19/25	0.83	0.18	114,115,118,120	0	
2	97Y	А	1201	37/37	0.95	0.18	83,95,108,109	0	
2	97Y	В	1202	37/37	0.96	0.19	74,89,105,106	0	

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.























6.5 Other polymers (i)

There are no such residues in this entry.

