



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 25, 2023 – 12:43 PM EDT

PDB ID : 5VEX  
Title : Structure of the human GLP-1 receptor complex with NNC0640  
Authors : Song, G.; Yang, D.; Wang, Y.; Graaf, C.D.; Zhou, Q.; Jiang, S.; Liu, K.; Cai, X.; Dai, A.; Lin, G.; Liu, D.; Wu, F.; Wu, Y.; Zhao, S.; Ye, L.; Han, G.W.; Lau, J.; Wu, B.; Hanson, M.A.; Liu, Z.-J.; Wang, M.-W.; Stevens, R.C.  
Deposited on : 2017-04-05  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	455	 2% 87% 6% 7%
1	B	455	 3% 85% 7% 7%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 6707 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucagon-like peptide 1 receptor, Endolysin chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	423	3309	2183	542	566	18	0	0	0
1	B	423	3316	2185	546	567	18	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	SER	-	expression tag	UNP P43220
A	193	CYS	SER	engineered mutation	UNP P43220
A	196	PHE	ILE	engineered mutation	UNP P43220
A	212	GLY	TYR	linker	UNP P43220
A	?	-	THR	deletion	UNP P43220
A	?	-	ALA	deletion	UNP P43220
A	?	-	ALA	deletion	UNP P43220
A	?	-	GLN	deletion	UNP P43220
A	?	-	GLN	deletion	UNP P43220
A	?	-	HIS	deletion	UNP P43220
A	?	-	GLN	deletion	UNP P43220
A	214	GLY	TRP	linker	UNP P43220
A	225	ALA	SER	engineered mutation	UNP P43220
A	233	CSD	MET	engineered mutation	UNP P43220
A	1012	GLY	ARG	engineered mutation	UNP P00720
A	1054	THR	CYS	engineered mutation	UNP P00720
A	1097	ALA	CYS	engineered mutation	UNP P00720
A	1137	ARG	ILE	engineered mutation	UNP P00720
A	271	ALA	SER	engineered mutation	UNP P43220
A	317	CYS	ILE	engineered mutation	UNP P43220
A	318	ILE	GLY	engineered mutation	UNP P43220
A	346	ALA	LYS	engineered mutation	UNP P43220
A	347	PHE	CYS	engineered mutation	UNP P43220
A	361	CYS	GLY	engineered mutation	UNP P43220
B	127	SER	-	expression tag	UNP P43220

*Continued on next page...*




<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>					<b>ZeroOcc</b>	<b>AltConf</b>
2	A	1	Total 41	C 29	N 7	O 4	S 1	0	0
2	B	1	Total 41	C 29	N 7	O 4	S 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glucagon-like peptide 1 receptor, Endolysin chimera



- Molecule 1: Glucagon-like peptide 1 receptor, Endolysin chimera



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.81Å 67.51Å 83.66Å 91.55° 89.88° 107.56°	Depositor
Resolution (Å)	30.00 – 3.00 29.84 – 3.00	Depositor EDS
% Data completeness (in resolution range)	91.4 (30.00-3.00) 91.4 (29.84-3.00)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 3.00Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.233 , 0.256 0.259 , 0.289	Depositor DCC
$R_{free}$ test set	1060 reflections (4.29%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	85.1	Xtriage
Anisotropy	0.710	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 77.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,l 0.000 for -k,-h,-l 0.000 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6707	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	111.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 97V, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/3380	0.53	0/4604
1	B	0.50	0/3387	0.54	0/4612
All	All	0.50	0/6767	0.54	0/9216

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3309	0	3251	12	0
1	B	3316	0	3259	17	0
2	A	41	0	0	0	0
2	B	41	0	0	0	0
All	All	6707	0	6510	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:227:ARG:HG2	1:B:296:CYS:HA	1.90	0.53
1:A:227:ARG:HG2	1:A:296:CYS:HA	1.91	0.53
1:B:293:ASP:HA	1:B:297:TRP:CD1	2.44	0.52
1:A:320:ASN:HB3	1:A:357:ILE:HG23	1.93	0.50
1:B:320:ASN:HB3	1:B:357:ILE:HG23	1.93	0.50
1:A:1005:GLU:O	1:A:1009:ILE:HG12	2.13	0.49
1:B:170:ARG:HA	1:B:173:HIS:HD2	1.78	0.48
1:A:170:ARG:HA	1:A:173:HIS:HD2	1.78	0.48
1:B:180:HIS:CD2	1:B:402:TYR:HB3	2.50	0.47
1:B:396:LEU:O	1:B:400:ILE:HG12	2.16	0.46
1:A:396:LEU:O	1:A:400:ILE:HG12	2.16	0.46
1:B:1034:THR:HG22	1:B:1036:SER:H	1.81	0.45
1:B:262:GLU:HB2	1:B:265:ILE:HG12	1.99	0.44
1:A:282:VAL:CG2	1:A:283:PRO:HD3	2.47	0.44
1:B:282:VAL:CG2	1:B:283:PRO:HD3	2.47	0.43
1:B:277:PRO:HA	1:B:280:PHE:CE2	2.53	0.43
1:A:1078:ILE:HG23	1:A:1084:LEU:HB3	2.01	0.43
1:A:1103:VAL:HG22	1:A:1111:VAL:HG21	2.01	0.42
1:B:1078:ILE:HG23	1:B:1084:LEU:HB3	2.01	0.42
1:B:1005:GLU:O	1:B:1009:ILE:HG12	2.20	0.42
1:B:1103:VAL:HG22	1:B:1111:VAL:HG21	2.02	0.42
1:B:243:TRP:CZ2	1:B:270:VAL:HA	2.56	0.41
1:A:1126:TRP:HB3	1:A:1154:ARG:HA	2.03	0.41
1:A:293:ASP:HA	1:A:297:TRP:CD1	2.56	0.41
1:B:1126:TRP:HB3	1:B:1154:ARG:HA	2.03	0.41
1:A:147:ILE:HD13	1:A:389:SER:HB3	2.04	0.40
1:B:147:ILE:HD13	1:B:389:SER:HB3	2.04	0.40
1:B:288:LYS:CE	1:B:297:TRP:HB3	2.52	0.40
1:A:287:VAL:HG21	1:A:308:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/455 (91%)	393 (94%)	23 (6%)	0	100	100
1	B	416/455 (91%)	394 (95%)	22 (5%)	0	100	100
All	All	832/910 (91%)	787 (95%)	45 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/394 (84%)	330 (100%)	1 (0%)	92	97
1	B	332/394 (84%)	330 (99%)	2 (1%)	86	95
All	All	663/788 (84%)	660 (100%)	3 (0%)	88	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1127	ASP
1	B	180	HIS
1	B	1127	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	173	HIS
1	A	1068	ASN
1	A	1141	GLN
1	B	173	HIS
1	B	1068	ASN
1	B	304	ASN
1	B	407	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSD	A	233	1	3,7,8	0.87	0	1,8,10	3.75	1 (100%)
1	CSD	B	233	1	3,7,8	0.92	0	1,8,10	3.93	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	233	1	-	2/2/6/8	-
1	CSD	B	233	1	-	2/2/6/8	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	233	CSD	OD1-SG-CB	3.93	113.01	105.54
1	A	233	CSD	OD1-SG-CB	3.75	112.67	105.54

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	233	CSD	CA-CB-SG-OD1

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	B	233	CSD	CA-CB-SG-OD1
1	A	233	CSD	N-CA-CB-SG
1	B	233	CSD	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	97V	B	1201	-	43,45,45	2.60	10 (23%)	55,63,63	1.48	4 (7%)
2	97V	A	1201	-	43,45,45	2.57	10 (23%)	55,63,63	1.33	5 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	97V	B	1201	-	-	7/32/42/42	0/5/5/5
2	97V	A	1201	-	-	7/32/42/42	0/5/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	97V	OAC-SAB	7.85	1.66	1.44
2	A	1201	97V	OAD-SAB	7.78	1.65	1.44
2	B	1201	97V	OAC-SAB	7.70	1.65	1.44
2	B	1201	97V	OAD-SAB	7.47	1.65	1.44
2	B	1201	97V	CAE-SAB	6.23	1.83	1.77
2	A	1201	97V	CAE-SAB	5.71	1.83	1.77
2	A	1201	97V	CAK-NAJ	5.43	1.47	1.37
2	B	1201	97V	CAK-NAJ	5.41	1.46	1.37
2	B	1201	97V	CAV-NAU	5.27	1.45	1.38
2	A	1201	97V	CAV-NAU	4.88	1.45	1.38
2	B	1201	97V	CAK-NAM	4.51	1.48	1.39
2	A	1201	97V	CAK-NAM	4.44	1.48	1.39
2	B	1201	97V	CAS-NAU	3.98	1.46	1.35
2	A	1201	97V	CAS-NAU	3.85	1.46	1.35
2	A	1201	97V	CAI-NAJ	3.01	1.47	1.41
2	B	1201	97V	CAI-NAJ	2.91	1.47	1.41
2	B	1201	97V	CBF-CBG	2.23	1.56	1.52
2	A	1201	97V	CBF-CBG	2.15	1.56	1.52
2	B	1201	97V	OAL-CAK	-2.04	1.19	1.23
2	A	1201	97V	OAL-CAK	-2.02	1.19	1.23

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1201	97V	OAD-SAB-OAC	-5.93	107.71	117.92
2	B	1201	97V	CAA-SAB-CAE	5.74	111.36	104.58
2	A	1201	97V	OAD-SAB-OAC	-5.04	109.25	117.92
2	A	1201	97V	CAA-SAB-CAE	4.22	109.57	104.58
2	B	1201	97V	OAC-SAB-CAE	2.82	110.55	108.25
2	A	1201	97V	OAD-SAB-CAE	2.69	110.44	108.25
2	B	1201	97V	CAI-CBO-CAE	2.43	120.72	118.89
2	A	1201	97V	CAI-CBO-CAE	2.38	120.68	118.89
2	A	1201	97V	CBO-CAE-SAB	2.03	120.68	119.06

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1201	97V	CBO-CAE-SAB-CAA
2	B	1201	97V	CAF-CAE-SAB-CAA
2	B	1201	97V	CBO-CAE-SAB-OAC
2	A	1201	97V	CAF-CAE-SAB-OAD

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	B	1201	97V	CAF-CAE-SAB-OAC
2	A	1201	97V	CBO-CAE-SAB-OAD
2	A	1201	97V	CAO-CAN-NAM-CAK
2	B	1201	97V	CAO-CAN-NAM-CAK
2	A	1201	97V	CAF-CAE-SAB-CAA
2	A	1201	97V	CBE-CBF-CBG-CBL
2	B	1201	97V	CBE-CBF-CBG-CBL
2	A	1201	97V	CBO-CAE-SAB-CAA
2	A	1201	97V	CBM-CBF-CBG-CBL
2	B	1201	97V	CBM-CBF-CBG-CBL

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	422/455 (92%)	-0.09	10 (2%) 59 30	69, 103, 173, 208	0
1	B	422/455 (92%)	-0.09	13 (3%) 49 21	68, 105, 177, 230	0
All	All	844/910 (92%)	-0.09	23 (2%) 54 26	68, 104, 177, 230	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1037	PRO	6.1
1	A	1037	PRO	5.9
1	A	1040	ASN	4.3
1	A	422	LEU	3.9
1	B	1034	THR	3.2
1	B	1040	ASN	3.1
1	A	302	ASN	3.1
1	B	419	ARG	2.8
1	A	1015	LEU	2.7
1	B	418	GLU	2.5
1	A	1025	TYR	2.4
1	B	218	LEU	2.4
1	A	1057	VAL	2.4
1	B	1015	LEU	2.4
1	A	218	LEU	2.3
1	B	302	ASN	2.2
1	B	301	SER	2.2
1	A	1144	ASN	2.2
1	B	1025	TYR	2.2
1	B	219	SER	2.2
1	A	382	ILE	2.1
1	B	417	TRP	2.0
1	B	1036	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSD	B	233	8/9	0.90	0.32	96,97,100,102	0
1	CSD	A	233	8/9	0.92	0.26	91,93,98,99	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	97V	A	1201	41/41	0.88	0.33	124,143,148,149	0
2	97V	B	1201	41/41	0.91	0.34	124,146,152,154	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 97V A 1201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.