



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2024 – 06:42 PM EDT

PDB ID : 1VHW
Title : Crystal structure of purine nucleoside phosphorylase with adenosine
Authors : Structural GenomiX
Deposited on : 2003-12-01
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	
1	B	253	
1	C	253	
1	D	253	
1	E	253	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	253	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into three segments: a green segment representing 87%, a yellow segment representing 6%, and a grey segment representing 8%. A small red square is at the beginning of the bar, and a '%' symbol is above it.</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called purine nucleoside phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	1776	1124	297	337	18	0	4	0
1	B	237	1768	1119	297	335	17	0	3	0
1	C	237	1794	1131	303	342	18	0	6	0
1	D	237	1799	1137	304	340	18	0	6	0
1	E	236	1783	1129	299	337	18	0	5	0
1	F	234	1755	1112	298	328	17	0	2	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	cloning artifact	UNP Q9KPM0
A	0	SER	-	cloning artifact	UNP Q9KPM0
A	1	LEU	-	cloning artifact	UNP Q9KPM0
A	57	ARG	LYS	variant	UNP Q9KPM0
A	242	GLU	-	cloning artifact	UNP Q9KPM0
A	243	GLY	-	cloning artifact	UNP Q9KPM0
A	244	GLY	-	cloning artifact	UNP Q9KPM0
A	245	SER	-	cloning artifact	UNP Q9KPM0
A	246	HIS	-	cloning artifact	UNP Q9KPM0
A	247	HIS	-	cloning artifact	UNP Q9KPM0
A	248	HIS	-	cloning artifact	UNP Q9KPM0
A	249	HIS	-	cloning artifact	UNP Q9KPM0
A	250	HIS	-	cloning artifact	UNP Q9KPM0
A	251	HIS	-	cloning artifact	UNP Q9KPM0
B	-1	MET	-	cloning artifact	UNP Q9KPM0
B	0	SER	-	cloning artifact	UNP Q9KPM0
B	1	LEU	-	cloning artifact	UNP Q9KPM0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	57	ARG	LYS	variant	UNP Q9KPM0
B	242	GLU	-	cloning artifact	UNP Q9KPM0
B	243	GLY	-	cloning artifact	UNP Q9KPM0
B	244	GLY	-	cloning artifact	UNP Q9KPM0
B	245	SER	-	cloning artifact	UNP Q9KPM0
B	246	HIS	-	cloning artifact	UNP Q9KPM0
B	247	HIS	-	cloning artifact	UNP Q9KPM0
B	248	HIS	-	cloning artifact	UNP Q9KPM0
B	249	HIS	-	cloning artifact	UNP Q9KPM0
B	250	HIS	-	cloning artifact	UNP Q9KPM0
B	251	HIS	-	cloning artifact	UNP Q9KPM0
C	-1	MET	-	cloning artifact	UNP Q9KPM0
C	0	SER	-	cloning artifact	UNP Q9KPM0
C	1	LEU	-	cloning artifact	UNP Q9KPM0
C	57	ARG	LYS	variant	UNP Q9KPM0
C	242	GLU	-	cloning artifact	UNP Q9KPM0
C	243	GLY	-	cloning artifact	UNP Q9KPM0
C	244	GLY	-	cloning artifact	UNP Q9KPM0
C	245	SER	-	cloning artifact	UNP Q9KPM0
C	246	HIS	-	cloning artifact	UNP Q9KPM0
C	247	HIS	-	cloning artifact	UNP Q9KPM0
C	248	HIS	-	cloning artifact	UNP Q9KPM0
C	249	HIS	-	cloning artifact	UNP Q9KPM0
C	250	HIS	-	cloning artifact	UNP Q9KPM0
C	251	HIS	-	cloning artifact	UNP Q9KPM0
D	-1	MET	-	cloning artifact	UNP Q9KPM0
D	0	SER	-	cloning artifact	UNP Q9KPM0
D	1	LEU	-	cloning artifact	UNP Q9KPM0
D	57	ARG	LYS	variant	UNP Q9KPM0
D	242	GLU	-	cloning artifact	UNP Q9KPM0
D	243	GLY	-	cloning artifact	UNP Q9KPM0
D	244	GLY	-	cloning artifact	UNP Q9KPM0
D	245	SER	-	cloning artifact	UNP Q9KPM0
D	246	HIS	-	cloning artifact	UNP Q9KPM0
D	247	HIS	-	cloning artifact	UNP Q9KPM0
D	248	HIS	-	cloning artifact	UNP Q9KPM0
D	249	HIS	-	cloning artifact	UNP Q9KPM0
D	250	HIS	-	cloning artifact	UNP Q9KPM0
D	251	HIS	-	cloning artifact	UNP Q9KPM0
E	-1	MET	-	cloning artifact	UNP Q9KPM0
E	0	SER	-	cloning artifact	UNP Q9KPM0
E	1	LEU	-	cloning artifact	UNP Q9KPM0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	57	ARG	LYS	variant	UNP Q9KPM0
E	242	GLU	-	cloning artifact	UNP Q9KPM0
E	243	GLY	-	cloning artifact	UNP Q9KPM0
E	244	GLY	-	cloning artifact	UNP Q9KPM0
E	245	SER	-	cloning artifact	UNP Q9KPM0
E	246	HIS	-	cloning artifact	UNP Q9KPM0
E	247	HIS	-	cloning artifact	UNP Q9KPM0
E	248	HIS	-	cloning artifact	UNP Q9KPM0
E	249	HIS	-	cloning artifact	UNP Q9KPM0
E	250	HIS	-	cloning artifact	UNP Q9KPM0
E	251	HIS	-	cloning artifact	UNP Q9KPM0
F	-1	MET	-	cloning artifact	UNP Q9KPM0
F	0	SER	-	cloning artifact	UNP Q9KPM0
F	1	LEU	-	cloning artifact	UNP Q9KPM0
F	57	ARG	LYS	variant	UNP Q9KPM0
F	242	GLU	-	cloning artifact	UNP Q9KPM0
F	243	GLY	-	cloning artifact	UNP Q9KPM0
F	244	GLY	-	cloning artifact	UNP Q9KPM0
F	245	SER	-	cloning artifact	UNP Q9KPM0
F	246	HIS	-	cloning artifact	UNP Q9KPM0
F	247	HIS	-	cloning artifact	UNP Q9KPM0
F	248	HIS	-	cloning artifact	UNP Q9KPM0
F	249	HIS	-	cloning artifact	UNP Q9KPM0
F	250	HIS	-	cloning artifact	UNP Q9KPM0
F	251	HIS	-	cloning artifact	UNP Q9KPM0

- Molecule 2 is ADENOSINE (three-letter code: ADN) (formula: C₁₀H₁₃N₅O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	10	5	4		
2	B	1	Total	C	N	O	0	0
			19	10	5	4		
2	C	1	Total	C	N	O	0	0
			19	10	5	4		
2	D	1	Total	C	N	O	0	0
			19	10	5	4		
2	E	1	Total	C	N	O	0	0
			19	10	5	4		
2	F	1	Total	C	N	O	0	0
			19	10	5	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	301	Total	O	0	0
			301	301		
3	B	220	Total	O	0	0
			220	220		
3	C	292	Total	O	0	0
			292	292		
3	D	290	Total	O	0	0
			290	290		
3	E	294	Total	O	0	0
			294	294		
3	F	220	Total	O	0	0
			220	220		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: purine nucleoside phosphorylase




- Molecule 1: purine nucleoside phosphorylase




- Molecule 1: purine nucleoside phosphorylase




- Molecule 1: purine nucleoside phosphorylase




- Molecule 1: purine nucleoside phosphorylase



- Molecule 1: purine nucleoside phosphorylase

Chain F:  %
87% 6% 8%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.60Å 152.72Å 96.72Å 90.00° 103.91° 90.00°	Depositor
Resolution (Å)	33.52 – 1.54 33.56 – 1.54	Depositor EDS
% Data completeness (in resolution range)	(Not available) (33.52-1.54) 94.0 (33.56-1.54)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 1.54Å)	Xtrriage
Refinement program	REFMAC 4.0	Depositor
R, R_{free}	0.162 , 0.193 0.155 , 0.182	Depositor DCC
R_{free} test set	9091 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	11.4	Xtrriage
Anisotropy	0.479	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.477 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	12406	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ADN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/1822	1.00	8/2461 (0.3%)
1	B	0.47	0/1808	0.89	1/2442 (0.0%)
1	C	0.49	0/1848	0.96	3/2494 (0.1%)
1	D	0.49	0/1851	0.97	2/2497 (0.1%)
1	E	0.50	0/1831	0.99	5/2472 (0.2%)
1	F	0.47	0/1790	0.88	1/2413 (0.0%)
All	All	0.49	0/10950	0.95	20/14779 (0.1%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	120	ARG	NE-CZ-NH2	7.07	123.83	120.30
1	A	120	ARG	NE-CZ-NH2	6.42	123.51	120.30
1	C	161	TYR	CB-CG-CD2	-6.37	117.18	121.00
1	C	120	ARG	NE-CZ-NH2	6.18	123.39	120.30
1	E	120	ARG	NE-CZ-NH1	-6.13	117.24	120.30
1	A	193	TYR	CB-CG-CD2	6.11	124.67	121.00
1	A	187	TYR	CB-CG-CD1	-5.98	117.41	121.00
1	A	193	TYR	CB-CG-CD1	-5.82	117.51	121.00
1	A	187	TYR	CB-CG-CD2	5.70	124.42	121.00
1	E	118	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	F	120	ARG	NE-CZ-NH2	5.48	123.04	120.30
1	E	47	PHE	CB-CG-CD1	-5.44	116.99	120.80
1	E	187	TYR	CB-CG-CD1	-5.37	117.78	121.00
1	A	44	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	D	57	ARG	NE-CZ-NH1	-5.31	117.65	120.30
1	A	145	ARG	NE-CZ-NH2	5.27	122.94	120.30
1	D	120	ARG	NE-CZ-NH2	5.21	122.91	120.30
1	C	161	TYR	CB-CG-CD1	5.20	124.12	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	161	TYR	CB-CG-CD2	-5.10	117.94	121.00
1	A	47	PHE	CB-CG-CD1	-5.07	117.25	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1776	0	1735	14	0
1	B	1768	0	1729	17	0
1	C	1794	0	1761	19	0
1	D	1799	0	1781	13	0
1	E	1783	0	1763	13	0
1	F	1755	0	1734	14	0
2	A	19	0	13	0	0
2	B	19	0	13	0	0
2	C	19	0	13	0	0
2	D	19	0	13	0	0
2	E	19	0	13	0	0
2	F	19	0	13	0	0
3	A	301	0	0	4	0
3	B	220	0	0	3	0
3	C	292	0	0	3	0
3	D	290	0	0	2	0
3	E	294	0	0	3	0
3	F	220	0	0	1	0
All	All	12406	0	10581	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108[A]:MET:SD	1:E:108[A]:MET:SD	2.67	0.93
1:B:108[A]:MET:SD	1:D:108[A]:MET:SD	2.67	0.92
1:C:108[A]:MET:SD	1:F:108[A]:MET:SD	2.69	0.90
1:C:100:LYS:HZ2	1:C:101:VAL:H	1.33	0.77
1:B:96:ASN:HB2	3:B:468:HOH:O	1.92	0.69
1:D:96:ASN:HB2	3:D:537:HOH:O	1.96	0.65
1:C:96:ASN:HB2	3:C:364:HOH:O	1.96	0.65
1:B:108[A]:MET:CG	1:D:108[A]:MET:SD	2.87	0.62
1:C:108[A]:MET:SD	1:F:108[A]:MET:CG	2.89	0.61
1:B:108[A]:MET:SD	1:D:108[A]:MET:CG	2.90	0.60
1:A:108[A]:MET:SD	1:E:108[A]:MET:CG	2.91	0.59
1:B:104:VAL:CG2	1:B:147:ILE:HD11	2.33	0.58
1:C:108[A]:MET:CG	1:F:108[A]:MET:SD	2.92	0.58
1:A:108[A]:MET:CG	1:E:108[A]:MET:SD	2.92	0.56
1:D:204:SER:HB3	3:D:384:HOH:O	2.03	0.56
1:C:108[A]:MET:SD	1:F:108[A]:MET:HG3	2.46	0.56
1:B:108[A]:MET:HG3	1:D:108[A]:MET:SD	2.46	0.55
1:B:172:ASP:HA	3:B:468:HOH:O	2.05	0.55
1:A:2:ALA:HB3	3:A:530:HOH:O	2.06	0.55
1:C:108[A]:MET:HG3	1:F:108[A]:MET:SD	2.47	0.55
1:B:108[A]:MET:SD	1:D:108[A]:MET:HG3	2.48	0.54
1:B:101:VAL:HG11	1:B:213:THR:HG23	1.89	0.53
1:B:147:ILE:HG21	1:B:228:ILE:HD11	1.91	0.53
1:C:102:ARG:HD2	1:C:213:THR:HG21	1.92	0.52
1:B:101:VAL:CG1	1:B:213:THR:HG23	2.40	0.52
1:F:101:VAL:O	1:F:102:ARG:HB2	2.10	0.51
1:E:3:THR:HB	1:E:4:PRO:HD2	1.92	0.51
1:E:149:VAL:HG22	3:E:435:HOH:O	2.11	0.49
1:E:39[A]:GLN:NE2	1:E:42:ASP:HB3	2.28	0.49
1:A:145:ARG:NE	1:A:227:GLU:HG2	2.28	0.49
1:C:216:GLU:OE2	1:C:217:GLU:HG3	2.13	0.48
1:A:123:ASP:HB2	3:A:543:HOH:O	2.14	0.48
1:F:96:ASN:HB3	1:F:99:ILE:CD1	2.44	0.48
1:F:147:ILE:HD13	1:F:228:ILE:HD11	1.96	0.47
1:C:206:HIS:HB3	1:C:209:THR:HG22	1.95	0.47
1:F:145[B]:ARG:HH21	1:F:145[B]:ARG:HG3	1.79	0.47
1:A:108[A]:MET:SD	1:E:108[A]:MET:HG3	2.55	0.47
1:A:108[A]:MET:HG3	1:E:108[A]:MET:SD	2.55	0.47
1:B:104:VAL:HG21	1:B:147:ILE:HD11	1.98	0.47
1:C:100:LYS:NZ	1:C:101:VAL:H	2.07	0.46
1:C:173:LYS:HD3	1:C:174:TYR:CE2	2.51	0.46
1:C:108[A]:MET:CE	1:F:108[A]:MET:HG3	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:136:LYS:O	1:B:140:GLU:HG3	2.16	0.45
1:C:84:LYS:HD3	3:C:348:HOH:O	2.15	0.45
1:C:3:THR:HB	1:C:4:PRO:HD2	1.99	0.45
1:F:96:ASN:HB3	1:F:99:ILE:HD13	1.99	0.45
1:D:63:HIS:CE1	1:D:182:GLU:HG2	2.53	0.44
1:C:63:HIS:CE1	1:C:182:GLU:HG2	2.52	0.44
1:B:102:ARG:HD2	1:B:221:THR:OG1	2.18	0.44
1:D:136:LYS:O	1:D:140:GLU:HG3	2.18	0.44
3:A:551:HOH:O	1:E:133:LYS:HE3	2.18	0.43
1:F:158:GLU:HG3	3:F:472:HOH:O	2.17	0.43
1:F:63:HIS:CE1	1:F:182:GLU:HG2	2.54	0.43
1:A:145:ARG:HD2	3:A:398:HOH:O	2.19	0.43
1:C:173:LYS:HE2	3:C:531:HOH:O	2.18	0.42
1:E:63:HIS:CE1	1:E:182:GLU:HG2	2.55	0.42
1:B:204:SER:O	1:B:213:THR:OG1	2.37	0.42
1:C:100:LYS:HG3	1:C:101:VAL:N	2.34	0.42
1:E:211:GLU:HG3	3:E:397:HOH:O	2.20	0.41
1:F:145[B]:ARG:HG3	1:F:145[B]:ARG:NH2	2.34	0.41
1:A:3:THR:HB	1:A:4:PRO:HD2	2.02	0.41
1:B:101:VAL:O	1:B:102:ARG:HB2	2.20	0.41
1:E:196:LYS:HE2	3:E:489:HOH:O	2.20	0.41
1:A:45:ASN:HB3	1:D:45:ASN:HB3	2.01	0.41
1:A:44:ARG:HA	1:D:22:ASP:OD1	2.20	0.41
1:D:22:ASP:HB3	1:D:25:ARG:HG3	2.03	0.41
1:A:145:ARG:HE	1:A:227:GLU:HG2	1.86	0.41
1:C:22:ASP:OD1	1:E:44:ARG:HA	2.21	0.41
1:B:213:THR:HG22	3:B:376:HOH:O	2.21	0.40
1:A:22:ASP:OD1	1:D:44:ARG:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/253 (94%)	233 (98%)	6 (2%)	0	100	100
1	B	238/253 (94%)	231 (97%)	7 (3%)	0	100	100
1	C	241/253 (95%)	235 (98%)	6 (2%)	0	100	100
1	D	241/253 (95%)	234 (97%)	7 (3%)	0	100	100
1	E	239/253 (94%)	233 (98%)	6 (2%)	0	100	100
1	F	232/253 (92%)	225 (97%)	7 (3%)	0	100	100
All	All	1430/1518 (94%)	1391 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/201 (90%)	181 (100%)	1 (0%)	88	77
1	B	180/201 (90%)	176 (98%)	4 (2%)	52	21
1	C	188/201 (94%)	185 (98%)	3 (2%)	62	33
1	D	189/201 (94%)	186 (98%)	3 (2%)	62	33
1	E	187/201 (93%)	186 (100%)	1 (0%)	88	77
1	F	179/201 (89%)	176 (98%)	3 (2%)	60	31
All	All	1105/1206 (92%)	1090 (99%)	15 (1%)	67	39

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	GLU
1	B	56	ARG
1	B	180	GLU
1	B	213	THR
1	B	219	GLN
1	C	100	LYS
1	C	180	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	216	GLU
1	D	100	LYS
1	D	180	GLU
1	D	218	ARG
1	E	180	GLU
1	F	3	THR
1	F	56	ARG
1	F	180	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	223	ASN
1	B	39	GLN
1	B	219	GLN
1	B	223	ASN
1	C	220	ASN
1	C	223	ASN
1	C	238	GLN
1	D	219	GLN
1	D	220	ASN
1	D	223	ASN
1	D	238	GLN
1	E	220	ASN
1	E	223	ASN
1	E	238	GLN
1	F	39	GLN
1	F	220	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADN	D	252	-	17,21,21	1.57	3 (17%)	17,31,31	1.82	4 (23%)
2	ADN	B	252	-	17,21,21	1.42	4 (23%)	17,31,31	1.62	3 (17%)
2	ADN	E	252	-	17,21,21	1.55	4 (23%)	17,31,31	1.60	4 (23%)
2	ADN	C	252	-	17,21,21	1.54	4 (23%)	17,31,31	1.80	4 (23%)
2	ADN	F	252	-	17,21,21	1.48	3 (17%)	17,31,31	1.77	4 (23%)
2	ADN	A	252	-	17,21,21	1.55	4 (23%)	17,31,31	1.66	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADN	D	252	-	-	0/2/22/22	0/3/3/3
2	ADN	B	252	-	-	0/2/22/22	0/3/3/3
2	ADN	E	252	-	-	0/2/22/22	0/3/3/3
2	ADN	C	252	-	-	0/2/22/22	0/3/3/3
2	ADN	F	252	-	-	0/2/22/22	0/3/3/3
2	ADN	A	252	-	-	0/2/22/22	0/3/3/3

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	252	ADN	C8-N7	-3.62	1.28	1.34
2	B	252	ADN	C8-N7	-3.53	1.28	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	252	ADN	C8-N7	-3.49	1.28	1.34
2	E	252	ADN	C8-N7	-3.43	1.28	1.34
2	D	252	ADN	C2-N1	3.24	1.39	1.33
2	A	252	ADN	C8-N7	-3.17	1.28	1.34
2	C	252	ADN	C2-N1	3.07	1.39	1.33
2	C	252	ADN	C8-N7	-3.01	1.29	1.34
2	A	252	ADN	C2-N1	2.94	1.39	1.33
2	E	252	ADN	C2-N1	2.92	1.39	1.33
2	F	252	ADN	C2-N1	2.52	1.38	1.33
2	A	252	ADN	C1'-N9	-2.38	1.44	1.49
2	C	252	ADN	O4'-C1'	-2.33	1.37	1.40
2	D	252	ADN	O4'-C1'	-2.32	1.37	1.40
2	F	252	ADN	C1'-N9	-2.31	1.44	1.49
2	E	252	ADN	C1'-N9	-2.30	1.44	1.49
2	A	252	ADN	C5-N7	-2.28	1.31	1.39
2	B	252	ADN	C1'-N9	-2.20	1.44	1.49
2	C	252	ADN	C1'-N9	-2.16	1.44	1.49
2	E	252	ADN	C5-N7	-2.15	1.32	1.39
2	B	252	ADN	C5-N7	-2.14	1.32	1.39
2	B	252	ADN	C2-N1	2.09	1.37	1.33

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	252	ADN	C1'-N9-C4	4.31	134.22	126.64
2	C	252	ADN	C1'-N9-C4	4.21	134.03	126.64
2	F	252	ADN	C5-C6-N6	3.98	126.38	120.31
2	D	252	ADN	C5-C6-N6	3.84	126.17	120.31
2	F	252	ADN	C1'-N9-C4	3.79	133.30	126.64
2	C	252	ADN	C5-C6-N6	3.73	125.99	120.31
2	B	252	ADN	C5-C6-N6	3.63	125.84	120.31
2	A	252	ADN	C4-C5-N7	3.50	113.04	109.34
2	B	252	ADN	C1'-N9-C4	3.40	132.61	126.64
2	A	252	ADN	C1'-N9-C4	3.29	132.41	126.64
2	E	252	ADN	C1'-N9-C4	3.06	132.02	126.64
2	E	252	ADN	C4-C5-N7	3.01	112.52	109.34
2	E	252	ADN	O4'-C1'-N9	3.00	112.72	108.75
2	F	252	ADN	C4-C5-N7	2.95	112.46	109.34
2	B	252	ADN	C4-C5-N7	2.88	112.38	109.34
2	D	252	ADN	O4'-C1'-N9	2.84	112.52	108.75
2	A	252	ADN	C5-C6-N6	2.80	124.57	120.31
2	A	252	ADN	O4'-C1'-N9	2.67	112.29	108.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	252	ADN	C5-C6-N6	2.55	124.19	120.31
2	D	252	ADN	C4-C5-N7	2.43	111.90	109.34
2	C	252	ADN	C4-C5-N7	2.39	111.86	109.34
2	F	252	ADN	O4'-C1'-N9	2.32	111.82	108.75
2	C	252	ADN	O4'-C1'-N9	2.29	111.79	108.75


There are no chirality outliers.



There are no torsion outliers.



There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/253 (93%)	-0.64	2 (0%) 86 88	6, 10, 26, 38	0
1	B	237/253 (93%)	-0.36	4 (1%) 70 74	7, 15, 36, 46	0
1	C	237/253 (93%)	-0.59	6 (2%) 57 63	6, 11, 29, 44	0
1	D	237/253 (93%)	-0.56	7 (2%) 50 57	6, 11, 28, 43	0
1	E	236/253 (93%)	-0.65	2 (0%) 86 88	5, 10, 25, 38	0
1	F	234/253 (92%)	-0.39	3 (1%) 77 81	6, 15, 33, 43	0
All	All	1418/1518 (93%)	-0.53	24 (1%) 70 74	5, 12, 30, 46	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	209	THR	5.3
1	B	213	THR	4.6
1	C	209	THR	4.3
1	B	2	ALA	4.2
1	D	207	ILE	4.0
1	D	2	ALA	3.9
1	D	212	GLN	3.9
1	C	207	ILE	3.9
1	F	209	THR	3.3
1	A	212	GLN	3.2
1	D	211	GLU	3.0
1	D	210	GLY	3.0
1	C	212	GLN	2.9
1	E	212	GLN	2.9
1	C	98	GLY	2.9
1	B	209	THR	2.7
1	A	213	THR	2.7
1	B	208	LYS	2.5
1	C	2	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	2	ALA	2.4
1	E	211	GLU	2.4
1	D	98	GLY	2.2
1	F	213	THR	2.2
1	C	210	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ADN	F	252	19/19	0.96	0.06	11,13,15,15	0
2	ADN	D	252	19/19	0.97	0.06	7,9,12,14	0
2	ADN	C	252	19/19	0.97	0.06	7,10,11,14	0
2	ADN	B	252	19/19	0.98	0.05	11,13,14,14	0
2	ADN	E	252	19/19	0.98	0.05	8,9,11,13	0
2	ADN	A	252	19/19	0.98	0.05	8,9,12,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ADN F 252:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADN D 252:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADN C 252:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADN B 252:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ADN E 252:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.