



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 31, 2024 – 06:28 PM EDT

PDB ID : 8VIH
Title : EgtB-IV from *Crocospaera subtropica*, an ergothioneine-biosynthetic type IV sulfoxide synthase in complex with hercynine
Authors : Ireland, K.A.; Davis, K.M.
Deposited on : 2024-01-04
Resolution : 1.55 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	468	 92% 5%
1	B	468	 90% 7%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 8006 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfoxide synthase EgtB-IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	444	3621	2357	589	660	15	0	4	0
1	B	435	3477	2267	568	628	14	0	3	1

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP B1WTS6
A	-18	GLY	-	expression tag	UNP B1WTS6
A	-17	SER	-	expression tag	UNP B1WTS6
A	-16	SER	-	expression tag	UNP B1WTS6
A	-15	HIS	-	expression tag	UNP B1WTS6
A	-14	HIS	-	expression tag	UNP B1WTS6
A	-13	HIS	-	expression tag	UNP B1WTS6
A	-12	HIS	-	expression tag	UNP B1WTS6
A	-11	HIS	-	expression tag	UNP B1WTS6
A	-10	HIS	-	expression tag	UNP B1WTS6
A	-9	SER	-	expression tag	UNP B1WTS6
A	-8	SER	-	expression tag	UNP B1WTS6
A	-7	GLY	-	expression tag	UNP B1WTS6
A	-6	LEU	-	expression tag	UNP B1WTS6
A	-5	VAL	-	expression tag	UNP B1WTS6
A	-4	PRO	-	expression tag	UNP B1WTS6
A	-3	ARG	-	expression tag	UNP B1WTS6
A	-2	GLY	-	expression tag	UNP B1WTS6
A	-1	SER	-	expression tag	UNP B1WTS6
A	0	HIS	-	expression tag	UNP B1WTS6
B	-19	MET	-	expression tag	UNP B1WTS6
B	-18	GLY	-	expression tag	UNP B1WTS6
B	-17	SER	-	expression tag	UNP B1WTS6
B	-16	SER	-	expression tag	UNP B1WTS6
B	-15	HIS	-	expression tag	UNP B1WTS6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP B1WTS6
B	-13	HIS	-	expression tag	UNP B1WTS6
B	-12	HIS	-	expression tag	UNP B1WTS6
B	-11	HIS	-	expression tag	UNP B1WTS6
B	-10	HIS	-	expression tag	UNP B1WTS6
B	-9	SER	-	expression tag	UNP B1WTS6
B	-8	SER	-	expression tag	UNP B1WTS6
B	-7	GLY	-	expression tag	UNP B1WTS6
B	-6	LEU	-	expression tag	UNP B1WTS6
B	-5	VAL	-	expression tag	UNP B1WTS6
B	-4	PRO	-	expression tag	UNP B1WTS6
B	-3	ARG	-	expression tag	UNP B1WTS6
B	-2	GLY	-	expression tag	UNP B1WTS6
B	-1	SER	-	expression tag	UNP B1WTS6
B	0	HIS	-	expression tag	UNP B1WTS6

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0


- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0
3	B	1	Total Na 1 1	0	0

- Molecule 4 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Fe 1 1	0	0
4	B	1	Total Fe 1 1	0	0

- Molecule 5 is N,N,N-trimethyl-histidine (three-letter code: AVJ) (formula: C₉H₁₆N₃O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
5	A	1	Total	C	N	O	0	0
			14	9	3	2		
5	B	1	Total	C	N	O	0	0
			14	9	3	2		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	470	Total O 470 470	0	0
7	B	384	Total O 384 384	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sulfoxide synthase EgtB-IV



- Molecule 1: Sulfoxide synthase EgtB-IV



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.20Å 94.51Å 83.75Å 90.00° 100.16° 90.00°	Depositor
Resolution (Å)	47.26 – 1.55 47.26 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.8 (47.26-1.55) 98.8 (47.26-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 1.55Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.180 , 0.198 0.177 , 0.197	Depositor DCC
R_{free} test set	1999 reflections (1.46%)	wwPDB-VP
Wilson B-factor (Å ²)	19.0	Xtrriage
Anisotropy	0.094	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8006	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, AVJ, FE, EDO, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.82	0/3750	0.78	0/5106
1	B	0.78	0/3600	0.75	0/4903
All	All	0.80	0/7350	0.77	0/10009

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3621	0	3348	7	0
1	B	3477	0	3150	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	14	0	15	1	0
5	B	14	0	15	2	0
6	A	12	0	18	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	8	0	12	1	0
7	A	470	0	0	0	0
7	B	384	0	0	0	0
All	All	8006	0	6558	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:VAL:HG13	1:A:396:LEU:HD21	1.82	0.59
1:B:141:ALA:HA	1:B:146:TRP:CD2	2.46	0.50
1:A:141:ALA:HA	1:A:146:TRP:CD2	2.50	0.47
1:B:157:ILE:HG21	5:B:504:AVJ:HAPA	1.98	0.46
1:B:73:VAL:HG13	1:B:396:LEU:HD21	1.99	0.45
1:A:301:VAL:HB	1:A:305:GLU:HB2	2.00	0.44
1:A:80:ILE:CD1	1:A:122:ALA:HA	2.48	0.43
1:A:188:VAL:HG22	1:A:189:PRO:HD2	2.01	0.42
1:B:80:ILE:HD11	1:B:122:ALA:HA	2.01	0.42
1:B:357:PRO:HG2	1:B:360:LEU:HD13	2.01	0.42
5:B:504:AVJ:HAB	6:B:506:EDO:H11	2.01	0.42
1:A:157:ILE:HG21	5:A:504:AVJ:HAPA	2.02	0.42
1:B:320:MET:HG3	1:B:445:VAL:HG11	2.01	0.41
1:A:11:PHE:HB3	1:A:13:ASN:OD1	2.21	0.41
1:B:139:ILE:HD12	1:B:145:LEU:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/468 (94%)	435 (98%)	7 (2%)	0	100	100
1	B	429/468 (92%)	422 (98%)	7 (2%)	0	100	100
All	All	871/936 (93%)	857 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/417 (89%)	373 (100%)	0	100	100
1	B	344/417 (82%)	343 (100%)	1 (0%)	92	85
All	All	717/834 (86%)	716 (100%)	1 (0%)	93	86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	397	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	132	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	B	505	-	3,3,3	0.36	0	2,2,2	0.49	0
6	EDO	A	506	-	3,3,3	0.71	0	2,2,2	0.69	0
6	EDO	A	507	-	3,3,3	0.70	0	2,2,2	0.20	0
5	AVJ	A	504	4	9,14,14	0.86	1 (11%)	11,20,20	0.86	1 (9%)
5	AVJ	B	504	4	9,14,14	0.80	0	11,20,20	0.86	1 (9%)
6	EDO	A	505	-	3,3,3	0.62	0	2,2,2	0.25	0
6	EDO	B	506	-	3,3,3	0.59	0	2,2,2	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	B	505	-	-	0/1/1/1	-
6	EDO	A	506	-	-	0/1/1/1	-
6	EDO	A	507	-	-	1/1/1/1	-
5	AVJ	A	504	4	-	3/14/14/14	0/1/1/1
5	AVJ	B	504	4	-	1/14/14/14	0/1/1/1
6	EDO	A	505	-	-	1/1/1/1	-
6	EDO	B	506	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	504	AVJ	OAL-CAX	-2.01	1.24	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	504	AVJ	OAL-CAX-CBC	2.10	116.85	111.87
5	B	504	AVJ	OAL-CAX-CBC	2.08	116.81	111.87

There are no chirality outliers.

All (6) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	A	505	EDO	O1-C1-C2-O2
5	A	504	AVJ	OAL-CAX-CBC-NBE
5	B	504	AVJ	OAL-CAX-CBC-NBE
5	A	504	AVJ	OAH-CAX-CBC-CAP
5	A	504	AVJ	OAL-CAX-CBC-CAP
6	A	507	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	504	AVJ	1	0
5	B	504	AVJ	2	0
6	B	506	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/468 (94%)	0.12	7 (1%) 72 77	10, 17, 33, 52	0
1	B	435/468 (92%)	0.18	18 (4%) 37 43	11, 20, 40, 61	0
All	All	879/936 (93%)	0.14	25 (2%) 53 60	10, 19, 37, 61	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	137	LEU	5.4
1	B	78	LYS	3.5
1	A	91	GLY	3.3
1	A	90	PHE	3.1
1	B	106	HIS	3.0
1	B	101	ASN	3.0
1	A	96	ASN	2.9
1	B	105	ALA	2.8
1	B	335	ASN	2.6
1	B	277	PHE	2.6
1	B	86	ILE	2.5
1	A	185	SER	2.5
1	B	279	ASN	2.4
1	B	448	HIS	2.4
1	B	14	ASN	2.3
1	B	108	ASN	2.3
1	A	92	VAL	2.3
1	B	133	THR	2.3
1	B	134	THR	2.3
1	B	139	ILE	2.2
1	A	186	GLN	2.2
1	A	102	GLN	2.1
1	B	143	HIS	2.1
1	B	2	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	136	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	AVJ	B	504	14/14	0.74	0.15	16,20,22,25	14
5	AVJ	A	504	14/14	0.83	0.15	14,18,20,21	14
6	EDO	A	505	4/4	0.86	0.15	24,34,35,35	0
6	EDO	A	507	4/4	0.86	0.11	24,26,30,36	0
6	EDO	B	506	4/4	0.91	0.10	19,19,20,28	0
6	EDO	B	505	4/4	0.92	0.11	24,24,28,30	0
6	EDO	A	506	4/4	0.95	0.07	16,18,21,23	0
2	CL	A	501	1/1	0.97	0.09	24,24,24,24	0
3	NA	B	502	1/1	0.98	0.10	19,19,19,19	0
2	CL	B	501	1/1	0.99	0.08	20,20,20,20	0
3	NA	A	502	1/1	0.99	0.07	23,23,23,23	0
4	FE	B	503	1/1	1.00	0.03	20,20,20,20	0
4	FE	A	503	1/1	1.00	0.04	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AVJ B 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.