



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2023 – 07:13 AM EDT

PDB ID : 6VLS
Title : Structure of C-terminal fragment of Vip3A toxin
Authors : Jiang, K.; Zhang, Y.; Chen, Z.; Gao, X.
Deposited on : 2020-01-25
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 30179 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose/maltodextrin-binding periplasmic protein,Vip3Aa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	953	7475	4773	1218	1467	17	0	0	0
1	B	950	7449	4755	1215	1462	17	0	0	0
1	C	954	7480	4773	1220	1470	17	0	0	0
1	D	963	7556	4820	1232	1487	17	0	0	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	initiating methionine	UNP P0AEX9
A	387	ALA	-	linker	UNP P0AEX9
A	388	ALA	-	linker	UNP P0AEX9
A	389	ARG	-	linker	UNP P0AEX9
A	390	ALA	-	linker	UNP P0AEX9
A	391	PHE	-	linker	UNP P0AEX9
A	392	ALA	-	linker	UNP P0AEX9
A	393	ALA	-	linker	UNP P0AEX9
A	394	ALA	-	linker	UNP P0AEX9
A	395	SER	-	linker	UNP P0AEX9
B	20	MET	-	initiating methionine	UNP P0AEX9
B	387	ALA	-	linker	UNP P0AEX9
B	388	ALA	-	linker	UNP P0AEX9
B	389	ARG	-	linker	UNP P0AEX9
B	390	ALA	-	linker	UNP P0AEX9
B	391	PHE	-	linker	UNP P0AEX9
B	392	ALA	-	linker	UNP P0AEX9
B	393	ALA	-	linker	UNP P0AEX9
B	394	ALA	-	linker	UNP P0AEX9
B	395	SER	-	linker	UNP P0AEX9
C	20	MET	-	initiating methionine	UNP P0AEX9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	387	ALA	-	linker	UNP P0AEX9
C	388	ALA	-	linker	UNP P0AEX9
C	389	ARG	-	linker	UNP P0AEX9
C	390	ALA	-	linker	UNP P0AEX9
C	391	PHE	-	linker	UNP P0AEX9
C	392	ALA	-	linker	UNP P0AEX9
C	393	ALA	-	linker	UNP P0AEX9
C	394	ALA	-	linker	UNP P0AEX9
C	395	SER	-	linker	UNP P0AEX9
D	20	MET	-	initiating methionine	UNP P0AEX9
D	387	ALA	-	linker	UNP P0AEX9
D	388	ALA	-	linker	UNP P0AEX9
D	389	ARG	-	linker	UNP P0AEX9
D	390	ALA	-	linker	UNP P0AEX9
D	391	PHE	-	linker	UNP P0AEX9
D	392	ALA	-	linker	UNP P0AEX9
D	393	ALA	-	linker	UNP P0AEX9
D	394	ALA	-	linker	UNP P0AEX9
D	395	SER	-	linker	UNP P0AEX9

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			7	4	3		
2	C	1	Total	C	O	0	0
			7	4	3		
2	D	1	Total	C	O	0	0
			7	4	3		
2	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	40	Total	O	0	0
			40	40		
3	B	49	Total	O	0	0
			49	49		
3	C	36	Total	O	0	0
			36	36		
3	D	52	Total	O	0	0
			52	52		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	136.09Å 127.88Å 149.18Å 90.00° 91.38° 90.00°	Depositor
Resolution (Å)	28.93 – 3.20	Depositor
% Data completeness (in resolution range)	97.2 (28.93-3.20)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 3.19Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.198 , 0.239	Depositor
Wilson B-factor (Å ²)	70.2	Xtrriage
Anisotropy	0.147	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.037 for h,-k,-l	Xtrriage
Total number of atoms	30179	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PEG	B	1001	-	6,6,6	0.49	0	5,5,5	0.28	0
2	PEG	D	1001	-	6,6,6	0.49	0	5,5,5	0.25	0
2	PEG	D	1002	-	6,6,6	0.49	0	5,5,5	0.25	0
2	PEG	A	1002	-	6,6,6	0.49	0	5,5,5	0.27	0
2	PEG	A	1001	-	6,6,6	0.49	0	5,5,5	0.27	0
2	PEG	C	1001	-	6,6,6	0.49	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	B	1001	-	-	2/4/4/4	-
2	PEG	D	1001	-	-	1/4/4/4	-
2	PEG	D	1002	-	-	2/4/4/4	-
2	PEG	A	1002	-	-	3/4/4/4	-
2	PEG	A	1001	-	-	2/4/4/4	-
2	PEG	C	1001	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1001	PEG	O1-C1-C2-O2
2	A	1002	PEG	O1-C1-C2-O2
2	A	1001	PEG	C4-C3-O2-C2
2	D	1002	PEG	C1-C2-O2-C3
2	B	1001	PEG	C1-C2-O2-C3
2	A	1002	PEG	C1-C2-O2-C3
2	B	1001	PEG	C4-C3-O2-C2
2	D	1002	PEG	O2-C3-C4-O4
2	A	1002	PEG	O2-C3-C4-O4
2	D	1001	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1001	PEG	C1-C2-O2-C3

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.