



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 16, 2024 – 12:05 PM EDT

PDB ID : 8VM8  
Title : The crystal structure of coxsackievirus B3 RNA replication element sD-loop mutant in complex with Fab BL3-6  
Authors : Das, N.K.; Koirala, D.  
Deposited on : 2024-01-13  
Resolution : 1.54 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)
RNA backbone	3102	1015 (2.36-0.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	93	
2	H	232	
3	L	215	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6189 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (93-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	R	93	1970	880	342	655	93	0	0	0

- Molecule 2 is a protein called Heavy Chain of Fab BL3-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	227	1692	1062	289	335	6	0	0	0

- Molecule 3 is a protein called Light Chain of Fab BL3-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	215	1643	1025	275	337	6	0	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	R	98	Total 98	O 98	0	0
4	H	365	Total 365	O 365	0	0
4	L	421	Total 421	O 421	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (93-MER)



- Molecule 2: Heavy Chain of Fab BL3-6



- Molecule 3: Light Chain of Fab BL3-6



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	145.99Å 80.37Å 87.34Å 90.00° 111.17° 90.00°	Depositor
Resolution (Å)	38.31 – 1.54 38.31 – 1.54	Depositor EDS
% Data completeness (in resolution range)	99.5 (38.31-1.54) 86.8 (38.31-1.54)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.52 (at 1.54Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.186 , 0.213 0.185 , 0.213	Depositor DCC
$R_{free}$ test set	1997 reflections (1.44%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.3	Xtrriage
Anisotropy	0.424	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6189	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	R	0.27	0/2198	0.80	1/3421 (0.0%)
2	H	0.39	0/1733	0.60	0/2358
3	L	0.41	0/1678	0.60	0/2277
All	All	0.35	0/5609	0.69	1/8056 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	R	65	A	O4'-C1'-N9	-5.43	103.86	108.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	1970	0	1001	14	0
2	H	1692	0	1646	1	0
3	L	1643	0	1594	0	0
4	H	365	0	0	0	0
4	L	421	0	0	0	0
4	R	98	0	0	1	0
All	All	6189	0	4241	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:7:A:H61	1:R:87:U:H3	1.39	0.71
1:R:21:G:H1	1:R:26:C:H42	1.41	0.66
1:R:49:U:H4'	1:R:50:A:H3'	1.94	0.48
1:R:32:C:H2'	1:R:33:A:C8	2.48	0.48
2:H:15:VAL:HG11	2:H:89:LEU:HD13	1.96	0.47
1:R:8:C:H2'	1:R:9:A:C8	2.50	0.46
1:R:23:U:H2'	1:R:24:C:O4'	2.15	0.46
1:R:30:C:H2'	1:R:31:A:C8	2.51	0.46
1:R:26:C:H2'	1:R:27:A:C8	2.52	0.45
1:R:75:U:H2'	1:R:76:U:C6	2.51	0.45
1:R:57:U:H2'	1:R:58:G:O4'	2.18	0.44
1:R:10:G:N3	1:R:47:G:H1'	2.33	0.43
1:R:72:A:OP2	4:R:101:HOH:O	2.21	0.42
1:R:11:C:H4'	1:R:47:G:H5'	2.02	0.41
1:R:10:G:C2	1:R:47:G:H1'	2.55	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	225/232 (97%)	221 (98%)	4 (2%)	0	100	100
3	L	213/215 (99%)	208 (98%)	5 (2%)	0	100	100
All	All	438/447 (98%)	429 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	H	188/193 (97%)	188 (100%)	0	100	100
3	L	190/190 (100%)	190 (100%)	0	100	100
All	All	378/383 (99%)	378 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	92/93 (98%)	17 (18%)	0

All (17) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	R	7	A
1	R	16	G
1	R	19	U
1	R	20	U
1	R	21	G
1	R	22	A
1	R	25	C
1	R	26	C
1	R	27	A
1	R	42	U
1	R	43	G
1	R	49	U
1	R	50	A
1	R	51	G
1	R	56	C
1	R	59	G

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	R	89	U

There are no RNA pucker outliers to report.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	R	93/93 (100%)	3.56	53 (56%) <b>0</b> <b>0</b>	20, 95, 187, 206	0
2	H	227/232 (97%)	0.46	19 (8%) <b>11</b> <b>12</b>	17, 30, 70, 108	0
3	L	215/215 (100%)	0.14	4 (1%) <b>66</b> <b>72</b>	18, 25, 42, 62	0
All	All	535/540 (99%)	0.87	76 (14%) <b>2</b> <b>2</b>	17, 30, 120, 206	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	R	21	G	19.9
1	R	25	C	12.9
1	R	16	G	12.1
1	R	22	A	10.7
1	R	33	A	10.1
1	R	18	G	10.1
1	R	17	G	9.8
1	R	34	G	9.8
1	R	1	G	9.3
1	R	14	G	9.2
1	R	32	C	9.2
1	R	35	G	9.0
2	H	145	GLY	8.9
1	R	31	A	8.9
1	R	27	A	8.8
1	R	24	C	8.7
1	R	20	U	8.7
1	R	26	C	7.9
1	R	23	U	7.3
2	H	144	SER	6.6
1	R	49	U	6.4
1	R	28	C	6.1
1	R	30	C	6.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	R	50	A	5.9
1	R	19	U	5.8
1	R	86	G	5.4
1	R	29	C	5.3
1	R	53	A	5.3
2	H	3	SER	5.2
2	H	143	THR	5.2
2	H	203	THR	5.1
1	R	15	U	4.9
2	H	140	SER	4.8
1	R	36	G	4.7
2	H	142	SER	4.4
1	R	13	U	4.4
2	H	228	CYS	4.1
1	R	51	G	4.1
1	R	82	G	3.8
1	R	48	C	3.7
1	R	2	G	3.7
2	H	229	ASP	3.6
1	R	12	C	3.5
1	R	93	C	3.5
2	H	227	SER	3.4
2	H	216	ASN	3.4
1	R	54	C	3.1
1	R	89	U	3.0
1	R	78	G	2.9
2	H	141	LYS	2.8
1	R	84	C	2.8
3	L	49	ILE	2.7
1	R	56	C	2.7
1	R	88	U	2.7
1	R	9	A	2.7
1	R	83	C	2.6
1	R	52	C	2.6
1	R	81	C	2.6
1	R	7	A	2.6
1	R	85	U	2.5
2	H	202	GLY	2.5
2	H	116	GLY	2.4
3	L	37	TYR	2.4
2	H	152	CYS	2.4
1	R	87	U	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	L	99	PHE	2.3
2	H	4	GLU	2.3
2	H	117	GLN	2.2
2	H	226	LYS	2.2
1	R	80	G	2.2
2	H	112	PHE	2.1
3	L	48	LEU	2.1
1	R	11	C	2.1
1	R	43	G	2.1
1	R	8	C	2.0
1	R	10	G	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.