

wwPDB X-ray Structure Validation Summary Report (i)

Oct 5, 2023 - 06:40 AM EDT

PDB ID : 6VQF Title : CRYSTAL STRUCTURE OF RAR-RELATED ORPHAN RECEPTOR C (NHIS-RORGT(244- 487)-L6-SRC1(678-692)) IN COMPLEX WITH AN INVERSE AGONIST Authors : Sack, J.S. Deposited on : 2020-02-05

Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity Mogul		FAILED 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)		
EDS	:	FAILED
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2096 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Nuclear receptor ROR-gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	252	Total 2005	C 1287	N 353	O 351	S 14	0	0	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	244	MET	_	expression tag	UNP P51449
А	245	GLY	_	expression tag	UNP P51449
А	246	SER	-	expression tag	UNP P51449
А	247	SER	-	expression tag	UNP P51449
А	248	HIS	-	expression tag	UNP P51449
А	249	HIS	-	expression tag	UNP P51449
А	250	HIS	-	expression tag	UNP P51449
А	251	HIS	-	expression tag	UNP P51449
А	252	HIS	-	expression tag	UNP P51449
А	253	HIS	-	expression tag	UNP P51449
А	254	SER	-	expression tag	UNP P51449
А	255	SER	-	expression tag	UNP P51449
А	256	GLY	-	expression tag	UNP P51449
А	257	LEU	-	expression tag	UNP P51449
A	258	VAL	-	expression tag	UNP P51449
А	259	PRO	-	expression tag	UNP P51449
А	260	ARG	-	expression tag	UNP P51449
А	261	GLY	-	expression tag	UNP P51449
А	262	SER	-	expression tag	UNP P51449
А	263	HIS	-	expression tag	UNP P51449
А	264	MET	-	expression tag	UNP P51449
А	509	SER	-	expression tag	UNP P51449
А	510	GLY	-	expression tag	UNP P51449
А	511	GLY	-	expression tag	UNP P51449
А	512	SER	-	expression tag	UNP P51449
А	513	GLY	-	expression tag	UNP P51449
А	514	GLY	_	expression tag	UNP P51449

There are 41 discrepancies between the modelled and reference sequences:

Continued on next page...




6	V	QF
U	v	જા

Chain	Residue	Modelled	Actual	Comment	Reference
А	515	LEU	-	expression tag	UNP P51449
А	516	THR	-	expression tag	UNP P51449
А	517	GLU	-	expression tag	UNP P51449
А	518	ARG	-	expression tag	UNP P51449
А	519	HIS	-	expression tag	UNP P51449
А	520	LYS	-	expression tag	UNP P51449
A	521	ILE	-	expression tag	UNP P51449
А	522	LEU	-	expression tag	UNP P51449
А	523	HIS	-	expression tag	UNP P51449
А	524	ARG	-	expression tag	UNP P51449
A	525	LEU	-	expression tag	UNP P51449
А	526	LEU	-	expression tag	UNP P51449
А	527	GLN	-	expression tag	UNP P51449
А	528	GLU	-	expression tag	UNP P51449

Continued from previous page...


• Molecule 2 is (1R,3S,4R)-4-[(3aR,9bR)-9b-[(4-fluorophenyl)sulfonyl]-7-(1,1,1,2,3,3,3-hep tafluoropropan-2-yl)-1,2,3a,4,5,9b-hexahydro-3H-benzo[e]indole-3-carbonyl]-3-methylcyc lohexane-1-carboxylic acid (three-letter code: R7V) (formula: C₃₀H₂₉F₈NO₅S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	А	1	Total 45	-		N 1	-	S 1	0	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	40	Total O 40 40	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	61.51Å 61.51 Å 157.91 Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.00 - 2.00	Depositor
% Data completeness	96.3 (57.00-2.00)	Depositor
(in resolution range)		-
R _{merge}	0.12	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.34 (at 2.25 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.11.7 (3-OCT-2019)	Depositor
R, R_{free}	0.203 , 0.241	Depositor
Wilson B-factor $(Å^2)$	42.0	Xtriage
Anisotropy	0.212	Xtriage
L-test for twinning ²	$ \langle L \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2096	wwPDB-VP
Average B, all atoms $(Å^2)$	44.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.31% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	ype Chain Res Link			Bo	ond leng	ths	Bond angles		
	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	R7V	А	601	-	46,49,49	0.60	2 (4%)	67,80,80	0.90	3 (4%)
3	GOL	А	602	-	$5,\!5,\!5$	0.05	0	$5,\!5,\!5$	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R7V	А	601	-	-	8/51/91/91	0/5/5/5
3	GOL	А	602	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	А	601	R7V	O41-C39	3.04	1.31	1.22
2	А	601	R7V	O40-C39	-2.36	1.22	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	А	601	R7V	O41-C39-C26	-3.85	113.51	122.93
2	А	601	R7V	C9-C8-C7	3.63	118.04	112.69
2	А	601	R7V	O40-C39-C26	3.17	122.50	114.21

There are no chirality outliers.

5 of 10 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	А	601	R7V	C35-C14-C6-C1
3	А	602	GOL	O1-C1-C2-C3
3	А	602	GOL	O1-C1-C2-O2
2	А	601	R7V	C35-C14-C6-C5
2	А	601	R7V	N13-C22-C23-C28

There are no ring outliers.

No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

