



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2024 – 12:07 PM EDT

PDB ID : 8VQK  
Title : YcjN from Escherichia coli  
Authors : Trevino, M.A.; Fernandez, D.; Sharaf, N.G.  
Deposited on : 2024-01-18  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	 2% 81% 10% 9%
1	B	440	 2% 80% 11% 9%

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6501 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative ABC transporter periplasmic-binding protein YcjN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	401	3065	1936	506	609	14	0	0	0
1	B	402	3074	1941	507	612	14	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	431	HIS	-	expression tag	UNP P76042
A	432	HIS	-	expression tag	UNP P76042
A	433	HIS	-	expression tag	UNP P76042
A	434	HIS	-	expression tag	UNP P76042
A	435	HIS	-	expression tag	UNP P76042
A	436	HIS	-	expression tag	UNP P76042
A	437	HIS	-	expression tag	UNP P76042
A	438	HIS	-	expression tag	UNP P76042
A	439	HIS	-	expression tag	UNP P76042
A	440	HIS	-	expression tag	UNP P76042
B	431	HIS	-	expression tag	UNP P76042
B	432	HIS	-	expression tag	UNP P76042
B	433	HIS	-	expression tag	UNP P76042
B	434	HIS	-	expression tag	UNP P76042
B	435	HIS	-	expression tag	UNP P76042
B	436	HIS	-	expression tag	UNP P76042
B	437	HIS	-	expression tag	UNP P76042
B	438	HIS	-	expression tag	UNP P76042
B	439	HIS	-	expression tag	UNP P76042
B	440	HIS	-	expression tag	UNP P76042

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total K 2 2	0	0
5	B	2	Total K 2 2	0	0

- Molecule 6 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	2	Total Cd 2 2	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	155	Total O 155 155	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
7	B	164	Total 164	O 164	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative ABC transporter periplasmic-binding protein YcjN



- Molecule 1: Putative ABC transporter periplasmic-binding protein YcjN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.22Å 63.29Å 74.31Å 110.01° 91.09° 118.66°	Depositor
Resolution (Å)	35.65 – 1.95 35.65 – 1.95	Depositor EDS
% Data completeness (in resolution range)	91.2 (35.65-1.95) 91.3 (35.65-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 1.95Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.215 , 0.264 0.211 , 0.260	Depositor DCC
$R_{free}$ test set	3378 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.8	Xtrriage
Anisotropy	0.360	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 40.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6501	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CD, K, SO4, GOL, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3118	0.55	0/4237
1	B	0.37	0/3127	0.55	0/4249
All	All	0.38	0/6245	0.55	0/8486

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3065	0	3062	31	0
1	B	3074	0	3068	28	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	6	0	8	3	0
4	A	7	0	10	2	0
4	B	14	0	20	5	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	B	2	0	0	0	0
7	A	155	0	0	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	164	0	0	4	0
All	All	6501	0	6168	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:422:LYS:HD2	3:A:502:GOL:H12	1.64	0.79
1:A:92:SER:H	4:A:503:PEG:H22	1.50	0.75
1:B:92:SER:H	4:B:504:PEG:H31	1.55	0.70
1:A:190:LEU:HD11	1:A:237:LEU:HG	1.81	0.63
1:B:422:LYS:HD2	4:B:505:PEG:H31	1.81	0.61
1:A:308:LYS:NZ	7:A:602:HOH:O	2.32	0.61
1:A:397:VAL:HG13	3:A:502:GOL:H11	1.86	0.57
1:B:190:LEU:HD23	1:B:264:ILE:HG13	1.85	0.57
1:B:165:PRO:O	1:B:285:VAL:HG21	2.08	0.54
1:A:94:ASP:HB2	1:A:297:MET:HE1	1.89	0.54
1:A:37:SER:HB3	1:A:42:ARG:HB3	1.90	0.53
1:A:41:GLU:H	1:A:41:GLU:CD	2.12	0.52
1:A:27:THR:OG1	1:A:27:THR:O	2.28	0.51
1:A:165:PRO:O	1:A:285:VAL:HG21	2.09	0.51
1:A:80:SER:HB3	1:A:85:LEU:HD13	1.92	0.51
1:B:180:ASP:HB3	1:B:185:LYS:HB2	1.93	0.51
1:A:234:TYR:O	1:A:238:THR:HG23	2.11	0.51
1:A:130:VAL:HG13	1:A:298:LEU:HD12	1.94	0.50
1:A:333:MET:HA	1:A:333:MET:HE2	1.92	0.50
1:B:355:ASN:O	1:B:359:LYS:HD3	2.13	0.49
1:A:274:ILE:HD13	1:A:279:PRO:HG3	1.94	0.49
1:B:234:TYR:O	1:B:238:THR:HG23	2.13	0.49
1:B:80:SER:HB3	1:B:85:LEU:HG	1.95	0.48
1:B:286:VAL:HG13	1:B:294:VAL:HG23	1.96	0.48
1:A:42:ARG:NH2	1:A:335:SER:OG	2.42	0.48
1:A:238:THR:O	1:A:241:THR:HG22	2.13	0.48
1:A:285:VAL:HG22	1:A:286:VAL:H	1.79	0.48
1:A:93:HIS:CE1	1:A:297:MET:SD	3.07	0.47
1:B:409:LYS:NZ	7:B:614:HOH:O	2.48	0.47
1:B:329:ALA:O	1:B:333:MET:HG2	2.15	0.46
1:A:422:LYS:NZ	7:A:612:HOH:O	2.47	0.46
1:A:191:PRO:HG2	1:A:201:SER:OG	2.16	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:GLU:HB2	1:A:86:PRO:HB3	1.98	0.45
1:A:94:ASP:H	1:A:297:MET:HE3	1.82	0.45
1:A:310:GLU:OE1	1:A:310:GLU:N	2.49	0.45
4:A:503:PEG:H21	4:A:503:PEG:H41	1.68	0.44
1:B:336:PRO:HA	7:B:603:HOH:O	2.17	0.44
1:B:150:ILE:HG12	1:B:264:ILE:CD1	2.48	0.43
1:A:54:GLU:HG2	1:A:60:ILE:O	2.18	0.43
1:B:188:ILE:HD11	1:B:264:ILE:HD11	1.98	0.43
1:B:186:TYR:CD2	1:B:242:MET:HG3	2.53	0.43
1:B:322:MET:HB3	1:B:322:MET:HE2	1.82	0.43
1:A:252:LYS:HD2	1:A:269:ILE:HD12	2.01	0.42
1:B:328:ILE:HG23	1:B:346:VAL:HG22	2.01	0.42
1:B:92:SER:HB2	1:B:297:MET:HE3	2.01	0.42
1:B:85:LEU:HD23	1:B:85:LEU:HA	1.92	0.42
1:B:91:THR:HA	4:B:504:PEG:H31	2.01	0.42
1:B:417:GLN:NE2	7:B:620:HOH:O	2.52	0.42
1:B:422:LYS:HZ2	4:B:505:PEG:H31	1.84	0.42
1:B:340:LEU:HD11	1:B:368:LEU:HB3	2.02	0.42
1:A:44:ALA:O	1:A:48:LYS:HG2	2.20	0.41
1:A:173:ASP:O	1:A:177:LYS:HD3	2.20	0.41
1:A:422:LYS:NZ	3:A:502:GOL:H31	2.35	0.41
1:B:274:ILE:HD13	1:B:279:PRO:HG3	2.03	0.41
1:B:162:LEU:HD21	1:B:177:LYS:HD2	2.03	0.41
1:A:105:ILE:HD13	1:A:139:THR:O	2.21	0.41
1:B:92:SER:H	4:B:504:PEG:C3	2.30	0.40
1:B:257:ASN:OD1	1:B:259:THR:HG23	2.21	0.40
1:A:79:LEU:HD13	1:A:86:PRO:HD3	2.02	0.40
1:B:81:ARG:NE	7:B:621:HOH:O	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/440 (91%)	390 (98%)	9 (2%)	0	100	100
1	B	400/440 (91%)	395 (99%)	4 (1%)	1 (0%)	37	29
All	All	799/880 (91%)	785 (98%)	13 (2%)	1 (0%)	48	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	427	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/372 (90%)	332 (99%)	4 (1%)	67	65
1	B	337/372 (91%)	328 (97%)	9 (3%)	40	31
All	All	673/744 (90%)	660 (98%)	13 (2%)	52	47

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	THR
1	A	32	GLU
1	A	72	TYR
1	A	427	ILE
1	B	27	THR
1	B	82	SER
1	B	133	GLU
1	B	173	ASP
1	B	201	SER
1	B	276	GLU
1	B	294	VAL
1	B	309	THR
1	B	388	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are

no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEG	A	503	-	6,6,6	0.16	0	5,5,5	0.11	0
2	SO4	A	501	-	4,4,4	0.25	0	6,6,6	0.28	0
2	SO4	B	501	-	4,4,4	0.36	0	6,6,6	0.16	0
3	GOL	A	502	-	5,5,5	1.01	0	5,5,5	1.05	0
4	PEG	B	504	-	6,6,6	0.20	0	5,5,5	0.08	0
4	PEG	B	505	-	6,6,6	0.17	0	5,5,5	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	B	504	-	-	3/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	503	-	-	2/4/4/4	-
3	GOL	A	502	-	-	4/4/4/4	-
4	PEG	B	505	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	GOL	C1-C2-C3-O3
4	A	503	PEG	O1-C1-C2-O2
4	B	505	PEG	O2-C3-C4-O4
4	B	504	PEG	C4-C3-O2-C2
3	A	502	GOL	O1-C1-C2-C3
3	A	502	GOL	O2-C2-C3-O3
4	B	504	PEG	O2-C3-C4-O4
4	B	505	PEG	O1-C1-C2-O2
4	A	503	PEG	C4-C3-O2-C2
3	A	502	GOL	O1-C1-C2-O2
4	B	505	PEG	C4-C3-O2-C2
4	B	505	PEG	C1-C2-O2-C3
4	B	504	PEG	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	PEG	2	0
3	A	502	GOL	3	0
4	B	504	PEG	3	0
4	B	505	PEG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	401/440 (91%)	0.48	11 (2%) 56 62	25, 36, 58, 72	0
1	B	402/440 (91%)	0.52	9 (2%) 62 68	26, 37, 58, 74	0
All	All	803/880 (91%)	0.50	20 (2%) 58 65	25, 37, 58, 74	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	THR	3.9
1	B	27	THR	3.8
1	B	40	GLN	2.8
1	B	211	ALA	2.8
1	A	427	ILE	2.6
1	A	182	ALA	2.5
1	B	313	GLU	2.4
1	A	424	ASP	2.3
1	A	426	LEU	2.3
1	A	84	SER	2.3
1	B	424	ASP	2.3
1	B	104	LEU	2.2
1	B	282	VAL	2.2
1	A	216	ALA	2.2
1	B	277	GLY	2.1
1	A	157	LEU	2.1
1	A	162	LEU	2.1
1	A	70	ASP	2.0
1	A	173	ASP	2.0
1	B	41	GLU	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	K	B	506	1/1	0.83	0.12	70,70,70,70	0
4	PEG	A	503	7/7	0.84	0.12	35,38,46,54	0
3	GOL	A	502	6/6	0.85	0.14	40,41,46,47	0
4	PEG	B	504	7/7	0.89	0.11	31,38,47,51	0
4	PEG	B	505	7/7	0.93	0.09	40,47,52,59	0
2	SO4	A	501	5/5	0.96	0.07	37,39,43,44	0
5	K	B	507	1/1	0.96	0.14	35,35,35,35	0
2	SO4	B	501	5/5	0.97	0.07	35,36,41,46	0
5	K	A	504	1/1	0.98	0.16	35,35,35,35	0
5	K	A	505	1/1	0.98	0.04	38,38,38,38	0
6	CD	B	502	1/1	0.99	0.06	57,57,57,57	0
6	CD	B	503	1/1	0.99	0.02	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.