



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 5, 2023 – 06:14 AM EDT

PDB ID : 6VR1
Title : Complex of HLA-A2, a class I MHC, with a p53 peptide
Authors : Wu, D.; Pierce, B.G.; Gallagher, D.T.; Mariuzza, R.A.
Deposited on : 2020-02-06
Resolution : 2.37 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6514 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	2219	1389	402	419	9	0	0	0
1	D	275	2184	1371	388	416	9	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q861F7
A	276	GLY	-	expression tag	UNP Q861F7
A	277	GLY	-	expression tag	UNP Q861F7
A	278	GLY	-	expression tag	UNP Q861F7
A	279	LEU	-	expression tag	UNP Q861F7
A	280	ASN	-	expression tag	UNP Q861F7
A	281	ASP	-	expression tag	UNP Q861F7
A	282	ILE	-	expression tag	UNP Q861F7
A	283	PHE	-	expression tag	UNP Q861F7
A	284	GLU	-	expression tag	UNP Q861F7
A	285	ALA	-	expression tag	UNP Q861F7
A	286	GLN	-	expression tag	UNP Q861F7
A	287	LYS	-	expression tag	UNP Q861F7
A	288	ILE	-	expression tag	UNP Q861F7
A	289	GLU	-	expression tag	UNP Q861F7
A	290	TRP	-	expression tag	UNP Q861F7
A	291	HIS	-	expression tag	UNP Q861F7
A	292	GLU	-	expression tag	UNP Q861F7
D	0	MET	-	initiating methionine	UNP Q861F7
D	276	GLY	-	expression tag	UNP Q861F7
D	277	GLY	-	expression tag	UNP Q861F7
D	278	GLY	-	expression tag	UNP Q861F7
D	279	LEU	-	expression tag	UNP Q861F7
D	280	ASN	-	expression tag	UNP Q861F7
D	281	ASP	-	expression tag	UNP Q861F7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	282	ILE	-	expression tag	UNP Q861F7
D	283	PHE	-	expression tag	UNP Q861F7
D	284	GLU	-	expression tag	UNP Q861F7
D	285	ALA	-	expression tag	UNP Q861F7
D	286	GLN	-	expression tag	UNP Q861F7
D	287	LYS	-	expression tag	UNP Q861F7
D	288	ILE	-	expression tag	UNP Q861F7
D	289	GLU	-	expression tag	UNP Q861F7
D	290	TRP	-	expression tag	UNP Q861F7
D	291	HIS	-	expression tag	UNP Q861F7
D	292	GLU	-	expression tag	UNP Q861F7

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			801	511	137	150	3			
2	E	100	Total	C	N	O	S	0	0	0
			806	516	137	149	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP P61769
E	1	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Cellular tumor antigen p53 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	9	Total	C	N	O	S	0	0	0
			77	45	17	13	2			
3	Q	9	Total	C	N	O	S	0	0	0
			77	45	17	13	2			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	161	Total	O	0	0
			161	161		
4	B	28	Total	O	0	0
			28	28		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	95	Total 95	O 95	0	0
4	E	52	Total 52	O 52	0	0
4	P	10	Total 10	O 10	0	0
4	Q	4	Total 4	O 4	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.33Å 79.73Å 85.85Å 90.00° 101.68° 90.00°	Depositor
Resolution (Å)	20.00 – 2.37	Depositor
% Data completeness (in resolution range)	99.6 (20.00-2.37)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.37Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.217 , 0.274	Depositor
Wilson B-factor (Å ²)	37.4	Xtrriage
Anisotropy	0.620	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	6514	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

There are no ligands in this entry.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.