



Full wwPDB X-ray Structure Validation Report i

Mar 8, 2018 – 11:00 pm GMT

PDB ID : 5VS6
Title : Structure of DUB complex
Authors : Seo, H.-S.; Dhe-Paganon, S.
Deposited on : 2017-05-11
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	6121 (2.30-2.26)
Clashscore	122126	6842 (2.30-2.26)
Ramachandran outliers	120053	6755 (2.30-2.26)
Sidechain outliers	120020	6755 (2.30-2.26)
RSRZ outliers	108989	5992 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)


There are 5 unique types of molecules in this entry. The entry contains 5754 atoms, of which 14 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 7.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total 2691	C 1704	N 457	O 514	S 16	0	0	0
1	B	337	Total 2665	C 1690	N 450	O 509	S 16	0	0	0

- Molecule 2 is N-[3-(4-hydroxy-1-[(3R)-3-phenylbutanoyl]piperidin-4-yl)methyl]-4-oxo-3,4-dihydroquinazolin-7-yl]-3-(4-methylpiperazin-1-yl)propanamide (three-letter code: 9QD) (formula: C₃₂H₄₂N₆O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 42	C 32	N 6	O 4		0	0
2	B	1	Total 42	C 32	N 6	O 4		0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H O 7 2 3 2	0	0
3	B	1	Total C H O 7 2 3 2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C H O 14 3 8 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	151	Total O 151 151	0	0
5	B	135	Total O 135 135	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7



- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.80 Å 68.52 Å 80.31 Å 90.00° 96.17° 90.00°	Depositor
Resolution (Å)	58.01 – 2.27 58.01 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.4 (58.01-2.27) 99.5 (58.01-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.10 (at 2.27 Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R , R_{free}	0.182 , 0.232 0.183 , 0.233	Depositor DCC
R_{free} test set	1867 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	34.8	Xtriage
Anisotropy	0.318	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.2	EDS
L-test for twinning ²	$< L > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5754	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.64 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1119e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT, 9QD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2748	0.54	0/3718
1	B	0.42	0/2721	0.53	0/3682
All	All	0.40	0/5469	0.54	0/7400

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2691	0	2565	47	0
1	B	2665	0	2538	55	0
2	A	42	0	0	0	0
2	B	42	0	0	2	0
3	A	4	3	3	0	0
3	B	4	3	3	0	0
4	B	6	8	8	0	0
5	A	151	0	0	6	0
5	B	135	0	0	3	0
All	All	5740	14	5117	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 10.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:MET:CE	1:A:310:LYS:HB3	1.98	0.93
1:A:245:MET:HE2	1:A:310:LYS:HB3	1.50	0.90
1:A:245:MET:HE1	1:A:307:VAL:HA	1.52	0.89
1:B:245:MET:HE2	1:B:310:LYS:HB3	1.54	0.88
1:B:532:THR:HG23	1:B:534:HIS:H	1.39	0.86
1:B:329:VAL:HG13	1:B:396:LEU:HD11	1.58	0.84
1:B:245:MET:CE	1:B:310:LYS:HB3	2.07	0.83
1:B:492:GLU:OE2	5:B:701:HOH:O	1.97	0.82
1:B:510:CYS:N	5:B:703:HOH:O	2.13	0.81
1:B:439:LYS:HD2	1:B:440:THR:H	1.51	0.74
1:B:329:VAL:CG1	1:B:396:LEU:HD11	2.17	0.74
1:A:245:MET:HE2	1:A:310:LYS:CB	2.18	0.72
1:B:526:GLU:OE2	5:B:702:HOH:O	2.07	0.71
1:B:532:THR:HG23	1:B:534:HIS:N	2.05	0.70
1:A:245:MET:CE	1:A:307:VAL:HA	2.22	0.70
1:A:366:ASP:OD1	5:A:702:HOH:O	2.11	0.69
1:B:439:LYS:HD2	1:B:440:THR:N	2.07	0.68
1:B:245:MET:CE	1:B:307:VAL:HA	2.23	0.68
1:B:245:MET:HE3	1:B:307:VAL:HG13	1.77	0.67
1:B:532:THR:CG2	1:B:534:HIS:HB3	2.26	0.66
1:B:245:MET:HE3	1:B:307:VAL:HA	1.78	0.65
1:A:322:LYS:HG2	1:B:534:HIS:ND1	2.14	0.63
1:B:412:ASP:CB	1:B:419:ILE:HD12	2.28	0.63
1:A:262:ARG:HD2	5:A:740:HOH:O	1.99	0.62
1:A:211:THR:HG22	1:A:213:TYR:CD1	2.36	0.60
1:B:245:MET:HE3	1:B:307:VAL:CA	2.32	0.60
1:A:245:MET:HE3	1:A:310:LYS:HB3	1.81	0.59
1:B:411:TYR:CE1	1:B:413:PRO:HA	2.37	0.59
1:A:366:ASP:OD2	5:A:703:HOH:O	2.17	0.59
1:A:448:TYR:HB3	1:A:518:TYR:HB3	1.83	0.59
1:A:424:ARG:HB2	1:A:500:GLY:HA2	1.84	0.59
1:B:468:TYR:CE2	1:B:479:LYS:HG3	2.37	0.59
1:A:228:LEU:HD23	1:A:299:LEU:HD13	1.85	0.59
1:B:279:LEU:HG	1:B:283:PHE:HE2	1.69	0.58
1:A:279:LEU:HD13	1:A:283:PHE:HE2	1.70	0.56
1:B:252:SER:HA	1:B:258:LEU:HD23	1.86	0.56
1:B:329:VAL:HG13	1:B:396:LEU:CD1	2.33	0.56
1:A:279:LEU:HD13	1:A:283:PHE:CE2	2.41	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:245:MET:HE1	1:B:310:LYS:HB3	1.89	0.55
1:A:412:ASP:OD2	1:A:415:THR:HG23	2.08	0.54
1:A:353:SER:O	1:A:363:SER:OG	2.23	0.53
2:B:601:9QD:C29	2:B:601:9QD:O3	2.57	0.52
1:B:252:SER:HB2	1:B:543:ARG:HH21	1.75	0.52
1:B:439:LYS:HE3	1:B:440:THR:O	2.10	0.52
1:A:509:HIS:N	5:A:717:HOH:O	2.43	0.52
1:B:536:ILE:HG22	1:B:541:VAL:HG23	1.92	0.51
1:B:536:ILE:CG2	1:B:541:VAL:HG23	2.42	0.50
1:B:246:PRO:O	1:B:310:LYS:HE3	2.12	0.49
1:A:307:VAL:O	1:A:311:MET:HG3	2.11	0.49
1:A:251:ASP:OD2	1:A:254:LYS:N	2.45	0.49
1:B:448:TYR:HB3	1:B:518:TYR:HB3	1.95	0.49
1:A:245:MET:HE2	1:A:310:LYS:CG	2.43	0.49
1:B:245:MET:HE1	1:B:307:VAL:HA	1.95	0.48
1:A:240:LYS:O	1:A:244:MET:HG2	2.14	0.48
1:A:259:ALA:HA	1:A:262:ARG:NH1	2.29	0.47
1:B:470:ASN:HD22	1:B:470:ASN:H	1.60	0.47
1:A:371:GLU:OE2	1:A:391:LYS:HD2	2.14	0.47
1:B:468:TYR:CZ	1:B:479:LYS:HG3	2.49	0.47
1:A:429:GLU:HG2	1:A:430:GLN:HG2	1.96	0.47
1:B:468:TYR:HD2	1:B:488:CYS:HG	1.62	0.47
1:B:353:SER:O	1:B:357:LYS:HG3	2.14	0.47
1:A:209:LYS:N	5:A:712:HOH:O	2.47	0.47
1:A:231:THR:HG21	1:A:517:VAL:HG21	1.97	0.47
1:A:279:LEU:CD1	1:A:283:PHE:HE2	2.28	0.46
1:A:536:ILE:HG22	1:A:541:VAL:HG23	1.96	0.46
1:B:350:ILE:HD12	1:B:402:LEU:HD11	1.97	0.46
1:A:350:ILE:HD12	1:A:402:LEU:HD11	1.98	0.46
1:A:347:TYR:OH	1:A:398:LEU:HD23	2.15	0.45
1:B:289:ASP:O	1:B:293:GLN:HG2	2.17	0.45
1:A:354:ILE:HG22	1:A:425:PHE:CE1	2.53	0.44
1:A:454:LEU:HB2	1:A:515:MET:HB3	1.98	0.44
1:B:223:CYS:HB3	1:B:465:TYR:CZ	2.52	0.44
1:A:523:LYS:NZ	5:A:701:HOH:O	2.01	0.44
1:B:328:MET:SD	1:B:393:VAL:HG21	2.57	0.44
1:A:456:HIS:HA	1:A:464:HIS:O	2.18	0.44
1:B:256:VAL:N	1:B:257:PRO:CD	2.81	0.44
1:A:225:MET:HG3	1:A:299:LEU:HD11	2.00	0.44
1:B:411:TYR:HE1	1:B:413:PRO:HA	1.83	0.44
1:A:245:MET:HE2	1:A:310:LYS:HG2	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:337:VAL:HG11	1:A:384:HIS:ND1	2.34	0.43
1:B:307:VAL:O	1:B:311:MET:HG3	2.19	0.43
1:B:318:GLY:C	1:B:321:PRO:HD2	2.38	0.43
1:B:403:HIS:CD2	1:B:517:VAL:HG22	2.53	0.43
1:B:248:GLU:CD	1:B:248:GLU:H	2.21	0.43
1:B:318:GLY:O	1:B:321:PRO:HD2	2.18	0.43
1:B:492:GLU:HA	1:B:496:HIS:ND1	2.34	0.43
1:B:354:ILE:HG22	1:B:425:PHE:CE1	2.54	0.43
1:B:265:TYR:HD1	1:B:536:ILE:HD13	1.84	0.42
1:B:532:THR:HG23	1:B:533:ASP:N	2.34	0.42
1:A:371:GLU:O	1:A:390:GLU:HA	2.20	0.42
1:A:269:HIS:NE2	1:A:533:ASP:OD2	2.52	0.42
1:A:223:CYS:HB3	1:A:465:TYR:CZ	2.55	0.42
1:B:252:SER:HA	1:B:258:LEU:CD2	2.49	0.42
1:A:283:PHE:CE1	1:A:285:TRP:HB2	2.55	0.42
1:A:245:MET:HE3	1:A:307:VAL:O	2.20	0.41
1:A:245:MET:HE1	1:A:307:VAL:CA	2.37	0.41
1:A:534:HIS:HB3	1:B:322:LYS:HE3	2.02	0.41
1:B:297:GLN:HE22	2:B:601:9QD:C30	2.32	0.41
1:A:370:VAL:HG13	1:A:390:GLU:HB2	2.03	0.41
1:B:245:MET:HE3	1:B:307:VAL:CG1	2.48	0.40
1:B:325:ARG:O	1:B:399:PRO:HB3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	335/353 (95%)	319 (95%)	14 (4%)	2 (1%)	27 31
1	B	333/353 (94%)	323 (97%)	10 (3%)	0	100 100
All	All	668/706 (95%)	642 (96%)	24 (4%)	2 (0%)	43 52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	414	GLN
1	A	416	ASP

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/320 (90%)	277 (96%)	11 (4%)	36 48
1	B	284/320 (89%)	276 (97%)	8 (3%)	47 61
All	All	572/640 (89%)	553 (97%)	19 (3%)	41 54

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	SER
1	A	299	LEU
1	A	322	LYS
1	A	438	GLN
1	A	469	LEU
1	A	470	ASN
1	A	487	ARG
1	A	525	SER
1	A	531	VAL
1	A	533	ASP
1	A	542	GLU
1	B	287	THR
1	B	299	LEU
1	B	315	CYS
1	B	386	LEU
1	B	409	PHE
1	B	438	GLN
1	B	470	ASN
1	B	533	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	9QD	A	601	-	45,46,46	1.26	4 (8%)	55,65,65	1.37	5 (9%)
3	ACT	A	602	-	1,3,3	1.16	0	0,3,3	0.00	-
2	9QD	B	601	-	45,46,46	1.06	5 (11%)	55,65,65	1.88	16 (29%)
3	ACT	B	602	-	1,3,3	1.46	0	0,3,3	0.00	-
4	GOL	B	603	-	5,5,5	0.39	0	5,5,5	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9QD	A	601	-	-	0/26/48/48	0/5/5/5
3	ACT	A	602	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9QD	B	601	-	-	0/26/48/48	0/5/5/5
3	ACT	B	602	-	-	0/0/0/0	0/0/0/0
4	GOL	B	603	-	-	0/4/4/4	0/0/0/0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	9QD	C20-N3	-2.91	1.35	1.41
2	B	601	9QD	C20-N3	-2.89	1.35	1.41
2	B	601	9QD	C30-C31	-2.11	1.36	1.41
2	B	601	9QD	C17-N2	2.28	1.35	1.30
2	A	601	9QD	C-N	2.28	1.41	1.38
2	B	601	9QD	C-C31	2.41	1.45	1.41
2	B	601	9QD	C31-C18	3.66	1.48	1.41
2	A	601	9QD	C31-C18	3.75	1.49	1.41
2	A	601	9QD	C-C31	4.52	1.48	1.41

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	9QD	C29-C30-C31	-3.48	116.68	121.57
2	B	601	9QD	C31-C18-N2	-3.44	117.14	122.40
2	B	601	9QD	C20-C19-C18	-3.01	117.21	120.50
2	A	601	9QD	C20-N3-C21	-2.69	122.73	127.48
2	B	601	9QD	C25-C24-N4	-2.59	105.54	110.64
2	B	601	9QD	C27-C28-N4	-2.35	106.00	110.64
2	B	601	9QD	O3-C21-C22	-2.33	117.68	122.00
2	A	601	9QD	C29-C30-C31	-2.22	118.46	121.57
2	B	601	9QD	C30-C31-C	-2.09	118.55	121.61
2	B	601	9QD	C20-N3-C21	-2.05	123.87	127.48
2	B	601	9QD	O3-C21-N3	2.00	127.34	123.66
2	A	601	9QD	C4-N1-C15	2.09	116.56	112.61
2	B	601	9QD	C27-N5-C25	2.20	112.67	109.53
2	B	601	9QD	C31-C-N	2.26	117.69	116.15
2	B	601	9QD	C1-N-C17	2.51	122.91	118.76
2	A	601	9QD	C19-C18-N2	2.71	120.95	117.91
2	B	601	9QD	C4-N1-C15	2.74	117.79	112.61
2	B	601	9QD	C30-C31-C18	3.26	122.49	117.57
2	A	601	9QD	C17-N2-C18	5.42	121.62	116.72
2	B	601	9QD	C19-C18-N2	5.59	124.17	117.91
2	B	601	9QD	C17-N2-C18	6.11	122.23	116.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	9QD	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	339/353 (96%)	-0.11	3 (0%)	84	87	27, 45, 74, 102
1	B	337/353 (95%)	-0.17	1 (0%)	93	95	26, 43, 73, 121
All	All	676/706 (95%)	-0.14	4 (0%)	89	92	26, 44, 73, 121

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	413	PRO	3.2
1	B	414	GLN	3.1
1	A	208	LYS	2.4
1	A	416	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	603	6/6	0.68	0.28	70,85,99,101	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	B	602	4/4	0.87	0.14	57,61,73,73	0
3	ACT	A	602	4/4	0.88	0.13	64,69,82,82	0
2	9QD	B	601	42/42	0.95	0.14	22,35,87,91	0
2	9QD	A	601	42/42	0.96	0.12	24,36,78,79	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.