



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2023 – 03:06 AM EDT

PDB ID : 6VSF
Title : Mycobacterium tuberculosis dihydrofolate reductase in complex with 4-(3,4-dihydro-2H-benzo[b][1,4]dioxepin-7-yl)-4-oxobutanoic acid(fragment 16)
Authors : Ribeiro, J.A.; Dias, M.V.B.
Deposited on : 2020-02-11
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 2960 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydrofolate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	159	Total	C	N	O	S	0	0	0
			1244	783	228	228	5			
1	B	159	Total	C	N	O	S	0	0	0
			1244	783	228	228	5			

- Molecule 2 is 4-(3,4-dihydro-2H-1,5-benzodioxepin-7-yl)-4-oxobutanoic acid (three-letter code: RKY) (formula: C₁₃H₁₄O₅) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total	C	O	0	0
			18	13	5		
2	B	1	Total	C	O	0	0
			18	13	5		
2	B	1	Total	C	O	0	0
			18	13	5		

- Molecule 3 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Co 1 1	0	0
3	B	1	Total Co 1 1	0	0

- Molecule 4 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 48 21 7 17 3	0	0
4	B	1	Total C N O P 48 21 7 17 3	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		
6	B	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	170	Total	O	0	0
			170	170		
7	B	123	Total	O	0	0
			123	123		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.56Å 70.90Å 71.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.77 – 2.01	Depositor
% Data completeness (in resolution range)	99.3 (46.77-2.01)	Depositor
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.01Å)	Xtrriage
Refinement program	PHENIX 1.14	Depositor
R, R_{free}	0.163 , 0.211	Depositor
Wilson B-factor (Å ²)	20.3	Xtrriage
Anisotropy	0.183	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.026 for -h,l,k	Xtrriage
Total number of atoms	2960	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RKY	B	201	-	19,19,19	2.77	8 (42%)	25,25,25	2.64	11 (44%)
4	NAP	B	204	-	45,52,52	2.66	11 (24%)	56,80,80	1.95	12 (21%)
6	PO4	A	205	-	4,4,4	0.95	0	6,6,6	0.45	0
2	RKY	A	201	-	19,19,19	2.42	7 (36%)	25,25,25	1.19	4 (16%)
5	GOL	A	204	-	5,5,5	0.55	0	5,5,5	0.80	0
5	GOL	B	205	-	5,5,5	0.75	0	5,5,5	1.48	1 (20%)
2	RKY	B	202	-	19,19,19	2.50	7 (36%)	25,25,25	1.18	4 (16%)
4	NAP	A	203	-	45,52,52	2.56	14 (31%)	56,80,80	1.86	13 (23%)
6	PO4	B	207	-	4,4,4	0.74	0	6,6,6	0.62	0
6	PO4	B	206	-	4,4,4	0.91	0	6,6,6	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RKY	B	201	-	-	2/9/17/17	0/2/2/2
4	NAP	B	204	-	-	7/31/67/67	0/5/5/5
2	RKY	A	201	-	-	0/9/17/17	0/2/2/2
5	GOL	A	204	-	-	0/4/4/4	-
5	GOL	B	205	-	-	0/4/4/4	-
2	RKY	B	202	-	-	2/9/17/17	0/2/2/2
4	NAP	A	203	-	-	4/31/67/67	0/5/5/5

All (47) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	204	NAP	P2B-O2B	11.20	1.80	1.59
4	A	203	NAP	P2B-O2B	10.24	1.78	1.59
4	B	204	NAP	C4N-C3N	6.39	1.50	1.39
4	A	203	NAP	C4N-C3N	6.07	1.49	1.39
2	B	202	RKY	C05-C06	5.89	1.59	1.51
4	B	204	NAP	C5N-C4N	5.39	1.50	1.38
2	A	201	RKY	C05-C06	5.38	1.59	1.51
2	B	201	RKY	O16-C17	5.33	1.48	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	RKY	C08-C06	5.22	1.57	1.49
2	B	201	RKY	C05-C06	5.08	1.58	1.51
4	A	203	NAP	C5N-C4N	4.84	1.49	1.38
4	B	204	NAP	C3N-C7N	-4.64	1.43	1.50
4	B	204	NAP	PN-O5D	4.37	1.77	1.59
2	A	201	RKY	O16-C17	4.32	1.46	1.38
2	B	202	RKY	O16-C17	4.26	1.46	1.38
2	A	201	RKY	C08-C06	3.95	1.55	1.49
2	B	202	RKY	C08-C06	3.73	1.55	1.49
4	A	203	NAP	O4B-C1B	3.62	1.46	1.41
2	B	201	RKY	O07-C06	-3.45	1.16	1.22
4	A	203	NAP	C2N-N1N	3.41	1.39	1.35
2	B	201	RKY	O16-C15	-3.41	1.39	1.44
2	A	201	RKY	O16-C15	-3.39	1.39	1.44
4	A	203	NAP	O4D-C1D	3.34	1.45	1.41
2	B	202	RKY	O16-C15	-3.30	1.39	1.44
4	B	204	NAP	O4B-C1B	3.22	1.45	1.41
4	B	204	NAP	O2B-C2B	-3.19	1.32	1.44
4	A	203	NAP	PN-O5D	3.16	1.72	1.59
2	B	202	RKY	O07-C06	-3.07	1.17	1.22
4	A	203	NAP	C2A-N3A	2.93	1.36	1.32
2	B	202	RKY	O12-C13	-2.90	1.39	1.44
2	A	201	RKY	O07-C06	-2.89	1.17	1.22
4	A	203	NAP	O2B-C2B	-2.82	1.33	1.44
4	B	204	NAP	C2N-N1N	2.68	1.38	1.35
2	B	201	RKY	O12-C11	2.65	1.43	1.38
4	B	204	NAP	C6N-N1N	2.62	1.41	1.35
2	A	201	RKY	O12-C11	2.60	1.43	1.38
4	A	203	NAP	C6N-N1N	2.59	1.41	1.35
4	A	203	NAP	C3B-C2B	2.48	1.58	1.52
2	B	202	RKY	O12-C11	2.47	1.43	1.38
2	B	201	RKY	O12-C13	-2.42	1.40	1.44
4	B	204	NAP	C2N-C3N	-2.39	1.35	1.39
2	A	201	RKY	O12-C13	-2.21	1.41	1.44
4	A	203	NAP	C2A-N1A	2.07	1.37	1.33
4	A	203	NAP	O4B-C4B	-2.06	1.40	1.45
4	B	204	NAP	C2A-N1A	2.03	1.37	1.33
4	A	203	NAP	C6N-C5N	-2.03	1.34	1.38
2	B	201	RKY	C04-C02	2.02	1.55	1.50

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	204	NAP	C5N-C4N-C3N	-7.04	112.01	120.34
4	A	203	NAP	C5N-C4N-C3N	-6.49	112.67	120.34
2	B	201	RKY	C18-C08-C06	-5.42	108.45	120.23
4	B	204	NAP	O7N-C7N-C3N	5.10	125.73	119.63
2	B	201	RKY	C09-C08-C06	5.00	131.90	120.61
2	B	201	RKY	O12-C11-C17	-4.82	117.98	123.17
4	B	204	NAP	PN-O3-PA	-4.14	118.63	132.83
4	A	203	NAP	O7N-C7N-C3N	4.11	124.55	119.63
2	B	201	RKY	O16-C15-C14	-4.09	107.63	112.72
4	A	203	NAP	O7N-C7N-N7N	-3.89	117.05	122.58
4	B	204	NAP	C2N-C3N-C4N	3.78	122.55	118.26
4	A	203	NAP	PN-O3-PA	-3.73	120.02	132.83
2	B	201	RKY	C05-C06-C08	3.67	124.22	119.13
2	B	201	RKY	O16-C17-C11	3.62	127.06	123.17
4	B	204	NAP	O2B-P2B-O1X	-3.43	96.16	109.39
4	A	203	NAP	C2N-C3N-C4N	3.28	121.97	118.26
2	B	201	RKY	O07-C06-C08	-3.22	116.18	120.74
4	A	203	NAP	O2B-P2B-O1X	-3.07	97.52	109.39
4	B	204	NAP	C3N-C7N-N7N	-2.87	114.31	117.75
2	B	201	RKY	O01-C02-C04	2.73	122.80	114.03
2	B	201	RKY	O16-C17-C18	-2.73	113.00	116.95
4	A	203	NAP	O3X-P2B-O2X	2.66	117.81	107.64
2	B	201	RKY	C04-C05-C06	-2.63	109.57	112.76
2	B	202	RKY	O16-C17-C11	-2.58	120.39	123.17
4	A	203	NAP	C3B-C2B-C1B	-2.56	98.07	102.89
2	A	201	RKY	C15-O16-C17	-2.56	112.48	116.03
4	B	204	NAP	O3X-P2B-O2X	2.55	117.40	107.64
5	B	205	GOL	C3-C2-C1	-2.47	102.08	111.70
4	B	204	NAP	PA-O5B-C5B	-2.42	107.47	121.68
4	A	203	NAP	C6N-C5N-C4N	2.40	122.92	119.44
4	B	204	NAP	C6N-N1N-C2N	-2.36	119.82	121.97
2	B	202	RKY	O12-C13-C14	2.30	115.58	112.72
4	A	203	NAP	PA-O5B-C5B	-2.28	108.30	121.68
4	B	204	NAP	C2B-C3B-C4B	-2.25	97.10	101.99
2	A	201	RKY	C15-C14-C13	-2.21	109.63	113.54
2	A	201	RKY	O01-C02-C04	2.20	121.10	114.03
4	B	204	NAP	C3B-C2B-C1B	-2.16	98.83	102.89
4	A	203	NAP	O4D-C1D-C2D	-2.15	103.79	106.93
2	A	201	RKY	C04-C05-C06	-2.13	110.19	112.76
4	B	204	NAP	C6N-C5N-C4N	2.11	122.51	119.44
2	B	202	RKY	C05-C04-C02	-2.08	109.12	113.60
4	A	203	NAP	C6N-N1N-C2N	-2.06	120.10	121.97
2	B	202	RKY	O01-C02-C04	2.05	120.60	114.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	201	RKY	C09-C10-C11	-2.03	116.19	120.06
4	A	203	NAP	O3X-P2B-O2B	-2.01	97.00	105.99

There are no chirality outliers.



All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	203	NAP	PA-O3-PN-O5D
4	A	203	NAP	O4D-C1D-N1N-C6N
4	A	203	NAP	C2D-C1D-N1N-C2N
4	B	204	NAP	PA-O3-PN-O5D
4	B	204	NAP	O4D-C1D-N1N-C2N
4	B	204	NAP	O4D-C1D-N1N-C6N
2	B	202	RKY	O01-C02-C04-C05
2	B	202	RKY	O03-C02-C04-C05
2	B	201	RKY	O01-C02-C04-C05
2	B	201	RKY	O03-C02-C04-C05
4	B	204	NAP	C5B-O5B-PA-O3
4	B	204	NAP	C2D-C1D-N1N-C2N
4	B	204	NAP	C2D-C1D-N1N-C6N
4	A	203	NAP	C5B-O5B-PA-O1A
4	B	204	NAP	C5B-O5B-PA-O1A


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







4.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.