



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 28, 2026 – 04:36 PM UTC

PDB ID : 8VS8 / pdb\_00008vs8  
Title : Crystal structure of ADI-19425 Fab in complex with anti-idiotypic 1D3 Fab  
Authors : Kher, G.; Homad, L.J.; McGuire, A.T.; Pancera, M.  
Deposited on : 2024-01-23  
Resolution : 2.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>10%</div> <div>75%</div> <div>14%</div> <div>11%</div> </div>
1	E	227	<div> <div>2%</div> <div>77%</div> <div>13%</div> <div>9%</div> </div>
1	I	227	<div> <div>2%</div> <div>44%</div> <div>9%</div> <div>47%</div> </div>
1	M	227	<div> <div>5%</div> <div>83%</div> <div>9%</div> <div>7%</div> </div>
1	Q	227	<div> <div>3%</div> <div>48%</div> <div>7%</div> <div>45%</div> </div>



*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	U	227	
1	Y	227	
1	c	227	
2	B	218	
2	F	218	
2	J	218	
2	N	218	
2	R	218	
2	V	218	
2	Z	218	
2	d	218	
3	C	214	
3	G	214	
3	K	214	
3	O	214	
3	S	214	
3	W	214	
3	a	214	
3	e	214	
4	D	226	
4	H	226	
4	L	226	
4	P	226	
4	T	226	
4	X	226	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	b	226	
4	f	226	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 45030 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADI-19425 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	0	0
			1512	958	253	293	8			
1	E	206	Total	C	N	O	S	0	0	0
			1547	979	259	301	8			
1	I	121	Total	C	N	O	S	0	0	0
			920	578	156	180	6			
1	M	210	Total	C	N	O	S	0	0	0
			1576	996	264	308	8			
1	Q	125	Total	C	N	O	S	0	0	0
			940	588	159	187	6			
1	U	178	Total	C	N	O	S	0	0	0
			1347	850	227	262	8			
1	Y	121	Total	C	N	O	S	0	0	0
			920	578	156	180	6			
1	c	98	Total	C	N	O	S	0	0	0
			748	474	126	143	5			

- Molecule 2 is a protein called ADI-19425 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	213	Total	C	N	O	S	1	0	0
			1568	983	261	320	4			
2	F	214	Total	C	N	O	S	1	0	0
			1587	996	263	324	4			
2	J	110	Total	C	N	O	S	0	0	0
			802	502	133	165	2			
2	N	214	Total	C	N	O	S	1	0	0
			1581	993	260	324	4			
2	R	110	Total	C	N	O	S	0	0	0
			806	504	134	166	2			
2	V	201	Total	C	N	O	S	0	0	0
			1484	933	245	302	4			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Z	108	Total	C	N	O	S	0	0	0
			792	496	131	163	2			
2	d	81	Total	C	N	O	S	0	0	0
			602	379	97	124	2			

- Molecule 3 is a protein called 1D3 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	214	Total	C	N	O	S	0	0	0
			1649	1027	279	338	5			
3	G	214	Total	C	N	O	S	0	0	0
			1649	1027	280	337	5			
3	K	212	Total	C	N	O	S	0	0	0
			1641	1024	279	334	4			
3	O	214	Total	C	N	O	S	0	0	0
			1657	1032	281	339	5			
3	S	214	Total	C	N	O	S	0	0	0
			1647	1026	277	339	5			
3	W	200	Total	C	N	O	S	0	0	0
			1549	967	261	317	4			
3	a	214	Total	C	N	O	S	0	0	0
			1645	1023	278	339	5			
3	e	214	Total	C	N	O	S	0	0	0
			1635	1018	275	337	5			

- Molecule 4 is a protein called 1D3 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	221	Total	C	N	O	S	9	0	0
			1660	1053	274	327	6			
4	H	220	Total	C	N	O	S	0	0	0
			1648	1047	270	325	6			
4	L	218	Total	C	N	O	S	2	0	0
			1632	1036	271	319	6			
4	P	217	Total	C	N	O	S	3	0	0
			1637	1041	270	320	6			
4	T	214	Total	C	N	O	S	2	0	0
			1616	1029	266	315	6			
4	X	216	Total	C	N	O	S	0	0	0
			1622	1031	266	319	6			
4	b	218	Total	C	N	O	S	5	0	0
			1639	1041	270	322	6			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	f	215	Total	C	N	O	S	5	0	0
			1594	1014	260	314	6			

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	9	Total	O	0	0
			9	9		
5	B	7	Total	O	0	0
			7	7		
5	C	5	Total	O	0	0
			5	5		
5	D	13	Total	O	0	0
			13	13		
5	E	6	Total	O	0	0
			6	6		
5	F	6	Total	O	0	0
			6	6		
5	G	10	Total	O	0	0
			10	10		
5	H	6	Total	O	0	0
			6	6		
5	I	5	Total	O	0	0
			5	5		
5	J	1	Total	O	0	0
			1	1		
5	K	3	Total	O	0	0
			3	3		
5	L	6	Total	O	0	0
			6	6		
5	M	14	Total	O	0	0
			14	14		
5	N	5	Total	O	0	0
			5	5		
5	O	13	Total	O	0	0
			13	13		
5	P	2	Total	O	0	0
			2	2		
5	Q	6	Total	O	0	0
			6	6		
5	S	5	Total	O	0	0
			5	5		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	T	8	Total 8	O 8	0	0
5	U	5	Total 5	O 5	0	0
5	V	4	Total 4	O 4	0	0
5	W	7	Total 7	O 7	0	0
5	X	6	Total 6	O 6	0	0
5	Y	3	Total 3	O 3	0	0
5	Z	1	Total 1	O 1	0	0
5	a	7	Total 7	O 7	0	0
5	b	10	Total 10	O 10	0	0
5	c	2	Total 2	O 2	0	0
5	e	2	Total 2	O 2	0	0
5	f	1	Total 1	O 1	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADI-19425 Heavy Chain




- Molecule 1: ADI-19425 Heavy Chain




- Molecule 1: ADI-19425 Heavy Chain




- Molecule 1: ADI-19425 Heavy Chain








• Molecule 1: ADI-19425 Heavy Chain





• Molecule 1: ADI-19425 Heavy Chain




• Molecule 1: ADI-19425 Heavy Chain




• Molecule 1: ADI-19425 Heavy Chain







• Molecule 2: ADI-19425 Light Chain




• Molecule 2: ADI-19425 Light Chain




• Molecule 2: ADI-19425 Light Chain




• Molecule 2: ADI-19425 Light Chain












• Molecule 2: ADI-19425 Light Chain




• Molecule 2: ADI-19425 Light Chain




• Molecule 2: ADI-19425 Light Chain





• Molecule 2: ADI-19425 Light Chain






### • Molecule 3: 1D3 Light Chain





### • Molecule 3: 1D3 Light Chain




### • Molecule 3: 1D3 Light Chain




### • Molecule 3: 1D3 Light Chain





### • Molecule 3: 1D3 Light Chain









• Molecule 3: 1D3 Light Chain




• Molecule 3: 1D3 Light Chain




• Molecule 3: 1D3 Light Chain





• Molecule 4: 1D3 Heavy Chain









• Molecule 4: 1D3 Heavy Chain





• Molecule 4: 1D3 Heavy Chain




• Molecule 4: 1D3 Heavy Chain




• Molecule 4: 1D3 Heavy Chain





• Molecule 4: 1D3 Heavy Chain







• Molecule 4: 1D3 Heavy Chain



• Molecule 4: 1D3 Heavy Chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.07Å 149.98Å 164.43Å 83.98° 89.90° 74.61°	Depositor
Resolution (Å)	49.33 – 2.67 49.33 – 2.67	Depositor EDS
% Data completeness (in resolution range)	92.5 (49.33-2.67) 92.5 (49.33-2.67)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.17 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, $R_{free}$	0.231 , 0.273 0.233 , 0.275	Depositor DCC
$R_{free}$ test set	11502 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.6	Xtriage
Anisotropy	0.250	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 36.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	45030	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.12	0/1548	0.31	0/2105
1	E	0.23	0/1584	0.38	0/2153
1	I	0.09	0/941	0.28	0/1272
1	M	0.12	0/1614	0.31	0/2195
1	Q	0.10	0/961	0.29	0/1301
1	U	0.23	0/1377	0.41	0/1863
1	Y	0.09	0/941	0.27	0/1272
1	c	0.21	0/757	0.41	0/1010
2	B	0.17	0/1609	0.35	0/2201
2	F	0.11	0/1621	0.29	0/2218
2	J	0.09	0/822	0.29	0/1121
2	N	0.12	0/1615	0.33	0/2211
2	R	0.10	0/825	0.28	0/1123
2	V	0.19	0/1512	0.36	0/2064
2	Z	0.09	0/811	0.26	0/1106
2	d	0.19	0/610	0.36	0/820
3	C	0.18	0/1685	0.37	0/2288
3	G	0.11	0/1685	0.32	0/2288
3	K	0.10	0/1677	0.32	0/2277
3	O	0.11	0/1693	0.30	0/2297
3	S	0.11	0/1683	0.30	0/2286
3	W	0.11	0/1582	0.33	0/2145
3	a	0.15	0/1681	0.36	0/2285
3	e	0.18	0/1671	0.35	0/2273
4	D	0.18	0/1697	0.38	1/2314 (0.0%)
4	H	0.12	0/1684	0.30	0/2296
4	L	0.12	0/1667	0.30	0/2272
4	P	0.19	0/1673	0.38	1/2280 (0.0%)
4	T	0.20	0/1652	0.43	0/2253
4	X	0.12	0/1656	0.31	0/2257
4	b	0.12	0/1675	0.32	0/2284
4	f	0.19	0/1637	0.40	1/2237 (0.0%)

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
All	All	0.15	0/45846	0.34	3/62367 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	f	100(A)	ASP	N-CA-C	-5.87	106.10	113.20
4	D	134	GLY	N-CA-C	-5.24	106.80	114.61
4	P	100(D)	PHE	CA-CB-CG	5.04	118.84	113.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1512	0	1459	21	0
1	E	1547	0	1501	16	0
1	I	920	0	876	13	0
1	M	1576	0	1528	14	0
1	Q	940	0	887	7	0
1	U	1347	0	1301	14	0
1	Y	920	0	876	13	0
1	c	748	0	705	9	0
2	B	1568	0	1502	19	0
2	F	1587	0	1532	22	0
2	J	802	0	761	7	0
2	N	1581	0	1521	21	0
2	R	806	0	766	14	1
2	V	1484	0	1425	18	0
2	Z	792	0	747	15	0
2	d	602	0	555	12	0
3	C	1649	0	1577	17	0
3	G	1649	0	1579	23	0
3	K	1641	0	1583	16	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	O	1657	0	1594	25	0
3	S	1647	0	1572	20	0
3	W	1549	0	1488	26	0
3	a	1645	0	1561	39	0
3	e	1635	0	1546	29	1
4	D	1660	0	1607	22	0
4	H	1648	0	1590	25	0
4	L	1632	0	1580	16	0
4	P	1637	0	1586	15	0
4	T	1616	0	1563	17	0
4	X	1622	0	1559	25	0
4	b	1639	0	1583	8	0
4	f	1594	0	1512	23	0
5	A	9	0	0	0	0
5	B	7	0	0	0	0
5	C	5	0	0	0	0
5	D	13	0	0	0	0
5	E	6	0	0	0	0
5	F	6	0	0	0	0
5	G	10	0	0	0	0
5	H	6	0	0	0	0
5	I	5	0	0	0	0
5	J	1	0	0	0	0
5	K	3	0	0	0	0
5	L	6	0	0	0	0
5	M	14	0	0	0	0
5	N	5	0	0	0	0
5	O	13	0	0	0	0
5	P	2	0	0	0	0
5	Q	6	0	0	0	0
5	S	5	0	0	0	0
5	T	8	0	0	0	0
5	U	5	0	0	0	0
5	V	4	0	0	0	0
5	W	7	0	0	0	0
5	X	6	0	0	0	0
5	Y	3	0	0	0	0
5	Z	1	0	0	0	0
5	a	7	0	0	0	0
5	b	10	0	0	0	0
5	c	2	0	0	0	0
5	e	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	f	1	0	0	0	0
All	All	45030	0	43022	539	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 539 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:Z:5:THR:HB	2:Z:24:THR:OG1	1.40	1.21
3:a:13:VAL:CG1	3:a:78:VAL:HG21	1.77	1.14
3:a:13:VAL:HG11	3:a:78:VAL:HG21	1.13	1.07
3:G:11:LEU:HD21	3:G:19:VAL:CG2	1.85	1.07
4:H:123:PRO:HB3	4:H:211:VAL:HG22	1.36	1.02

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:R:54:ARG:NH2	3:e:187:GLU:O[1_454]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/227 (87%)	183 (93%)	12 (6%)	2 (1%)	12	28
1	E	200/227 (88%)	193 (96%)	6 (3%)	1 (0%)	24	45
1	I	119/227 (52%)	114 (96%)	4 (3%)	1 (1%)	16	34
1	M	204/227 (90%)	197 (97%)	6 (3%)	1 (0%)	24	45
1	Q	123/227 (54%)	119 (97%)	2 (2%)	2 (2%)	7	18

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	U	170/227 (75%)	155 (91%)	13 (8%)	2 (1%)	10	24
1	Y	119/227 (52%)	113 (95%)	4 (3%)	2 (2%)	7	17
1	c	78/227 (34%)	72 (92%)	6 (8%)	0	100	100
2	B	211/218 (97%)	197 (93%)	14 (7%)	0	100	100
2	F	212/218 (97%)	200 (94%)	12 (6%)	0	100	100
2	J	108/218 (50%)	99 (92%)	8 (7%)	1 (1%)	14	31
2	N	212/218 (97%)	198 (93%)	14 (7%)	0	100	100
2	R	106/218 (49%)	96 (91%)	10 (9%)	0	100	100
2	V	191/218 (88%)	176 (92%)	14 (7%)	1 (0%)	24	45
2	Z	104/218 (48%)	96 (92%)	8 (8%)	0	100	100
2	d	63/218 (29%)	55 (87%)	8 (13%)	0	100	100
3	C	212/214 (99%)	201 (95%)	11 (5%)	0	100	100
3	G	212/214 (99%)	204 (96%)	8 (4%)	0	100	100
3	K	210/214 (98%)	199 (95%)	10 (5%)	1 (0%)	24	45
3	O	212/214 (99%)	202 (95%)	10 (5%)	0	100	100
3	S	212/214 (99%)	199 (94%)	13 (6%)	0	100	100
3	W	194/214 (91%)	178 (92%)	15 (8%)	1 (0%)	24	45
3	a	212/214 (99%)	198 (93%)	14 (7%)	0	100	100
3	e	212/214 (99%)	198 (93%)	13 (6%)	1 (0%)	24	45
4	D	219/226 (97%)	210 (96%)	8 (4%)	1 (0%)	24	45
4	H	216/226 (96%)	205 (95%)	11 (5%)	0	100	100
4	L	214/226 (95%)	205 (96%)	8 (4%)	1 (0%)	24	45
4	P	213/226 (94%)	203 (95%)	10 (5%)	0	100	100
4	T	210/226 (93%)	197 (94%)	12 (6%)	1 (0%)	24	45
4	X	208/226 (92%)	198 (95%)	8 (4%)	2 (1%)	12	28
4	b	214/226 (95%)	200 (94%)	14 (6%)	0	100	100
4	f	211/226 (93%)	200 (95%)	11 (5%)	0	100	100
All	All	5798/7080 (82%)	5460 (94%)	317 (6%)	21 (0%)	30	51

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Y	100(A)	GLY

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	Q	115	SER
1	U	113	SER
1	A	100(B)	SER
1	A	114	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/191 (87%)	166 (99%)	1 (1%)	78	90
1	E	173/191 (91%)	167 (96%)	6 (4%)	32	58
1	I	99/191 (52%)	99 (100%)	0	100	100
1	M	177/191 (93%)	176 (99%)	1 (1%)	78	90
1	Q	101/191 (53%)	100 (99%)	1 (1%)	68	84
1	U	151/191 (79%)	150 (99%)	1 (1%)	76	89
1	Y	99/191 (52%)	97 (98%)	2 (2%)	48	74
1	c	80/191 (42%)	79 (99%)	1 (1%)	61	81
2	B	174/181 (96%)	173 (99%)	1 (1%)	78	90
2	F	177/181 (98%)	176 (99%)	1 (1%)	78	90
2	J	88/181 (49%)	87 (99%)	1 (1%)	65	83
2	N	176/181 (97%)	176 (100%)	0	100	100
2	R	89/181 (49%)	89 (100%)	0	100	100
2	V	165/181 (91%)	164 (99%)	1 (1%)	78	90
2	Z	87/181 (48%)	86 (99%)	1 (1%)	65	83
2	d	66/181 (36%)	66 (100%)	0	100	100
3	C	189/191 (99%)	188 (100%)	1 (0%)	81	91
3	G	189/191 (99%)	188 (100%)	1 (0%)	81	91
3	K	189/191 (99%)	187 (99%)	2 (1%)	65	83
3	O	191/191 (100%)	189 (99%)	2 (1%)	68	84

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	S	189/191 (99%)	187 (99%)	2 (1%)	65	83
3	W	180/191 (94%)	177 (98%)	3 (2%)	53	77
3	a	188/191 (98%)	184 (98%)	4 (2%)	47	73
3	e	186/191 (97%)	184 (99%)	2 (1%)	65	83
4	D	181/186 (97%)	180 (99%)	1 (1%)	78	90
4	H	179/186 (96%)	178 (99%)	1 (1%)	78	90
4	L	177/186 (95%)	175 (99%)	2 (1%)	65	83
4	P	178/186 (96%)	178 (100%)	0	100	100
4	T	175/186 (94%)	173 (99%)	2 (1%)	65	83
4	X	176/186 (95%)	175 (99%)	1 (1%)	78	90
4	b	178/186 (96%)	177 (99%)	1 (1%)	78	90
4	f	171/186 (92%)	170 (99%)	1 (1%)	78	90
All	All	4985/5992 (83%)	4941 (99%)	44 (1%)	70	86

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	W	21	PHE
3	a	1	ASP
3	W	33	ILE
1	Y	13	LYS
3	a	155	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
3	W	92	ASN
4	b	5	GLN
3	W	198	HIS
3	a	92	ASN
2	d	38	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PCA	H	1	4	7,8,9	0.59	0	9,10,12	1.02	1 (11%)
4	PCA	L	1	4	7,8,9	0.45	0	9,10,12	0.79	1 (11%)
4	PCA	X	1	4	7,8,9	0.61	0	9,10,12	0.82	1 (11%)
2	PCA	N	1	2	7,8,9	0.48	0	9,10,12	1.24	1 (11%)
2	PCA	F	1	2	7,8,9	0.47	0	9,10,12	0.88	1 (11%)
2	PCA	V	1	2	7,8,9	0.46	0	9,10,12	1.27	1 (11%)
4	PCA	T	1	4	7,8,9	0.51	0	9,10,12	0.98	1 (11%)
4	PCA	b	1	4	7,8,9	0.60	0	9,10,12	1.19	1 (11%)
4	PCA	D	1	4	7,8,9	0.55	0	9,10,12	0.87	1 (11%)
4	PCA	P	1	4	7,8,9	0.52	0	9,10,12	0.91	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PCA	H	1	4	-	0/0/11/13	0/1/1/1
4	PCA	L	1	4	-	0/0/11/13	0/1/1/1
4	PCA	X	1	4	-	0/0/11/13	0/1/1/1
2	PCA	N	1	2	-	0/0/11/13	0/1/1/1
2	PCA	F	1	2	-	0/0/11/13	0/1/1/1
2	PCA	V	1	2	-	0/0/11/13	0/1/1/1
4	PCA	T	1	4	-	0/0/11/13	0/1/1/1
4	PCA	b	1	4	-	0/0/11/13	0/1/1/1
4	PCA	D	1	4	-	0/0/11/13	0/1/1/1
4	PCA	P	1	4	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	V	1	PCA	O-C-CA	-3.70	115.26	124.77
2	N	1	PCA	O-C-CA	-3.28	116.33	124.77
4	b	1	PCA	O-C-CA	-3.20	116.53	124.77
4	T	1	PCA	O-C-CA	-2.86	117.41	124.77
4	H	1	PCA	O-C-CA	-2.44	118.49	124.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	203/227 (89%)	0.47	22 (10%) 11 9	24, 40, 88, 103	0
1	E	206/227 (90%)	0.22	4 (1%) 66 63	24, 41, 75, 88	0
1	I	121/227 (53%)	0.21	4 (3%) 49 44	29, 45, 63, 83	0
1	M	210/227 (92%)	0.32	12 (5%) 29 25	26, 43, 73, 87	0
1	Q	125/227 (55%)	0.30	6 (4%) 35 31	29, 48, 68, 85	0
1	U	178/227 (78%)	0.80	25 (14%) 6 5	34, 52, 93, 109	0
1	Y	121/227 (53%)	0.90	12 (9%) 13 10	34, 56, 77, 89	0
1	c	98/227 (43%)	2.47	61 (62%) 0 0	83, 102, 112, 124	0
2	B	213/218 (97%)	0.48	15 (7%) 22 19	26, 49, 95, 103	1 (0%)
2	F	213/218 (97%)	0.21	4 (1%) 66 63	26, 47, 74, 95	1 (0%)
2	J	110/218 (50%)	0.75	9 (8%) 17 14	43, 58, 75, 83	0
2	N	213/218 (97%)	0.31	7 (3%) 49 44	26, 46, 73, 99	1 (0%)
2	R	110/218 (50%)	0.83	9 (8%) 17 14	45, 61, 84, 145	0
2	V	200/218 (91%)	1.03	46 (23%) 2 1	29, 58, 117, 131	0
2	Z	108/218 (49%)	1.65	31 (28%) 1 1	60, 82, 98, 107	0
2	d	81/218 (37%)	2.46	47 (58%) 0 0	81, 104, 116, 119	0
3	C	214/214 (100%)	0.31	8 (3%) 45 40	27, 48, 70, 105	0
3	G	214/214 (100%)	0.25	1 (0%) 87 86	29, 49, 70, 91	0
3	K	212/214 (99%)	0.27	6 (2%) 55 51	27, 49, 67, 79	0
3	O	214/214 (100%)	0.57	13 (6%) 27 23	29, 50, 81, 102	0
3	S	214/214 (100%)	0.64	14 (6%) 25 21	35, 57, 78, 110	0
3	W	200/214 (93%)	1.33	45 (22%) 2 1	36, 66, 98, 109	0
3	a	214/214 (100%)	0.77	16 (7%) 20 17	33, 61, 89, 110	0
3	e	214/214 (100%)	1.17	37 (17%) 4 3	49, 73, 92, 102	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
4	D	220/226 (97%)	0.39	19 (8%) 16 13	27, 43, 87, 141	5 (2%)
4	H	219/226 (96%)	0.35	18 (8%) 17 14	26, 43, 88, 137	0
4	L	217/226 (96%)	0.64	19 (8%) 15 13	29, 47, 82, 180	1 (0%)
4	P	216/226 (95%)	0.40	12 (5%) 30 26	26, 43, 79, 141	2 (0%)
4	T	213/226 (94%)	0.52	17 (7%) 18 15	29, 43, 76, 174	1 (0%)
4	X	215/226 (95%)	0.92	29 (13%) 7 6	29, 51, 101, 187	0
4	b	217/226 (96%)	0.61	21 (9%) 13 11	30, 45, 77, 171	2 (0%)
4	f	215/226 (95%)	1.27	32 (14%) 5 4	46, 70, 104, 155	5 (2%)
All	All	5938/7080 (83%)	0.67	621 (10%) 11 9	24, 52, 97, 187	19 (0%)

The worst 5 of 621 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	X	100(B)	GLY	10.1
4	L	100(C)	PRO	9.9
4	b	52(A)	SER	9.4
4	f	100(B)	GLY	9.1
4	b	100(D)	PHE	8.6

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PCA	b	1	8/9	0.50	0.33	20,20,20,20	0
4	PCA	T	1	8/9	0.56	0.30	20,20,20,20	0
4	PCA	L	1	8/9	0.67	0.26	20,20,20,20	0
2	PCA	F	1	8/9	0.68	0.27	20,20,20,20	0
2	PCA	V	1	8/9	0.71	0.36	20,20,20,20	0
2	PCA	N	1	8/9	0.79	0.32	20,20,20,20	0
4	PCA	P	1	8/9	0.85	0.23	20,20,20,20	0
4	PCA	H	1	8/9	0.88	0.22	20,20,20,20	0
4	PCA	X	1	8/9	0.89	0.19	20,20,20,20	0
4	PCA	D	1	8/9	0.90	0.16	20,20,20,20	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.